



IJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: ETPAM-2018 **Issue:** Conference **Month of publication:** March 2018

DOI:

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

Strongly Unique Best Co-approximation in Linear 2-Normed Spaces

Sumitha Solson¹, R. S. Karunya²

¹II PG-Mathematics, ²Assistant Professor, Department of Mathematics, St. Joseph's College of Arts and Science for Women, Hosur, Periyar university, salem.

Abstract: This paper deals with some fundamental properties of the set of strongly unique best co-approximation in a linear 2-normed space.

Keywords: Linear 2-normed space, best co- approximation and strongly unique best Co-approximation.

I. INTRODUCTION

The problem of best co-approximation was first introduced by Franchetti and Furi [1] to study some characteristic properties of real Hilbert spaces and was followed up by Papini and Singer [2].

This theory is largely concerned with the questions of existence, uniqueness and characterization of best co-approximation. Newman and Shapiro [3] studied the problems of strongly unique best approximation in the space of continuous functions under supremum norm. The notion of strongly unique best co-approximation in the context of linear 2-normed spaces is introduced in this paper, it provides some important definitions and results that are required and deals with some fundamental properties of the set of strongly unique best co-approximation with respect to 2-norm.

II. PRELIMINARIES

A. Definition 1.2.1[4]

Let G be a non-empty subset of a linear 2-normed space X . An element $g_0 \in G$ is called a best co-approximation to $x \in X$ from G if for every $g \in G$, $\|g - g_0, k\| \leq \|x - g, k\|$, for every $k \in X \setminus [G, x]$, where $[G, x]$ represents a linear space spanned by elements of G and x .

B. Definition 1.2.2[4]

Let G be a non-empty subset of a linear 2-normed space X . An element $g_0 \in G$ is called a strongly unique best co-approximation to $x \in X$ from G , if there exists a constant $t > 0$ such that for every $g \in G$, $\|g - g_0, k\| \leq \|x - g, k\| - t\|x - g_0, k\|$, for every $k \in X \setminus [G, x]$.

The set of all elements of strongly unique best co-approximations to $x \in X$ from G is denoted by $T_G(x)$.

The subset G is called an existence set if $T_G(x)$ contains at least one element for every $x \in X$.

G is called a uniqueness set if $T_G(x)$ contains at most one element for every $x \in X$. G is called an existence and uniqueness set if $T_G(x)$ contains exactly one element for every $x \in X$.

C. Definition 1.2.3[4]

A set $K \subset X$ is convex if $\lambda x + (1 - \lambda)y \in K$ whenever $x, y \in K$ and $\lambda \in [0, 1]$.

A convex combination of x_1, \dots, x_m is a sum of the form $\sum_{i=1}^n a_i x_i$, Where $\sum_{i=1}^n a_i = 1$, n is a positive integer, and all the a_i are non-negative.

If K is convex and $x_1, \dots, x_n \in K$, each $a_i \geq 0$, and $\sum_{i=1}^n a_i = 1$, then $\sum_{i=1}^n a_i x_i \in K$.

This can be proved easily using induction on n .

D. Definition 1.2.4[4]

A set $K \subset V$ is called closed if the limit of every sequence $\{x_n\} \subset K$, which converges in V , belongs to K . i.e., $\{x_n\} \subset K$ and $x_n \rightarrow x$ in V then $x \in K$.

E. Definition 1.2.5[4]

A function f defined on some set X with real or complex values is called **bounded**, if the set of its values is bounded. In other words, there exists a real number M such that

$$|f(x)| \leq M$$

For all x in X . A function that is not bounded is said to be **unbounded**.

III. SOME FUNDAMENTAL PROPERTIES OF $T_G(x)$

Some basic properties of strongly unique best co-approximation are obtained in the following Theorems.

A. Theorem 1.3.1[2]

For any points $a, b \in X$ and any $\alpha \in \mathbb{R}$,

$$\| \alpha a, b \| = \| \alpha a, b + \alpha a \|$$

B. Theorem 1.3.2[5]

Let G be a subset of a linear 2-normed space X and $x \in X$. Then the following statements hold.

- 1) $T_G(x)$ is closed if G is closed.
- 2) $T_G(x)$ is convex if G is convex.
- 3) $T_G(x)$ is bounded.

a) Proof:

(i) Let G be closed.

Let $\{g_m\}$ be a sequence in $T_G(x)$ such that $g_m \rightarrow \bar{g}$.

To prove that $T_G(x)$ is closed, it is enough to prove that $\bar{g} \in T_G(x)$.

Since G is closed, $\{g_m\} \in G$ and $g_m \rightarrow \bar{g}$, we have $\bar{g} \in G$.

Since $\{g_m\} \in T_G(x)$, we have

$$\| g - g_m, k \| \leq \| x - g, k \| - t \| x - g_m, k \|, \text{ for every } k \in X \setminus [G, x] \text{ and for some } t > 0$$

$$\Rightarrow \| g - g_m + \bar{g} - \bar{g}, k \| \leq \| x - g, k \| - t \| x - g_m, k \|$$

$$\Rightarrow \| g - \bar{g}, k \| - \| g_m - \bar{g}, k \| \leq \| x - g, k \| - t \| x - g_m, k \|, \text{ for every } g \in G$$

Since $g_m \rightarrow \bar{g}$, $g_m - \bar{g} \rightarrow 0$(1)

So $\| g_m - \bar{g}, k \| \rightarrow 0$, as 0 and k are linearly dependent.

Therefore, it follows from (1) that

$$\| g - \bar{g}, k \| \leq \| x - g, k \| - t \| x - \bar{g}, k \|, \text{ for every } g \in G \text{ and for some } t > 0.$$

Thus $\bar{g} \in T_G(x)$

Hence $T_G(x)$ is closed.

(ii). Let G be convex, $g_1, g_2 \in T_G(x)$ and $\alpha \in (0, 1)$.

To prove that $\alpha g_1 + (1 - \alpha)g_2 \in T_G(x)$,

Let $k \in X \setminus [G, x]$. Then

$$\begin{aligned} \| g - (\alpha g_1 + (1 - \alpha)g_2), k \| &= \| \alpha(g - g_1) + (1 - \alpha)(g - g_2), k \| \\ &\leq \alpha \| g - g_1, k \| + (1 - \alpha) \| g - g_2, k \| \\ &\leq \alpha \| x - g, k \| - \alpha t \| x - g_1, k \| \\ &\quad + (1 - \alpha) \| x - g, k \| - (1 - \alpha)t \| x - g_2, k \|, \text{ for every } g \in G \text{ and for some } t > 0. \\ &= \| x - g, k \| - t (\alpha \| x - g_1, k \| + (1 - \alpha) \| x - g_2, k \|) \\ &\leq \| x - g, k \| - t (\alpha \| x - g_1, k \| + (1 - \alpha) \| x - g_2, k \|) \\ &= \| x - g, k \| - t \| x - (\alpha g_1 + (1 - \alpha)g_2), k \|. \end{aligned}$$

Thus $\alpha g_1 + (1 - \alpha)g_2 \in T_G(x)$.

Hence $T_G(x)$ is convex.

(iii). To prove that $T_G(x)$ is bounded, it is enough to prove for arbitrary $g_0, \bar{g}_0 \in T_G(x)$ that $\| g_0 - \bar{g}_0, k \| < c$ for some $c > 0$, since $\| g_0 - \bar{g}_0, k \| < c$ implies that $\sup_{g_0, \bar{g}_0 \in T_G(x)} \| g_0 - \bar{g}_0, k \|$ is finite and hence the diameter of $T_G(x)$ is finite.

Let $g_0, \bar{g}_0 \in T_G(x)$. Then there exists a constant $t > 0$ such that for every

$g \in G$ and $k \in X \setminus [G, x]$,

$$\begin{aligned} \| g - g_0, k \| &\leq \| x - g, k \| - t \| x - g_0, k \| \text{ and} \\ \| g - \bar{g}_0, k \| &\leq \| x - g, k \| - t \| x - \bar{g}_0, k \|. \end{aligned}$$

Now,

$$\begin{aligned} \| x - g_0, k \| &\leq \| x - g, k \| + \| g - g_0, k \| \\ &\leq 2 \| x - g, k \| - t \| x - g_0, k \|. \end{aligned}$$

Hence $\|x - g_0, k\| \leq \frac{2}{1+t} d$,

where $d = \inf_{g \in G} \|x - g, k\|$

Similarly, $\|x - \bar{g}_0, k\| \leq \frac{2}{1+t} d$

Therefore, it follows that

$$\|g_0 - \bar{g}_0, k\| \leq \|g_0 - x, k\| + \|x - \bar{g}_0, k\| \\ \leq \frac{4}{1+t} d = c$$

Hence $T_G(x)$ is bounded

Hence the proof

C. Theorem 1.3.3[2]

Let G be a subset of a linear 2-normed space X , $x \in X$ and $K \in X \setminus [G, x]$. Then the following statements are equivalent for every $y \in [k]$.

- 1) $g_0 \in T_G(x)$.
 - 2) $g_0 \in T_G(x + y)$.
 - 3) $g_0 \in T_G(x - y)$.
 - 4) $g_0 + y \in T_G(x + y)$.
 - 5) $g_0 + y \in T_G(x - y)$.
- a) Proof: The proof follows immediately by using Theorem 1.3.1

D. Theorem 1.3.2

Let G be a subspace of a linear 2-normed space X and $x \in X$. Then the following statements hold.

- 1) $T_G(x + g) = T_G(x) + g$, for every $g \in G$.
- 2) $T_G(\alpha x) = \alpha T_G(x)$, for every $\alpha \in \mathbb{R}$.

a) Proof

(i). Let \bar{g} be an arbitrary but fixed element of G .

Let $g_0 \in T_G(x)$. It is clear that $g_0 + \bar{g} \in T_G(x) + \bar{g}$.

To prove that $T_G(x) + \bar{g} \subseteq T_G(x + \bar{g})$,

it is enough to prove that $g_0 + \bar{g} \in T_G(x + \bar{g})$.

Now,

$$\|g + \bar{g} - g_0 - \bar{g}, k\| \leq \|x - g, k\| - t \|x - g_0, k\|, \text{ for all } g \in G \text{ and for some } t > 0.$$

$$\Rightarrow \|g + \bar{g} - (g_0 + \bar{g}), k\| \leq \|x + \bar{g} - (g + \bar{g}), k\| - t \|x + \bar{g} - (g_0 + \bar{g}), k\|$$

for all $g \in G$

$$\Rightarrow g_0 + \bar{g} \in T_G(x + \bar{g}), \text{ since } g - \bar{g} \in G.$$

Conversely, let $g_0 + \bar{g} \in T_G(x + \bar{g})$.

To prove that $T_G(x + \bar{g}) \subseteq T_G(x) + \bar{g}$, it is enough to prove that $g_0 \in T_G(x)$.

Now,

$$\|g - g_0, k\| = \|g + \bar{g} - (g_0 + \bar{g}), k\|$$

$$\leq \|x + \bar{g} - (g + \bar{g}), k\| \leq -t \|x + \bar{g} - (g_0 + \bar{g}), k\|, \text{ for all } g \in G \text{ and for some } t > 0.$$

$$\Rightarrow g_0 \in T_G(x), \text{ thus the result follows.}$$

(ii). The proof is similar to that of (i).

Hence the proof.

E. Proposition 1.3.4[2]

Let G be a subset of a linear 2-normed space X , $x \in X$ and

$k \in X \setminus [G, x]$. If $g_0 \in T_G(x)$, then there exists a constant $t > 0$ such that for all

$g \in G$,

$$\|x - g_0, k\| \leq 2\|x - g, k\| - t\|x - g_0, k\|$$

Proof:

The proof is trivial.

F. Theorem: 1.3.5 [2]

Let G be a subspace of a linear 2-normed space X , $x \in X$ and

$K \in X \setminus \{G, x\}$. Then $g_0 \in T_G(x) \Leftrightarrow g_0 \in T_G(\alpha^m x + (1 - \alpha^m)g_0)$, for all $\alpha \in \mathbb{R}$ and $m = 0, 1, 2, \dots$.

Proof:

Claim: $g_0 \in T_G(x) \Leftrightarrow g_0 \in T_G(\alpha x + (1 - \alpha)g_0)$, for every $\alpha \in \mathbb{R}$.

Let $g_0 \in T_G(x)$. Then

$\|g - g_0, k\| \leq \|x - g, k\| - t \|x - g_0, k\|$, for all $g \in G$ and for some $t > 0$.

$\Rightarrow \| \alpha g - \alpha g_0, k \| \leq \| \alpha x - \alpha g, k \| - t \| \alpha x - \alpha g_0, k \|$, for all $g \in G$

$\Rightarrow \| \alpha (\frac{(\alpha-1)g_0+g}{\alpha}) - \alpha g_0, k \| \leq \| \alpha x - \alpha (\frac{(\alpha-1)g_0+g}{\alpha}), k \| - t \| \alpha x - \alpha g_0, k \|$, for all $g \in G$ and $\alpha \neq 0$,

Since $(\frac{(\alpha-1)g_0+g}{\alpha}) \in G$

$\Rightarrow \|g - g_0, k\| \leq \| \alpha x + (1 - \alpha)g_0 - g, k \| - t \| \alpha x + (1 - \alpha)g_0 - g_0, k \|$

$\Rightarrow g_0 \in T_G(\alpha x + (1 - \alpha)g_0)$, when $\alpha \neq 0$.

If $\alpha = 0$, then it is clear that $g_0 \in T_G(\alpha x + (1 - \alpha)g_0)$.

The converse is obvious by taking $\alpha = 1$.

Hence the claim is true.

By repeated application of the claim the result follows.

IV. CONCLUSION

Hence we proved that linear 2-normed space are closed, bounded and convex.

Therefore, strongly unique best co-approximation in linear 2-normed space is also closed.

Here also we proved some fundamental properties.

REFERENCE

- [1] Franchetti .C and Furi.M, some characteristic properties of real Hilbert spaces, Rev. Roumaine Math. Pures. Appl., 17(1972), 1045-1048.
- [2] Papini.P.L and I.Singer, Best approximation in normed linear spaces, Mh.Math.,88(1979), 27-44.
- [3] Newman D.J and H.S.Shapiro, some theorems on Chebyshev approximation, Duke Math.J., 30(1963), 673-684.
- [4] Cho.Y.J, "Theory of 2-inner product spaces", Nova science Publications, New York, 1994.
- [5] Geetha S.Rao and R.Saravanan, Strongly unique best coapproximation, kyungpook Math. J., 43(2003), 000-000.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)