iJRASET
International Journal For Research in Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: ETPAM-2018 **Issue:** Conference **Month of publication:** March 2018

DOI:

www.ijraset.com

Call: 08813907089 | E-mail ID: ijraset@gmail.com

ε-Best Approximation and E- Orthogonality

R.S. Karunya
Assistant Professor, Department of Mathematics, St. Joseph's College of Arts and Science for Women, Hosur.

Abstract: The purpose of this paper is to study the concept of ε-Best approximation and ε-orthogonality. I discussed their properties and noted that are similar to the properties of best approximation.
Keywords: ε-Best approximation, normed linear spaces, proximinal, ε-orthogonality, convex.

I. INTRODUCTION

The theory of best approximation is an important topic in functional analysis. It is a very extensive field which has various applications

What do we mean by "***Best approximation"*** in normed linear spaces?

To explain this, let X be a normed linear space, and let G be a nonempty subset of X. An element $g_0 \in G$ is called a best approximation to x from G if g_0 is closest to x from among all the elements of G.

That is, $\| x-g_0 \| \leq \| x-g \|$ for all $g \in G$.

The set of all such elements $g_0 \in G$ are called a ***best approximation*** to $x \in X$ is denoted by $P_G(x)$.

If $P_G(x)$ contains at least one element, then the subset G is called a ***proximinal*** set. If each element $x \in X$ has a unique best approximation in G, then G is called a ***Chebyshev*** set of X.

The theory of approximation is mainly concerned with the following fundamental questions.

1) *(Existence of best approximation)* Which subsets are proximinal?
2) *(Uniqueness of best approximation)* Which subsets are Chebyshev?
3) *(Characterization of best approximation)* How to recognize when a given $y \in G$ is a best approximation to x or not?
4) *(Error of approximation)* How to compute the error of approximation d(x, G)?
5) (*Computation of best approximation*) How to describe some useful algorithms for actually computing best approximation?
6) (*Continuity of best approximation*) How does the set of all best approximation vary as a function of x or (G)?

A. Definition 1.1[1]

Let G be a nonempty subset of a real normed linear space E and let an element $f \in E$ be given. The problem of ***best approximation*** is to determine an element $g_f \in G$ such that

$$\| f- g_f \| = \inf_{g \in G} \|f - g\|$$

such an element is called a *best approximation* to f from G ,*and*

$$d(f, G) = \inf_{g \in G} \|f - g\| \text{ is called the } \textit{minimal deviation} \text{ off from G.}$$

The set of all elements $g_0 \in G$ that are called best approximation to $x \in X$ is

$$P_G(x) = \{ g_0 \in G: \| x - g_0 \| \leq \| x - g \| \text{ for all } g \in G \}$$

Hence P_G defines a mapping from X into the power set of G is called the ***metric projection*** onto G, (other names nearest point mapping, proximity map)

B. Remark 1.2[1]

The set $P_G(x)$ of all *best approximation* to $x \in X$ can be written as

$$P_G(x) = \{g_0 \in G: \| x-g_0 \| = d(x, G)\}$$

©IJRASET (UGC Approved Journal): All Rights are Reserved

C. Definition 1.3.[3]

A set S, in a linear space is ***convex*** .if $s_1, s_2 \in S$ implies that

$$\lambda_1 s_1 + \lambda_2 s_2 \in S$$

If λ_1 and λ_2 are non negative and $\lambda_1 + \lambda_2 = 1$

If S is empty or consists of one point, then it is clearly *convex*

D. Definition 1.4[1]

If $P_G(x)$ contains at least one element, then the subset G is called a *proximinal set.*

In other words, if $P_G(x) \neq \varphi$ then G is called a *proximinal set*

The term *proximinal set* (is a combination of proximity and maximal)

E. Definition 1.5[1] (Quasi-Orthogonal Set)

Let X be a normed linear space, and G a nonempty subset of X. Then we say that G is *quasi-orthogonal set* if $G \perp_B \hat{G}$, that is $g \perp_B \hat{G}$ for every $g \in G$.

where $\hat{G} = \{x \in X ; \| x \| = d(x, G)\} = \{x \in X : x \perp_B G\}$.

F. Remark 1.6[1]

In a Hilbert space, any closed subspace is quasi-orthogonal.

Proof:

Let H be a Hilbert space and G a closed subspace of H.

Then $\hat{G} = G^{\perp} = \{y \in H : < x, y > = 0, \text{ for all } x \in G\}$. Then $G \perp \hat{G}$.

Therefore G is quasi-orthogonal subspace of H.

G. Definition 1.7[2]

Let X be a normed linear space and G be a subset of X, and $\varepsilon > 0$. A point $g_0 \in G$ is said to be ε-best approximation for $x \in X$ if and only if

$\| x - g_0 \| \leq \| x - g \| + \varepsilon$ for all $g \in G$

H. Remark 1.8[2]

For $x \in X$, the set of all ε-Best approximation of x in G is denoted by

$P_G(x, \varepsilon)$, in other words,

$P_G(x, \varepsilon) = \{g_0 \in G: \| x - g_0 \| \leq \| x - g \| + \varepsilon \text{ for all } g \in G\}$.

I. Theorem 1.9[2]

Let G be a subspace of a normed linear space X. Then $P_G(x, \varepsilon)$ is bounded.

Proof:

Let $g_1, g_2 \in P_G(x, \varepsilon)$, then $\| x - g_1 \| \leq \| x - g \| + \varepsilon$ for all $g \in G$, and

$$\| x - g_2 \| \leq \| x - g \| + \varepsilon \text{ for all } g \in G$$

Now, $\| g_1 - g_2 \| = \| g_1 - x + x - g_2 \| \leq \| x - g_1 \| + \| x - g_2 \|$

$$\leq \| x - g \| + \varepsilon + \| x - g \| + \varepsilon = 2 \| x - g \| + 2\varepsilon = k,$$

so we have $\| g_1 - g_2 \| \leq k$ where $k = 2d(x, G) + 2\varepsilon$.

Therefore, $P_G(x, \varepsilon)$ is bounded.

Hence the proof

©IJRASET (UGC Approved Journal): All Rights are Reserved

J. Theorem 1.10[2]

Let G be a subspace of normed linear space X, and $x \in X$. Then $P_G(x, \varepsilon)$ is convex.

Proof:

Let $g_1, g_2 \in P_G(x, \varepsilon)$, and $0 \le \lambda \le 1$, then $\| x - g_1 \| \le \| x - g \| + \varepsilon$ for all $g \in G$, and $\| x - g_2 \| \le \| x - g \| + \varepsilon$ for all $g \in G$

Now,
$$\begin{aligned} \| x - (\lambda g_1 + (1-\lambda) g_2) \| &= \| x - \lambda g_1 - g_2 + \lambda g_2 \| \\ &= \| x - \lambda g_1 - g_2 + \lambda g_2 + \lambda x - \lambda x \| \\ &= \| \lambda(x - g_1) + (1-\lambda)(x - g_2) \| \\ &\le \| x - g_1 \| + (1-\lambda) \| x - g_2 \| \\ &\le \lambda(\| x - g \| + \varepsilon) + (1 - \lambda)(\| x - g \| + \varepsilon) \\ &= \| x - g \| + \varepsilon. \end{aligned}$$

Thus, $\lambda g_1 + (1-\lambda) g_2 \in P_G(x, \varepsilon)$.

Hence $P_G(x, \varepsilon)$ is convex.

Hence the proof

K. Definition 1.11.[2] (ε-orthogonality)

Let X be a normed linear space, $\varepsilon > 0$, and $x, y \in X$. We call x is ***ε- orthogonal*** to y and is denoted by $x \perp_\varepsilon y$ if and only if $\| x + \alpha y \| + \varepsilon \ge \| x \|$ for all scaler α with $| \alpha | \le 1$

For subsets G_1, G_2 of X, $G_1 \perp_\varepsilon G_2$ if and only if, $g_1 \perp_\varepsilon g_2$ for all $g_1 \in G_1, g_2 \in G_2$.

L. Theorem: 1.12[2]

Let X be a normed linear space, G be a subspace of X, and $\varepsilon > 0$.Then for all $x \in X$, $g_0 \in P_G(x, \varepsilon)$ if and only if $(x - g_0) \perp_\varepsilon G$.

Proof:

(=>) Suppose $g_0 \in P_G(x, \varepsilon)$. Put $g_1 = g_0 - \alpha g$ for $g \in G$ and $| \alpha | \le 1$.

Since $g_0 \in P_G(x, \varepsilon)$ and $g_1 \in G$ so, then, $\| x - g_0 \| \le \| x - g_1 \| + \varepsilon$, then

$\| x - g_0 \| \le \| x - (g_0 - \alpha g) \| + \varepsilon$, and this implies that

$\| x - g_0 \| \le \| (x - g_0) + \alpha g \| + \varepsilon$.

Therefore, $(x - g_0) \perp_\varepsilon G$.

(<=) Let $(x - g_0) \perp_\varepsilon G$, then for all α with $| \alpha | \le 1$ and $g_1 \in G$

we have,

$\| x - g_0 \| \le \| x - g_0 + \alpha g_1 \| + \varepsilon$

For any $g \in G$ by putting $g_1 = g_0 - g$ and $\alpha = 1$, the last inequality implies,

$\| x - g_0 \| \le \| x - g \| + \varepsilon$

Therefore, $g_0 \in P_G(x, \varepsilon)$

Hence the proof

M. Notation 1.13

Let X be a normed linear space, and G a subspace of X, and for $\varepsilon > 0$, let

$P_G^{-1}(0, \varepsilon) = \{x \in X : \| x \| \le \| x - g \| + \varepsilon \text{ for all } g \in G\} = \{x \in X : x \perp_\varepsilon G\}$

Then, $\hat{G}_\varepsilon = \{x \in X : x \perp_\varepsilon G\}$.

N. Lemma 1.14[2]

Let G be a subspace of a normed linear space X. Then for all $x \in X$ and all $\varepsilon > 0$, we have, $g_0 \in P_G(x, \varepsilon)$ if and only if $(x - g_0) \in \hat{G}_\varepsilon$

Proof:

$g_0 \in P_G(x, \varepsilon)$ if and only if by [Theorem1.12], $(x - g_0) \perp_\varepsilon \hat{G}_\varepsilon$ if and only if $(x - g_0) \in \hat{G}_\varepsilon$.

O. Corollary 1.15

Let G be a subspace of a normed linear space X, and let $\varepsilon > 0$, $x \in X$. Then,

$P_G(x, \varepsilon) = G \cap (x - \hat{G}_\varepsilon)$

Proof:

©IJRASET (UGC Approved Journal): All Rights are Reserved

$g_0 \in G \cap (x - \hat{G}_\varepsilon)$ if and only if $g_0 \in G$, and $g_0 \in (x - \hat{G}_\varepsilon)$ if and only if $g_0 \in G$ and $g_0 = x - \hat{g}$, where $\hat{g} \in \hat{G}_\varepsilon$ if and only if $g_0 \in G$, $\hat{g} = (x - g_0) \in \hat{G}_\varepsilon$ if and only if $g_0 \in P_G(x, \varepsilon)$ by [Lemma 1.14].
Therefore, $P_G(x, \varepsilon) = G \cap (x - \hat{G}_\varepsilon)$
Hence the proof

P. Theorem 1.16

Let G be a subspace of a normed linear space X, $\varepsilon > 0$, and $\varepsilon \geq \alpha$. Then,
$\hat{G} \subseteq \hat{G}_\alpha \subseteq \hat{G}_\varepsilon$, and therefore $\cap_{\varepsilon>0}\hat{G}_\varepsilon = \hat{G}$

Proof:

Let $x \in \hat{G}$, then $\| x \| \leq \| x - g \|$ for all $g \in G$.
Now $\| x \| \leq \| x - g \| \leq \| x - g \| + \alpha$ [$\alpha > 0$], so, we have $x \in \hat{G}_\alpha$.
Hence $\hat{G} \subseteq \hat{G}_\alpha$ ……(1)
Let $x \in \hat{G}_\alpha$, then $\| x \| \leq \| x - g \| + \alpha \leq \| x - g \| + \varepsilon$ [$\varepsilon > \alpha$], this implies that $x \in \hat{G}_\varepsilon$, and so, $\hat{G}_\alpha \subseteq \hat{G}_\varepsilon$ ……..(2)
(1) and (2) together imply that $\hat{G} \subseteq \hat{G}_\alpha \subseteq \hat{G}_\varepsilon$,
Now, we show $\cap_{\varepsilon>0}\hat{G}_\varepsilon = \hat{G}$
From above we have $\hat{G} \subseteq \cap_{\varepsilon>0}\hat{G}_\varepsilon$
conversely, let $x \in \cap_{\varepsilon>0}\hat{G}_\varepsilon$,
Then for all $\varepsilon > 0$, $0 \leq \| x \| \leq \| x - g \| + \varepsilon$ for all $g \in G$, then for all $n \in N$,
$0 \leq \| x \| \leq \| x - g \| + \frac{1}{n}$ for all $g \in G$:
As $n \to \infty$, $\| x \| \leq \| x - g \|$ for all $g \in G$, then $x \in \hat{G}$,
and so,
$\cap_{\varepsilon>0}\hat{G}_\varepsilon \subseteq \hat{G}$
Therefore $\cap_{\varepsilon>0}\hat{G}_\varepsilon = \hat{G}$
Hence the proof.

Q. Lemma 1.17

Let G be a subspace of a normed linear space X. Then.

1) If $\varepsilon > 0$, $x, g \in X$ and $x \perp_\varepsilon g$, then $x \perp_\delta g$ for all $\delta \geq \varepsilon$.
2) If $x, g \in X$ and $x \perp_B g$, then $x \perp_\varepsilon g$ for all $\varepsilon > 0$.
3) If $x \in X$, and $\varepsilon > 0$, then $0 \perp_\varepsilon x$, $x \perp_\varepsilon 0$.
4) If $x \perp_\varepsilon g$ and $|\beta| < 1$, then $\beta x \perp_\varepsilon \beta g$.

Proof:

(a) Let $\varepsilon > 0$, $x, g \in X$ and $x \perp_\varepsilon g$, then by [Definition 1.11] we have
$\| x \| \leq \| x + \alpha g \| + \varepsilon$, where $| \alpha | \leq 1$ and $\varepsilon > 0$
Then, $\| x \| \leq \| x + \alpha g \| + \varepsilon \leq \| x + \alpha g \| + \delta$, [since $\delta \geq \varepsilon$]
Therefore, $x \perp_\delta g$
(b) Let $x, g \in X$ and $x \perp_B g$, then $\| x \| \leq \| x + \alpha g \|$ for all $\alpha \in \mathbb{R}$
Since $\varepsilon > 0$, then $\| x \| \leq \| x + \alpha g \| \leq \| x + \alpha g \| + \varepsilon$ for all $| \alpha | \leq 1$
Hence $x \perp_\varepsilon g$ for all $\varepsilon > 0$
(c) Let $x \in X$ and $\varepsilon > 0$, then $\| 0 \| \leq \| 0 + \alpha x \| + \varepsilon$, and so $0 \perp_\varepsilon x$.
We have also $\| x \| \leq \| x \| + \varepsilon$, then $\| x \| \leq \| x + \alpha 0 \| + \varepsilon$,
Hence $x \perp_\varepsilon 0$.
(d) Let $x \perp_\varepsilon g$, and $| \beta | < 1$, then $\| x \| \leq \| x + \alpha g \| + \varepsilon$.
Multiply both sides by $| \beta |$,
we get $|\beta| \| x \| \leq \| \beta x + \beta \alpha g \| + | \beta | \varepsilon$
$\leq \| \beta x + \alpha_1 g \| + \varepsilon$, and so
$\| \beta x \| \leq \| \beta x + \alpha_1 g \| + \varepsilon$
Therefore, $\beta x \perp_\varepsilon \beta g$
Hence the proof

©IJRASET (UGC Approved Journal): All Rights are Reserved

R. Theorem 1.18

Let G be a subspace of a normed linear space X. If $x \in X$, $\varepsilon > 0$ and $\delta \geq \varepsilon$, then $P_G(x, \varepsilon) \subseteq P_G(x, \delta)$.

Proof:

Let $g_0 \in P_G(x, \varepsilon)$. Then by [Definition 1.7], we have

$\| x - g_0 \| \leq \| x - g \| + \varepsilon$ for all $g \in G$ and $\varepsilon > 0$

Then $\| x - g_0 \| \leq \| x - g \| + \varepsilon \leq \| x - g \| + \delta$

[since $\delta > \varepsilon$], then, $g_0 \in P_G(x, \delta)$.

Therefore $P_G(x, \varepsilon) \subseteq P_G(x, \delta)$

Hence the proof


II. CONCLUSION

Here, I conclude my paper as ε-Best approximation and ε- orthogonality has the properties which are similar to the properties of best approximation.

REFERENCE

[1] Mazaheri .H ,Maalek Ghaini.F.m, ' Quasi –Orthogonality of the best approximation sets', Non linear Analysis ,65534-537(2006)
[2] White.A,2-Banach Spacs,Math.Nachr,42(1969),43-60
[3] Rivlin.Theodore J 'The Approximation of Function' 1969

©IJRASET (UGC Approved Journal): All Rights are Reserved



10.22214/IJRASET

45.98

IMPACT FACTOR: 7.129

IMPACT FACTOR: 7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)