



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 6 Issue: IV Month of publication: April 2018

DOI: <http://doi.org/10.22214/ijraset.2018.4047>

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

Geology and Strain History of Rocks Exposed around Village Modi, North East of Gogunda, Udaipur District, Rajasthan

Anju Sharma

Assistant Professor, Sunrise Group of Institutions, Udaipur (RAJ).

Abstract: The rocks exposed around village Modi, NE of Gogunda, Rajasthan includes metapelites, quartzites and metaultramafics. Lithologically the rocks of this area belong to deep sea facies condition (Jharol formation, Aravalli Supergroup, Udaipur, Rajasthan). Structural Analysis of rocks reveals subjected to the area has been subjected to three phases of deformation. The second deformation (AF₂) is the most dominant structural event which has controlled the whole outcrop pattern. The strain analysis suggest that the initiation of AF₁ and AF₂ deformations took place by buckling followed by flattening resulting in Class IC (flattened parallel folds)

Keywords: Aravalli Supergroup, Meta Ultramafics, Udaipur, Rajasthan.

I. INTRODUCTION

The rock types exposed towards NE of Gogunda belongs to Aravalli as well as Delhi Supergroup out of which the former covers greater part of the area. These rocks have been involved in complex superposed deformation and thus selected for the study of detailed geological and structural behavior of rocks. The present work has also revealed a detailed study of meta-ultramafics exposed in the area regarding to its tectonic, stratigraphic status and strain history of rocks. The area selected for the study is about 40 sq. km and has been mapped on 1:10,000 scale. This mapping has revealed a lot of details about the behavior of the small scale structural elements of deformed layers, structural geometry and strain history of the area which is briefly discussed in this paper.

A. Geological Setting

The geological details of the present area of investigation are given in the Fig.1. Stratigraphically, the rocks of the area belongs to Jharol formation of deep sea facies sequences of Aravalli Supergroup (Roy, et.al., 1991). According to GSI (1981), also rocks of Jharol Group are exposed around village Modi, and Chippala. This includes garnet-mica schist and meta-ultramafics with thin intercalatory bands of quartzite. On the western margin of the area, rock types of Delhi Supergroup are exposed which includes mainly quartzite. It belongs to Kelwara Formation of Gogunda Group (GSI, 1981).

The stratigraphic succession of the rock of the area is given in table1.


Table1: Stratigraphy of the area around Modi

| | | | | |
|--------------------------------------|---------------|--|---------------------------|--|
| Delhi Super Group | Gogunda Group | Kelwara Formation Antalia Formation Richer Formation | Kelwara Formation exposed | Quartzite |
| -----Structural discordance----- | | | | |
| Aravalli Jharol Super Group | Group | Samlaji Formation Goran Formation | Goran Formation exposed | Quartz veins/ meta-ultramafics |
| | | | | associated with Quartzite/mica-schist & garnet-mica schist |
| -----Unconformity(not exposed)----- | | | | |
| Banded Gneissic Complex | | not exposed | | Exposed further south near Bagdunda (chauhan,1983) |

Nomenclature as per GSI(1981).

NE of Gogunda Udaipur, Rajasthan

*Department of Geology, Mohan Lal Sukhadia University, Udaipur, Rajasthan



B. Structural Setting

The Area has been subjected to three episodes of deformations i.e. AF_1 , AF_2 and AF_3 . Out of which the second deformation is the most dominant one. The first deformation is less conspicuous due to superimposition by AF_2 folds. First deformation is invariably associated with axial planar schistosity (S_1) which are mostly observed parallel to bedding planes (S_0) indicating the isoclinal nature of mostly observed parallel to bedding planes (S_0) indicating the isoclinal first folding. The second deformation (AF_2) is observed as open to isoclinal, upright folds trending NNE-SSW. S_2 cleavages developed axial planar to these folds (Plate 1a, 1b) and are observed parallel to S_1 schistosity except the hinge of the folds. The third deformation (AF_3) is less penetrative and is observed at few places striking ESE-WNW with vertical axial plane. Among linear structures intersection lineations are most common features due to superposed deformation (Plate 1c). Early lineations deformed as a result of later deformation is preserved in different rocks (Plate 1d). Lineations are also marked by crenulations, small-scale folds axes, mullions and boudins.

C. Strain History

Strain determination of rocks of the area is done by studying the behavior of different layer contrasts. In the present attempt to strain estimation, geometric techniques as proposed by Ramsay (1967) and Huddleston (1977) are employed. For this purpose profiles of folds (traced by quartz veins in garnet-mica schist and meta-ultramafics) of different generations have been used (Fig. 2a)

Biot (1961) predicted that wavelength to thickness ratio of buckled layer would be constant if the viscosity ratio of the fold layer and the host medium is constant. Huddleston (1973) and Huddleston and Stephanson (1973) have concluded that in a layered

sequence of comparable viscosity ratio, folding is developed by buckling if the ratio of the dominant wavelength to thickness of competent layer in the holds of different viscosity is constant. The wave length is measured in terms of arc length. De Sitter (1958) suggested that due to compressive strain the parallel shape of buckle folds is modified due to flattening thus, resulting into flattened parallel folds. Taking into consideration all the above aspects, the following plots have been prepared for the different folds.





Fig.2 (a) :Profiles of AF_2 Folds traced by quartz Vein(Vq) in schistose rock(Sc)
(b) :Plots of arc length(L) v/s thickness(t) of AF_1 and AF_2 folds.

Arc length (L) vs. thickness (t) plots of AF_1 and AF_2 folds (Fig. 2b) is prepared on log-log scale which shows that arc length is directly proportional to wave length pointing to buckling origin of folds (Biot, 1961; Ramberg, 1963; Naha et.al., 1977; Sharma et. al., 1988).

Different folds of AF_1 and AF_2 generations are chosen for the measurement of orthogonal thickness (t_α) at different limb dip (α). The t_α'/α plots shows that majority of folds shows Class 1C geometry (i.e. flattened parallel folds) with Class 1B geometry at the hinge (Fig. 3a). Plots of $t' t_\alpha'/\alpha$ also give an idea about the apparent strain ratio $\sqrt{\lambda_2/\lambda_1}$.

The values shows a wide range from 0.1 to 0.5 for AF_1 and 0.1 to 1.0 for AF_2 folds

The range of $\sqrt{\lambda_2/\lambda_1}$ values indicates that different folds underwent varying amount of flattening (Roy, 1978) Fig. 3c).

The ϕ/α plots are prepared for the same set of folds for which t_α'/α plots have been prepared. Here, also most of the folds satisfies the condition of Class 1C geometry i.e. $\alpha > \phi_\alpha > \theta_2$ (Fig. 3b)

D. Origin of AF_1 and AF_2 folds:

Thus, the constant values of L/t ratio, nature of t_α'/α and ϕ_α/α plots, wide range of $\sqrt{\lambda_2/\lambda_1}$ values and different field evidences such as refraction of axial plane cleavages across layers of different competency contrasts, occurrence of radial tension cracks at hinge of the folds in competent layers, and reversal of curvature at the hinge of folds all of them indicate that both AF_1 and AF_2 folds are initiated by buckling followed by flattening, normal to their axial surface.

E. Tectonic and stratigraphic status of metaultramafics of Jharol Formation:

The meta-ultramafics exposed in the area belongs to Jharol Formation of deep sea facies sequences; the emplacement of these rocks took place along an important lineament (Kaliguman Lineament). It is emplaced parallel to the Aravalli Quartzite striking NS to NE-SW. Moreover, it has participated in all the three deformation episodes (i.e. AF_1 , AF_2 , and AF_3). These evidences suggest that the emplacement of these lithounits took place contemporaneous with sedimentation but prior to Aravalli deformation.

II. CONCLUSION

The lithological aspects suggest that the rocks of this area are deposited in deep sea facies conditions (Jharol Formation). The area has been subjected to three phases of deformation. The second deformation (AF_2) is the most dominant structural event which has controlled the whole outcrop pattern of the area. The strain analysis suggest that the initiation of both AF_1 and AF_2 deformations took place by buckling followed by flattening resulting in Class 1C (flattened parallel folds).

The compressive strain ratio varies from 0.1 to 1.0. The metaultramafics exposed in the area is early Proterozoic in age and have been emplaced prior to Aravalli deformation (AF_1) in the associated lithounits.

REFERENCES

- [1] Bakliwal, P.C., and Ramasamy, S.N., 1987. Lineament fabric of Rajasthan and Gujarat. *Tec. Geol. Surv. Ind.*, v.113 (7), pp. 54-64.
- [2] Biot, M.A., 1961, Theory of folding of stratified Viscoelastic media and its implications in tectonics and orogenesis, *Geol. Soc. Am. Bull.*, v. 72, pp. 1595-1632.
- [3] Chauhan, N.K., 1983. Geology and structure of the area around Bagdunda. Udaipur district, Raj. Unpublished Ph.D. Thesis, University of Rajasthan Jaipur, 156 p.
- [4] Chauhan, N.K., Sharma B.L., and Sabah A. Mohammed., 1996. Structural geometry and strain history of the Early Proterozoic Aravalli rocks of Gorimari, Udaipur District, Rajasthan, *Jour Geol. Soc., India.* v. 47, pp. 59-74.
- [5] De Sitter, I.U., 1958. Boudins and parasitic folds in relation to cleavage and folding. *Geologie Mijub.*, v. 20, pp. 277-286.
- [6] G.S.I., Publication 1981. Explanatory brochure to the geological Map of the Aravalli region southern Rajasthan and northeastern Gujarat, Published under the direction of V.S., Krishnaswamy, Director General, Geol, Surv, India, 38 p.
- [7] Heron. A.M., 1953. Geology of the central Rajasthan Mem Geol Surv. India., v.79, 389p.
- [8] Huddleston, P.J., and Stephansson, O., (1973): Layer shortening and fold -shape development in the buckling of single layers. *Tectonophysics*, v.17. pp. 299-321.
- [9] Huddleston, P.J., 1973a. Fold morphology and some geometrical implication of theories of fold development *Tectonophysics*, v. 16, pp. 1-46.
- [10] Huddleston, P.J., 1973b. An analysis of 'single-layer' folds developed experimentally in viscous media. *Tectonophysics*, v. 16, pp. 189-214.
- [11] Huddleston, P.J., 1973c. An analysis and interpretation of minor folds development in the Moine rocks of Monar, Scotland. *Tectonophysics*, v. 17, pp. 89-132.
- [12] Naha, K., and Halyburton, R.V., 1977. Structural pattern and strain history of a superposed fold system in the Precambrian of Central Rajasthan, India, II, strain history. *Precambrian Res.*, v. 4, pp. 85-111.
- [13] Ramsay, J.G., 1967. Folding and fracturing of rocks. Mc Graw Hill, New York, 568 p.
- [14] Roy, A.B., Sharma, B.L., Paliwal B.S., Chauhan, N.K., Nagori, D.K., Golani, P.R., Bejarniya, B.R., Bhu, H., and Ali, Sabah M., 1993. Lithostratigraphy and tectonic evolution of the Aravalli Supergroup: A Proterozoic synclinal Sequence. In: S.M. Casshyap (ed.), *Rift Basins and Aulacogens*. Gyanodaya Prakashan, Nainital, pp. 734-90.
- [15] Roy, A.B. and Jhakar, S.R., 2002. Geology of Rajasthan (Northwest India), Precambrian to Recent. Scientific Publishers (India), Jodhpur. 421 p.
- [16] Sharma B.L., Chauhan, N.K., and Bhu, H., 1988. Structural geometry and deformation history of the early Proterozoic Aravalli rocks from Bagdunda, district Udaipur, Rajasthan. In: A.B. Roy (ed.) *Precambrian of Aravalli Mountain, Rajasthan, India*, Mem. Geol. Soc. India, v.7p. 169-191.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)