



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 6 Issue: IV Month of publication: April 2018

DOI: <http://doi.org/10.22214/ijraset.2018.4784>

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

LICT Subdivision Double Domination in Graphs

M. H. Muddebihal¹, Suhas P. Gade²

¹Department of Mathematics Gulbarga University, Kalburgi-585106, Karnataka, India.

²Department of Mathematics, Sangameshwar College, Solapur – 413001, Maharashtra, India.

Abstract: Let $S(G)$ be the subdivision graph of G . The lict graph of $n[S(G)]$ of $S(G)$ is a graph whose vertex set is the union of the set of edges and set of cutvertices of $S(G)$ in which two vertices adjacent if and only if the corresponding members are adjacent or incident. A subset D^d of $V[n(S(G))]$ is double dominating set of $n[S(G)]$ if for every vertex $v \in V[n(S(G))]$, $|N[v] \cap D^d| \geq 2$, that is v is in D^d and has at least one neighbour in D^d or v is in $V[n(S(G))] - D^d$ and has at least two neighbours in D^d . The lict subdivision double dominating number $\gamma_{ddns}(G)$ is a minimum cardinality of the lict subdivision double dominating set of G and is denoted by $\gamma_{ddns}(G)$. In this paper, we establish some sharp bounds for $\gamma_{ddns}(G)$. Also some upper and lower bounds on $\gamma_{ddns}(G)$ in terms of the vertices, edges and other different parameters of G and not in terms of the element of $n[S(G)]$. Further, its relation with other different dominating parameters is also obtained.

Subject classification number: AMS – 05C69, 05C70.

Keyword: Lict subdivision graph/ Dominating set/Double domination

I. INTRODUCTION

In this paper, all the graphs considered here are simple, finite, non-trivial, undirected and connected. The vertex set and edge set of graph G are denoted by $V(G) = p$ and $E(G) = q$ respectively. Terms not defined here are used in the sense of Harary [1]. The neighbourhood of a vertex $v \in V$ is defined by $N(v) = \{u \in V/uv \in E\}$. The close neighbourhood of a vertex v is $N[v] = N(v) \cup \{v\}$. The order $|V(G)|$ of G is denoted by p . The degree of v is $d(v) = |N(v)|$. The maximum degree of a graph G is denoted by $\Delta(G)$ and the minimum degree is denoted by $\delta(G)$. A vertex cover in a graph G is a set of vertices that covers all the edges of G . The vertex covering number $\alpha_0(G)$ is the minimum cardinality of a vertex cover in G . A set of vertices in a graph G is called independent set if no two vertices in the set are adjacent. The vertex independence number $\beta_0(G)$ is the maximum cardinality of an independent set of vertices. For a vertex v of a graph G , the eccentricity $e(v)$ is the distance between v and a vertex farthest from v . The maximum eccentricity is its diameter, $diam(G)$. A set D of vertices in a graph G is called a dominating set of G if every vertex in $V - D$ is adjacent to some vertex in D . The domination number of G , denoted by $\gamma(G)$ is the minimum cardinality of a dominating set. The domination in graphs with many variations is now well studied in graph theory. A thorough study of domination appears in [2]. Let $S(G)$ be the subdivision graph of G . The lict graph of $n[S(G)]$ of $S(G)$ is a graph whose vertex set is the union of the set of edges and set of cutvertices of $S(G)$ in which two vertices adjacent if and only if the corresponding members are adjacent or incident. A subset D^d of $V[n(S(G))]$ is double dominating set of $n[S(G)]$ if for every vertex $v \in V[n(S(G))]$, $|N[v] \cap D^d| \geq 2$, that is v is in D^d and has at least one neighbour in D^d or v is in $V[n(S(G))] - D^d$ and has at least two neighbours in D^d . The lict subdivision double dominating number $\gamma_{ddnb}(G)$ is a minimum cardinality of the lict subdivision double dominating set of G and is denoted by $\gamma_{ddns}(G)$. The graph valued function related to domination parameters have been studied in [4,5,6,7,9,10,11,12,13,14,15]. Further in [8], the subdivision of G with graphvalued function and related with domination parameters has been established. In this paper, we establish some sharp bounds for $\gamma_{ddns}(G)$. Also some upper and lower bounds on $\gamma_{ddns}(G)$ in terms of the vertices, edges and other different parameters of G and not in terms of the element of $n[S(G)]$. Further, its relation with other different dominating parameters is also obtained.

II. RESULTS

We need the following theorems to prove our results.

- 1) **Theorem A[1]:** For any path P_n , the vertex covering number is $\alpha_0(P_n) = \begin{cases} \frac{n}{2} & \text{if } n \text{ is even} \\ \frac{n-1}{2} & \text{if } n \text{ is odd} \end{cases}$

- 2) *Theorem B[1]*: For any path P_n , the edge covering number is $\alpha_1(P_n) = \begin{cases} \frac{n}{2} & \text{if } n \text{ is even} \\ \frac{n+1}{2} & \text{if } n \text{ is odd} \end{cases}$
- 3) *Theorem C[2]*: A graph G is eulerian if and only if G is of even degree.
- 4) *Theorem D[3]*: For any connected graph G , with $p \geq 4$, $G \neq K_p$, $\gamma_{ss} = \alpha_0(G)$.

III. THE LICT SUBDIVISION DOUBLE DOMINATION NUMBER OF A GRAPH.

In this section, we characterise the lict subdivision double domination number by giving a necessary and sufficient condition for it and also establish the conditions for a lict subdivision double dominating set

- 1) *Theorem 3.1*: A double dominating set D^d of the $n[S(G)]$, is a lict subdivision double dominating set of G , if and only if the following conditions hold.
- (i) $n[S(G)] - D^d$ has at least two vertices.
- (ii) For any two vertices $u, v \in V[n(G)] - D^d$, every uv path contains a vertex of D^d .

We now give a characterization of lict subdivision double dominating set of G which is minimal.

- 2) *Theorem 3.2*: A double dominating set D^d of the $n[S(G)]$ is minimal if and only if for every vertex $v \in D^d$ either (i) $|N(v) \cap D^d| \leq 2$ or (ii) there $\exists$ a vertex $u, \in V[n(G)] - D^d$ such that $|N(v) \cap D^d| = 2$ and $u \in N(v)$.

Proof: Let D^d be a minimal lict subdivision double dominating set of G . Suppose that there exists a vertex $v \in D^d$ for which $|N(v) \cap D^d| \geq 2$ and for every vertex $u \in V[n(S(G))] - D^d$, either $|N(v) \cap D^d| \geq 2$ or $u \notin N(v)$. Then consider $D^{d'} = D^d - \{v\}$, since v is adjacent to at least two vertices of $D^{d'}$ it follows that $D^{d'}$ is double dominating set of $n[S(G)]$, which is contradicting to the minimality of D^d . Conversely, assume that D^d is double dominating set of $n[S(G)]$ satisfying conditions (i) and (ii). For that consider the set $D^{d'} = D^d - \{v\}$ for any vertex $v \in D^d$. If condition (i) holds, then $|N(v) \cap D^{d'}| \leq 2$, which implies that $D^{d'}$ is not a double dominating set of $n[S(G)]$. If the condition (ii) holds. Then there exists a vertex $u \in V[n(S(G))] - D^{d'}$ such that $|N(u) \cap D^{d'}| = 2$ and $u \in N(v)$. But in this case that set $D^{d'}$ would not double dominating to u and hence would not be a double dominating set of $n[S(G)]$. Thus in both condition $D^{d'}$ is not a double dominating set of $n[S(G)]$. Therefore D^d is a minimal double dominating set of $n[S(G)]$.

IV. LOWER BOUNDS FOR $\gamma_{ddns}(G)$.

We establish lower bounds for $\gamma_{ddns}(G)$ in terms of elements of G .

- 1) *Theorem 4.1*: For any connected (p, q) graph G with $p \geq 2$, $\left\lceil \frac{p}{2} \right\rceil + 1 \leq \gamma_{ddns}(G)$. Equality hold if G is P_2 .

Proof: Let $D^d = \{v_1, v_2, \dots, v_n\}$ be a double dominating set of $n[S(G)]$. By the definition of $n[S(G)]$, $V[n(S(G))] = E(S(G)) \cup C(S(G))$. Let D^d be the double dominating set of $n[S(G)]$ such that any vertex $v \in V[n(S(G))] - D^d$, $|N[v] \cap D^d| \geq 2$. Then $\{V[n(S(G))] - D^d\}$ contains at least one vertex which gives $\frac{p}{2} < \left\lceil \frac{p}{2} \right\rceil < \left\lceil \frac{p}{2} \right\rceil + 1 \leq \gamma_{ddns}(G)$.

- 2) *Theorem 4.2*: For any connected (p, q) graph G with $p \geq 2$, $\gamma_{ddns}(G) \leq p + q - 1$.

Proof: Let G be a (p, q) graph, then $V[S(G)] = p + q$. Let $F = \{v_1, v_2, \dots, v_k\}$ be the set of vertices in $S(G)$ and $F \subseteq V[S(G)]$ such that $|F| = \alpha_0(S(G))$ similarly for $E = \{e_1, e_2, \dots, e_n\}$ be the set of edges in $S(G)$, $E \subseteq E[S(G)]$ such that $|E| = \beta_0(S(G))$ and by the definition of $n[S(G)]$, $V[n(S(G))] = E(S(G)) \cup C(S(G))$. Where $E(S(G))$ is the set of edges and $C(S(G))$ is the set of cutvertices in $S(G)$. We consider a set $F \subseteq E(S(G))$. If F corresponding to such vertices $D^d = \{v_1, v_2, \dots, v_q\} \subseteq V[n(S(G))]$ such that $D^d = V[n(S(G))] - C(S(G))$ and any vertex $v \in V[n(S(G))] - D^d$ is dominated by at least two vertices of $n[S(G)]$. Thus D^d is double dominating set of $n[S(G)]$. Now consider $|D^d| \leq \alpha_0(S(G)) + \beta_0(S(G)) - 1 = V[S(G)] - 1 = p + q - 1$. Hence $\gamma_{ddns}(G) \leq p + q - 1$.

- 3) *Theorem 4.3*: For any connected (p, q) graph G with $p \geq 3$, $p + \gamma(G) \leq \gamma_{ddns}(G)$. Equality holds if $G \cong P_p$, $p \geq 3$.

Proof: Let $D = \{v_1, v_2, \dots, v_k\}$ be a minimal dominating set of G such that $|D| = \gamma(G)$. Since in $n(S(G))$, $V[n(S(G))] = E(S(G)) \cup C(S(G))$. Further let $E = \{e_1, e_2, \dots, e_q\}$ be the set of all edges which are incident to the vertices of D becomes $E(S(G)) = 2E$, in $n(S(G))$. Since $V[n(S(G))] = E(S(G)) \cup C(S(G))$. Let $D^d = \{v_1, v_2, \dots, v_n\} \subseteq E(S(G)) \subseteq V[n(S(G))]$ be the double dominating set of $n(S(G))$ such that $|N[v] \cap D^d| \geq 2$. Then D^d form a minimal double dominating set in $n[S(G)]$. Clearly

$|E(S(G))| - |V(G)| \geq |D|$. Thus it follows that $\gamma_{ddns}(G) - p \geq \gamma(G)$. Hence $p + \gamma(G) \leq \gamma_{ddns}(G)$. For equality if $G \cong P_3$. Then in this case $|D| = 1$, further $S(P_3) = P_5$, $\gamma_{ddns}(G) = 4$. Hence $\gamma_{ddns}(G) = 4 = 1 + 3\gamma(G) + p$.

4) **Theorem 4.4:** For any connected (p, q) graph G with $p \geq 3$, then $diam(G) + 2 \leq \gamma_{ddns}(G)$.

Proof: Let $I = \{e_1, e_2, \dots, e_n\} \subseteq E(G)$ be the set of edges which constitutes the longest path between any two distinct vertices of G such that $|I| = diam(G)$. Let $D^d = \{v_1, v_2, \dots, v_n\}$ be the set of vertices in $V[n(S(G))]$ such that for any vertex $v \in V[n(S(G))]$ - D^d is adjacent to at least two vertices of D^d and $|N[v] \cap D^d| \geq 2$. It follows that $|D^d| \geq 2$ and the diametral path includes at least two vertices. Thus $diam \leq \gamma_{ddns}(G) - 2$ it gives $diam(G) + 2 \leq \gamma_{ddns}(G)$.

5) **Theorem 4.5:** For any connected (p, q) graph G with $p \geq 2$, $\frac{(2q-p(p-3))}{2} \leq \gamma_{ddns}(G)$.

Proof: Let $D^d = \{v_1, v_2, \dots, v_n\} \subseteq V[n(S(G))]$ be the set of vertices and every vertex $v \in V[n(S(G))]$ - D^d is adjacent to at least two vertices of D^d , thus D^d itself is a double dominating set of $n[S(G)]$. Since $V[n(S(G))] = E(S(G)) \cup C(S(G))$. Then there exists a vertex $v \in D^d$ which is not adjacent to any vertex of $V[n(S(G))]$. This implies that $q \leq \frac{p(p-1)}{2} - (p - \gamma_{ddns}(G))$. Which is $2q \leq p^2 - p - 2p + 2\gamma_{ddns}(G)$. Hence the result.

6) **Theorem 4.6:** For any connected (p, q) graph G , $\left\lceil \frac{2p}{\Delta(G)+2} \right\rceil \leq \gamma_{ddns}(G)$.

Proof: Let D^d be a minimal double dominating set of $n[S(G)]$. Since in $n[S(G)]$, $V[n(S(G))] = E(S(G)) \cup C(S(G))$. Let k denote the number of edges between D^d and $V[n(S(G))]$ - D^d . Since for any connected graph G , there exists at least one vertex $v \in V(G)$ such that $\deg(v) = \Delta(G)$, $k \leq \Delta(G) \cdot \gamma_{ddns}(G)$. Also since each vertex v in $V[n(S(G))]$ - D^d is adjacent to at least two vertices in D^d , $k \geq 2(p - \gamma_{ddns}(G))$. From these two inequalities, $2p - 2p\gamma_{ddns}(G) \leq \Delta(G) \cdot \gamma_{ddns}(G)$. It follows that $\frac{2p}{\Delta(G)+2} \leq \left\lceil \frac{2p}{\Delta(G)+2} \right\rceil \leq \gamma_{ddns}(G)$.

7) **Theorem 4.7:** For any non-trivial tree T of order p with l leaves and s support vertices, $(p - l - s + 4) \leq \gamma_{ddns}(T)$.

Proof: To prove that if T is a tree of order $p \geq 2$ with l leaves and s support vertices then $\gamma_{ddns}(T) \geq (p - l - s + 4)$. We use mathematical induction on p . Let $p = 2$, then $T = P_2$ and $diam(T) = 1$ it implies that $\gamma_{ddns}(T) = p - l - s + 4 = 2$. Let $p = 3$ then T is a star and $diam(G) = 2$, it implies that $\gamma_{ddns}(T) = (p - l - s + 4) = 4$. Let $p = 4$ then T may be P_4 and $diam(G) = 3$ it implies that $\gamma_{ddns}(T) > (p - l - s + 4)$. For $p = n$, $\gamma_{ddns}(T) > (p - l - s + n)$. Further it is also true for $p = n + 1$. Hence we obtain the desired result.

8) **Theorem 4.8:** For any connected (p, q) graph G with $p \geq 2$, $\gamma(G) + 1 \leq \gamma_{ddns}(G)$. Equality hold if G is K_2 .

Proof: Let $D = \{v_1, v_2, \dots, v_k\}$ be a minimal dominating set of G such that $|D| = \gamma(G)$. Since in $n(S(G))$, $V[n(S(G))] = E(S(G)) \cup C(S(G))$, where $E(S(G))$ is the set of edges and $C(S(G))$ is the set of cutvertices in $S(G)$. We consider the following cases. Case 1: Suppose G is a tree. Then clearly for any tree T , $D^d = \{v_1, v_2, \dots, v_n\}$ set of vertices in $n[S(G)]$ such that $D^d = V[n(S(G))] - C(S(G))$ is dominated by at least two vertices of $n[S(G)]$. Since the number of edges of $S(G)$ are more than that of G , which gives $V(G) \subset V[n(S(G))]$. It follows that $|D^d| > |D| + 1$ which gives $\gamma_{ddns}(G) > \gamma(G) + 1$. Case 2: Suppose G is not a tree. Then there exists at least one cycle. Let D be a minimal dominating set of G , then $|D| = \gamma(G)$. Suppose there exists a cycle of length l , then in $S(G)$, the cycle length be $2l$. Let $E = \{e_1, e_2, \dots, e_k\} \subseteq E[S(G)]$, such that $|E| = \gamma(G)$. In $n[S(G)]$, $V[n(S(G))] = E(S(G)) \cup C(S(G))$. Suppose $F \subset C[S(G)]$. Then $\{E\} \cup \{F\} \subseteq V[n(S(G))]$ such that $\forall v_i \in V[n(S(G))] - \{E\} \cup \{F\}$ is adjacent to at least two vertices of $V[n(S(G))]$. Hence $|E| \cup |F|$ is a minimal double dominating set $n[S(G)]$ such that $|E| \cup |F| = \gamma_{ddns}(G)$ -set. Clearly $|D| \leq |E| \cup |F|$ which gives $\gamma(G) + 1 \leq \gamma_{ddns}(G)$. For equality, let $G = K_2$, we have $\gamma(G) = 1$. Further $S(K_2) = P_3$ and $n[S(K_2)] = C_3$ and $\gamma_{ddns}(G) = 2$, $\gamma(G) = 1$. Hence $\gamma_{ddns}(G) = 2 = \gamma(G) + 1$.

9) **Theorem 4.9:** For any connected (p, q) graph G , $\left\lceil \frac{p(9-p)-6}{6} \right\rceil \leq \gamma_{ddns}(G)$.

Proof: Let D^d be a γ_{ddns} -set of G . Since in $n[S(G)]$, $V[n(S(G))] = E(S(G)) \cup C(S(G))$. Let t_1 denote the number of edges in $n[S(G)]$ incident to the vertices of $V[n(S(G))]$ - D^d only. Also t_2 denotes the number of edges in $n[S(G)]$ incident to the vertices of D^d only. Then $\frac{p(p-1)}{2} \geq t_1 + t_2 \geq 4|V(G) - D^d| - 2 + |D^d| - 1$ it implies that $\frac{p(p-1)}{2} \geq 4p - 4|D^d| - 2 + |D^d| - 1 = 4p - 3|D^d| - 3$. It gives that $3|D^d| \geq \frac{p(9-p)-6}{2}$. Hence $\left\lceil \frac{p(9-p)-6}{6} \right\rceil \leq \gamma_{ddns}(G)$.

10) **Theorem 4.10:** For any connected (p, q) graph G with $p \geq 2$, $2(p - q) \leq \gamma_{ddns}(G)$.

Proof: Let $D^d = \{v_1, v_2, \dots, v_n\}$ be the minimal set of vertices which covers all the vertices in $V[n(S(G))]$. Suppose for every vertex of $v \in V[n(S(G))]$ is adjacent to at least two vertices of D^d , clearly D^d forms a double dominating set of $n(S(G))$. Let any vertex $v \in D^d$ which is not adjacent to any vertex of $V[n(S(G))] - D^d$. Then $2q \geq |D^d| + 2|V(G) - D^d|$ it gives that $2q \geq |D^d| + 2p - 2|D^d|$. Therefore $|D^d| \geq 2p - 2q$ this implies that $2(p - q) \leq \gamma_{ddns}(G)$.

V. SPECIFIC VALUES OF $\gamma_{ddns}(G)$.

We found some constraints for which $\gamma_{ddns}(G)$ follows the equality relation with other domination parameters of G .

1) **Theorem 5.1:** For any path P_p with $p \geq 3$, $\gamma_{ddns}(P_p) = \begin{cases} 4\alpha_0(P_p) - 2, & \text{if } p \text{ is even} \\ 4\alpha_0(P_p), & \text{if } p \text{ is odd} \end{cases}$.

Proof: Let P_p be the path with $p \geq 3$ vertices. Consider $V = \{v_1, v_2, \dots, v_n\}$ be the vertices and $E = \{(v_i, v_{i+1})\}$, $i = 1, 2, 3, \dots$ be the edge set of path P_p . Since in $n[S(G)]$, $V[n(S(G))] = E(S(G)) \cup C(S(G))$. We consider the following cases

a) **Case 1:** If p is even. Then by Theorem A[1], we have $\alpha_0(P_p) = \frac{p}{2}$ it implies that $p = 2\alpha_0(P_p)$. Since $\gamma_{ddns}(P_p) = 2q = 2(p - 1)$, we have $\gamma_{ddns}(P_p) = 2p - 2 = 2[2\alpha_0(P_p)] - 2 = 4\alpha_0(P_p) - 2$.

b) **Case 2:** If p is an odd. Then by Theorem A[1], we have $\alpha_0(P_p) = \frac{p-1}{2}$ it implies that $p - 1 = 2\alpha_0(P_p)$. Since $\gamma_{ddns}(P_p) = 2q = 2(p - 1) = 4\alpha_0(P_p)$.

2) **Theorem 5.2:** For any path P_p with $p \geq 3$, $\gamma_{ddns}(P_p) = \begin{cases} 4\alpha_1(P_p) - 2, & \text{if } p \text{ is even} \\ 4\alpha_1(P_p) - 4, & \text{if } p \text{ is odd} \end{cases}$.

Proof: Let P_p be the path with $p \geq 3$ vertices. Consider $V = \{v_1, v_2, \dots, v_n\}$ be the vertices and $E = \{(v_i, v_{i+1})\}$, $i = 1, 2, 3, \dots$ be the edge set of path P_p . Since in $n[S(G)]$, $V[n(S(G))] = E(S(G)) \cup C(S(G))$. We consider the following cases.

a) **Case 1:** If p is even. Then by Theorem B[1], we have $\alpha_1(P_p) = \frac{p}{2}$ it implies that $p = 2\alpha_1(P_p)$. Since $\gamma_{ddns}(P_p) = 2q = 2(p - 1)$, we have $\gamma_{ddns}(P_p) = 2p - 2 = 2[2\alpha_1(P_p)] - 2 = 4\alpha_1(P_p) - 2$.

b) **Case 2:** If p is an odd. Then by Theorem B[3], we have $\alpha_1(P_p) = \frac{p+1}{2}$ it implies that $p - 1 = 2\alpha_1(P_p)$. Since $\gamma_{ddns}(P_p) = 2q = 2(p - 1) = 2p - 2 = 4\alpha_1(P_p) - 4$.

VI. UPPER BOUNDS FOR $\gamma_{ddns}(G)$.

We establish upper bounds for $\gamma_{ddns}(G)$ in terms of elements of G .

1) **Theorem 6.1:** For any connected (p, q) graph G with $p \geq 2$, $\gamma_{ddns}(G) \leq 2q$.

Proof: Since $V[n(S(G))] = E(S(G)) \cup C(S(G))$, $V(S(G)) = p + q$. Let D^d be double dominating set of $n[S(G)]$. Then by definition of list subdivision double domination $|D^d| \geq 2$. Further by definition of $n[S(G)]$, $2q - \gamma_{ddns}(G) \geq 0$. Clearly it follows that $\gamma_{ddns}(G) \leq 2q$.

2) **Theorem 6.2:** For any connected (p, q) graph G with $p \geq 2$, $\gamma_{ddns}(G) \leq p + q - \delta(G)$.

Proof: Let $D^d = \{v_1, v_2, \dots, v_n\} \subseteq V[n(S(G))]$ be the set of vertices and every vertex $v \in V[n(S(G))] - D^d$ such that $|N[v] \cap D^d| \geq 2$. Thus it is clear that $|D^d| \geq 2$. Since for any graph G there exists at least one vertex $v \in V(G)$ such that $\deg(v) = \delta(G)$. By definition of $n[S(G)]$, $V[n(S(G))] = E(S(G)) \cup C(S(G))$. Then there exists a vertex $v \in G$ such that $\deg(v) = \delta(G)$. Thus $\delta(G) \leq p - |D^d| + q$, which implies that $|D^d| \leq p + q - \delta(G)$. Hence the result.

3) **Theorem 6.3:** For any connected (p, q) graph G with $n[S(G)] \neq K_p$, $\gamma_{ddns}(G) + \gamma_{ssns}(G) \leq 2q + C$. Where C is the number of cut vertices in $S(G)$.

Proof: Suppose G has $p \leq 3$ then γ_{ddns} -set does not exist. Now we consider any graph with $p \leq 4$, such that $n[S(G)] \neq K_p$. Since $\gamma_{ddns}(G) \leq 2q$ and from Theorem D[3] $\gamma_{ssns}(G) = \alpha_0[n(S(G))]$. Further $\gamma_{ddns}(G) + \gamma_{ssns}(G) \leq 2q + \alpha_0[n(S(G))] \leq V[n(S(G))] \leq E(S(G)) \cup C(S(G)) \leq 2q + C$. Hence the result.

4) **Theorem 6.4:** For any non-trivial tree T , the list subdivision of a tree is non-eulerian.

Proof: Let T be a non-trivial tree and $n[S(G)]$ always contain a point of odd degree. Hence by Theorem C[2], the result follows.

NORDHAUS-GADDUM TYPE RESULTS

5) **Theorem 6.5:** For any connected (p, q) graph G with $p \geq 3$ vertices,




$$(i) \gamma_{dms}(G) + \gamma_{dms}(\bar{G}) \leq 4q.$$

$$(ii) \gamma_{dms}(G) \cdot \gamma_{dms}(\bar{G}) \leq 4q^2.$$

REFERENCES

- [1] Gray Chartrand and Ping Zhan, Introduction to Graph Theory, Tata McGraw-Hill Pub.Co.Ltd. (2006).
- [2] F. Harary, Graph Theory, Addison-Wesley, Reading, Mass, 1972.
- [3] Kulli V.R., Theory Of Domination In Graphs, Vishwa International Publication, Gulbarga, India, 2010.
- [4] T.W. Haynes, S. T. Hedetniemi and P.J. Slater, Fundamentals of domination in graphs. Marcel, Dekker Publ. 1998 New York.
- [5] M.H. Muddebihal and Megha Khandelwal , Strong split list domination of a graph, International Journal of Engineering and Scientific Research, Vol.2(7), pp 187-198, 2014.
- [6] M.H. Muddebihal and Megha Khandelwal, Strong split block cut vertex domination of a graph, , International Journal of Mathematics and Computer Applications Research, vol.5(5), pp-73-80, 2015.
- [7] M.H. Muddebihal and Megha Khandelwal, On strong split middle domination of a graph, , International Journal of recent and innovation trends in computing and communication trends in computing and communication vol.4(6), pp-538-541,2016.
- [8] M.H. Muddebihal and Sumangaladevi, Roman list domination in Graphs, Ultra Scientist, vol.24 (3) A, pp 449-458, 2012.
- [9] M.H. Muddebihal and Sumangaladevi, Connected roman domination in graph, International Journal of research in engineering and technology, vol.2 (10) , pp 333-340, 2013.
- [10] M.H. Muddebihal and Sumangaladevi, Roman block domination in graphs, International Journal of research in science and advanced technologies, vol.6 (2), pp 267-275,2013
- [11] M.H. Muddebihal and Sumangaladevi, Roman subdivision domination in graphs, International Journal of engineering, science and research technology, vol.3(3), pp 1441-1447, 2014.
- [12] M.H. Muddebihal and Suhas P. Gade, Line double domination in graphs, International Journal of Mathematics and Computer Application Research vol.5(3), pp 75-80, 2015.
- [13] M.H. Muddebihal and Suhas P. Gade, List double domination in graphs, Global Journal of Pure and Applied Mathematics vol.13(7), pp 3113-3120, 2017.
- [14] M.H. Muddebihal and Suhas P. Gade, Semitotal block double domination in graphs, International Journal of Mathematics trends and Technology, vol.52 (7), 2017.
- [15] M.H. Muddebihal and Suhas P. Gade, Block double domination in graphs, International Journal of Mathematical Archive, vol.9(1), 2018, pp 1-5.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)