iJRASET
International Journal For Research in Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 6 **Issue:** V **Month of publication:** May 2018

DOI: http://doi.org/10.22214/ijraset.2018.5033

www.ijraset.com

Call: 08813907089 | E-mail ID: ijraset@gmail.com

International Journal for Research in Applied Science & Engineering Technology (IJRASET)
ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 6.887
Volume 6 Issue V, May 2018- Available at www.ijraset.com

MHD and Thermal Radiation Effects on Rotating Fluid Past a Moving Vertical Plate with Heat and Mass Transfer in the Presence of Chemical Reaction, OHMIC Heating

K. laxmaiah[1], Dr. D. Raju[2], M. Chenna Krishna Reddy[3]
[1]*Department of Mathematics, Guru Nanak Institutions Technical Campus, Ibrahimpatnam, Ranga Reddy, Telangana – 501 506.*
[2]*Department of Mathematics, Vidya Jyothi Institute of Technology, Hyderabad-500 070*
[3] *Department of Mathematics, Osmania University, Hyderabad, Telangana – 500 007*

Abstract: An analysis of MHD effects on unsteady free convection and mass transfer over a moving isothermal vertical plate in a rotating fluid in the presence of thermal radiation and viscous dissipation with Ohmic heating were presented. An exact solution is obtained for the axial and transverse components of the velocity by defining a complex velocity. The effects of velocity, temperature and concentration for different parameters like radiation parameter, rotation parameter, magnetic field parameter, Schmidt number, thermal Grashof number, mass Grashof number, Prandtl number and time on the plate are discussed.

Keywords: Magnetic field, Radiation, rotation, mass transfer, vertical plate, heat source.

I. INTRODUCTION

The study of hydromagnetic flow is called hydromagnetics or magnetohydrodynamics (MHD), which studies the dynamics of electrically conducting fluids. The set of equations which describe MHD are a combination of the Navier-Stokes equation of fluid dynamics and Maxwell's equations of electromagnetism. Radiative heat and mass transfer play an important role in manufacturing industries for the design of reliable equipment, Nuclear power plants, gas turbines and various propulsion devices for aircraft, missiles, satellites and space vehicles. The effect of Coriolis force has wide applications in science and technology. Viskanta and Grosh [15] studied the transfer of energy in boundary layer flow of an incompressible and radiating medium over wedge. The Rosseland approximation for the radiant heat flux vector was used to simplify the energy equation. Arpaci [22] studied the interaction between thermal radiation and laminar convection of heated vertical plate in a stagnant radiating gas. Greenspan [9] discussed the theory of rotating fluids owing to its numerous applications in cosmical and geophysical fluid dynamics, meteorology and engineering. England and Emery [23] have presented a correlation between the analytical solution and the experimental results of the convection-radiation interaction upon a vertical flat plate for absorbing and non absorbing gases. The flow past a horizontal plate has been studied by Debnath [10, 11], Puri and Kulshrestha [13]. Soundalgekar et al [20] presented an exact solution of flow past an impulsively started isothermal vertical plate under the action of transversely applied magnetic field. Again Soundalgekar et al [21] studied Stokes' problem for an infinite vertical plate whose temperature varies with time in the presence of transverse magnetic field for an incompressible fluid. Srinathuni Lavanya and Chenna Kesavaiah [17] considered heat transfer to MHD free convection flow of a viscoelastic dusty gas through a porous medium with chemical reaction, Srinathuni Lavanya and Chenna Kesavaiah [18] have radiation effects on MHD natural convection heat transfer flow from spirally enhanced wavy channel through a porous medium.

Many research works have been done based on the action of a uniform transverse magnetic field either fixed to the fluid or to the plate. Heat and mass transfer on MHD flows have applications in Meteorology, solar physics, cosmic fluid dynamics, astrophysics and geophysics. Magneto convection plays an important role in various industrial applications such as magnetic control of molten iron flow in the steel industry, liquid metal cooling in nuclear reactors, electromagnetic pumps, controlled fusion research, crystal growing, MHD couples and bearings, plasma jets, chemical synthesis and underground nuclear waste storage sites. Singh [1] studied the effects of coriolis as well as magnetic force on the flow field of an electrically conducting fluid past an impulsively started infinite vertical plate. Bestman and Adjepong [2] studied the magnetohydrodynamic free convection flow,

©IJRASET: All Rights are Reserved

with radiative heat transfer, past an infinite moving plate in rotating incompressible, viscous and optically transparent medium. Das et al. [19] have analyzed radiation effects on flow past an impulsively started infinite isothermal vertical plate. An analysis of the unsteady magneto hydrodynamic flow of a viscous and electrically conducting fluid past a vertical flat plate by the presence of radiation was done by Raptis and Massalas [5]. Basant Kumar Jha [6] presented the effects of uniform transverse magnetic field on the transient free convective flow of an electrically conducting fluid in a vertical channel. Raptis and Perdikis [3] considered the effects of radiation and free convection flow past a moving vertical plate. Saravanan and Kandaswamy [16] have analyzed the effect of temperature dependent thermal conductivity on buoyancy induced convection in the presence of a uniform magnetic field. It was inferred that it is advantageous to use low Prandtl number liquid metals or alloys as coolants in fast reactors whose thermal conductivity does not depend much on the temperature. Raptis and Perdikis [4] investigated free convection and mass transfer effects on optically thin gray gas past an infinite moving vertical plate. The governing equations were solved analytically. Cai and Zhang [14] studied the heat and mass transfer effects of an infinite vertical plate. Cavus and Karafistan [8] studied the effects of differential rotation in the lower convective region of the Sun, Ch Kesavaiah et.al. [7] has effects of the chemical reaction and radiation absorption on an unsteady MHD convective heat and mass transfer flow past a semi-infinite vertical permeable moving plate embedded in a porous medium with heat source and suction, Bhavana et.al [12] investigated the Soret effect on free convective unsteady MHD flow over a vertical plate with heat source.
However, MHD effects on a moving infinite vertical plate in a rotating fluid with mass transfer in the presence of thermal radiation is not studied in the literature. It is proposed to study MHD effects on flow past an impulsively started infinite isothermal vertical plate with uniform mass diffusion in a rotating fluid in the presence of thermal radiation and viscous dissipation with Ohmic heating. The dimensionless governing equations are solved by perturbation technique.

A. Mathematical Model

Three dimensional flow of a viscous, incompressible, electrically conducting fluid past an impulsively started infinite vertical isothermal plate with uniform mass diffusion in a rotating fluid [3, 6] is considered. On this plate, the x′-axis is taken along the plate in the vertically upward direction and the y′-axis is taken normal to x′-axis in the plane of the plate and z′-axis is normal to it. Both the fluid and the plate are in a state of rigid rotation with uniform angular velocity Ω′ about the z′-axis. The fluid considered here is a gray, absorbing-emitting radiation but a non-scattering medium. A transverse magnetic field B_0 of uniform strength is applied normal to the plate in the z′ direction. The induced magnetic field and viscous dissipation is assumed to be negligible. Initially, the plate and fluid are at rest with the temperature T'_∞ and concentration C'_∞ everywhere. At time t′ > 0, the plate is given an impulsive motion in the vertical direction against gravitational field with constant velocity u_0 in a fluid, in the presence of thermal radiation. At the same time the plate temperature is raised to T'_w and the concentration to T'_w, which are there after maintained constant. Since the plate occupying the plane z′ = 0 is of infinite extent, all the physical quantities depend only on z′ and t′. Then by Boussinesq's approximation, the unsteady flow is governed by the following equations:




Figure (1): Physical model of the problem

$$\frac{\partial u'}{\partial t'} - 2\Omega' v' = g\beta(T' - T'_\infty) + g\beta^*(C' - C'_\infty) + \nu\frac{\partial^2 u'}{\partial z'^2} - \sigma\frac{B_0^2}{\rho}u' \qquad (1)$$

©IJRASET: All Rights are Reserved

$$\frac{\partial v'}{\partial t'} + 2\Omega' u' = \nu \frac{\partial^2 v'}{\partial z'^2} - \frac{\sigma B_0^2}{\rho} v' \qquad (2)$$

$$\rho C_p \frac{\partial T'}{\partial t'} = k \frac{\partial^2 T'}{\partial z'^2} - \frac{\partial q_r}{\partial z'} - Q_0 (T' - T_\infty') + \sigma B_0^2 u'^2 \qquad (3)$$

$$\frac{\partial C'}{\partial t'} = D \frac{\partial^2 C'}{\partial z'^2} - Kr'(C' - C_\infty') \qquad (4)$$

The term $\frac{\partial q_r}{\partial z'}$ represents the change in the radiative flux with distance normal to the plate with the following initial and boundary conditions:

$$\begin{aligned} t' \le 0: \quad & u' = 0, \quad T' = T_\infty', \quad C' = C_\infty' \quad for all z' \\ t' > 0: \quad & u' = u_0, \quad T' = T_w', \quad C' = C_w' \quad at \quad z' = 0 \\ & u = 0, \quad T \to T_\infty, \quad C' \to C_\infty' \quad as \quad z' \to \infty. \end{aligned} \qquad (5)$$

By Rosseland approximation [1, 13], radiative heat flux of an optically thin gray gas is expressed by

$$\frac{\partial q_r}{\partial z'} = -4a^* \sigma (T_\infty'^4 - T'^4) \qquad (6)$$

It is assume that the temperature differences within the flow are sufficiently small such that T'^4 may be expressed as a linear function of the temperature. This is accomplished by expanding T'^4 in a Taylor series about T_∞' and neglecting higher-order terms, thus

$$T'^4 \cong 4T_\infty'^3 T' - 3T_\infty'^4 \qquad (7)$$

By using equations (6) and (7), equation (3) reduces to

$$\rho C_p \frac{\partial T'}{\partial t'} = k \frac{\partial^2 T'}{\partial y'^2} + 16a^* \sigma T_\infty'^3 (T_\infty' - T') - Q_0 (T' - T_\infty') + \sigma B_0^2 u'^2 \qquad (8)$$

On introducing the following dimensionless quantities:

$$(u, v) = \frac{(u', v')}{u_0}, \quad t = \frac{t' u_0^2}{\nu}, \quad z = \frac{z' u_0}{\nu}, \quad \theta = \frac{T' - T_\infty'}{T_w' - T_\infty'}, C = \frac{C' - C_\infty'}{C_w' - C_\infty'}$$

$$Gr = \frac{g\beta \nu (T_w' - T_\infty')}{u_0^3}, \quad Sc = \frac{\nu}{D}, Kr = \frac{\nu Kr'}{u_0^2}, \quad Gc = \frac{\nu g \beta^* (C_w' - c_\infty')}{u_0^3} \qquad (9)$$

$$M = \frac{\sigma B_0^2 u_0}{\rho}, \quad Pr = \frac{\mu C_p}{k}, \quad \Omega = \frac{\Omega' \nu}{u_0^2}, \quad R = \frac{16a^* \nu^2 \sigma T_\infty'^3}{k u_0^2}, Q = \frac{Q_0}{u_0^2 k}$$

and the complex velocity q = u + iv, $i = \sqrt{-1}$ in equations (1) to (5), the equations relevant to the problem reduces to

$$\frac{\partial q}{\partial t} + 2i\Omega i = Gr\,\theta + Gc\,C + \frac{\partial^2 q}{\partial z^2} - Mq, \qquad (10)$$

$$\frac{\partial \theta}{\partial t} = \frac{1}{Pr} \frac{\partial^2 \theta}{\partial z^2} - \frac{1}{Pr}(R+Q)\theta \qquad (11)$$

$$\frac{\partial C}{\partial t} = \frac{1}{Sc} \frac{\partial^2 C}{\partial z^2} - Kr\,C \qquad (12)$$

where magnetic field parameter (M), rotation parameter (Ω), radiation parameter (R), Schmidt number (Sc), heat source parameter (Q), chemical reaction parameter (Kr), thermal Grashof number (Gr) and mass Grashof number (Gm)

©IJRASET: All Rights are Reserved

The initial and boundary conditions in non-dimensional form are

$$\begin{aligned} & q = 0, \quad \theta = 0, \quad C = 0, \quad \text{forall} \quad z \le 0 \ \ \& \ \ t \le 0 \\ t > 0: \quad & q = 1, \quad \theta = 1, \quad C = 1, \quad \text{at} \quad z = 0 \\ & q = 0, \quad \theta \to 0, \quad C \to 0, \quad \text{as} \quad z \to \infty \end{aligned} \tag{13}$$

All the physical variables are defined in the nomenclature.

B. Method Of Solution

Equation (10) – (12) are coupled, non – linear partial differential equations and these cannot be solved in closed – form using the initial and boundary conditions (13). However, these equations can be reduced to a set of ordinary differential equations, which can be solved analytically. This can be done by representing the velocity, temperature and concentration of the fluid in the neighbourhood of the fluid in the neighbourhood of the plate as

$$q(z,t) = q_0(z) + \varepsilon e^{nt} q_1(z) + O(\varepsilon^2) \tag{14}$$

$$\theta(z,t) = \theta_0(z) + \varepsilon e^{nt} \theta_1(z) + O(\varepsilon^2) \tag{15}$$

$$C(z,t) = C_0(z) + \varepsilon e^{nt} C_1(z) + O(\varepsilon^2) \tag{16}$$

Substituting (13) in Equation (9) – (11) and equating the harmonic and non – harmonic terms, and neglecting the higher order terms of $O(\varepsilon^2)$, we obtain

$$q_0'' - M q_0 = -Gr\,\theta_0 - Gc\,C_0 - 2\Omega \tag{17}$$

$$q_1'' - (M + n) q_1 = -Gr\,\theta_1 - Gc\,C_1 \tag{18}$$

$$\theta_0'' - (R + Q)\theta_0 = 0 \tag{19}$$

$$\theta_1'' - (R + Q + n\,\mathrm{Pr})\theta_1 = 0 \tag{20}$$

$$C_0'' - Kr\,Sc\,C_0 = 0 \tag{21}$$

$$C_1'' - (Kr + n)\,Sc\,C_1 = 0 \tag{22}$$

The corresponding boundary conditions can be written as

$$\begin{aligned} & q = 0, \quad \theta = 0, \quad C = 0, \quad \text{forall} \quad z \le 0 \quad \& \quad t \le 0 \\ t > 0: \quad & q_0 = 1, \ \theta_0 = 1, \ C_0 = 1, \ q_1 = 0, \ \theta_1 = 0, \ C_1 = 0 \quad \text{at} \quad z = 0 \\ & q_0 \to 0, \theta_0 \to 0, C_0 \to 0, q_1 \to 0, \theta_1 \to 0, C_1 \to 0 \quad \text{as} \quad z \to \infty \end{aligned} \tag{23}$$

Solving Equations (17) - (22) under the boundary conditions (23) we obtain the velocity, temperature and concentration distributions in the boundary layer as

$$q_0 = A_1 e^{m_6 z} + A_2 e^{m_2 z} + A_3 + A_4 e^{m_8 z}$$

$$\theta_0 = e^{m_6 z}$$

$$C_0 = e^{m_2 z}$$

$$q_1 = 0, \theta_1 = 0, C_1 = 0$$

In view of the above equations (14) – (16) becomes

$$q(z,t) = A_1 e^{m_6 z} + A_2 e^{m_2 z} + A_3 + A_4 e^{m_8 z}$$

$$\theta(z,t) = e^{m_6 z}$$

$$C(z,t) = e^{m_2 z}$$

©IJRASET: All Rights are Reserved

C. Appendix

$$m_2 = -\sqrt{Kr\,Sc}\;\; m_4 = -\sqrt{(Kr+n)\,Sc},$$

$$m_6 = -\sqrt{R+Q}, m_8 = -\sqrt{M}, A_1 = -\frac{Gr}{m_6^2+M}$$

$$A_2 = -\frac{Gc}{m_2^2+M}, A_3 = -\frac{2\Omega}{M}, A_4 = (1-A_1-A_2-A_3)$$

Using equation (16) we get the following expression for skin-friction components τ_x and τ_y

$$\tau_x + i\tau_y = \left(\frac{\partial q}{\partial z}\right)_{z=0} = A_1 m_6 + A_2 m_2 + A_4 m_8$$

In equation (16), (17) the argument of the complementary error function and error function is complex. Hence in order to obtain the u and v components of the velocity and skin-friction.

II. RESULTS AND DISCUSSION

Using the above formula, expressions for u are obtained but they are omitted here to save the space. In order to get a physical view of the problem, these expressions are used to obtain the numerical values of velocity (u) for different values of the various parameter like Magnetic field parameter (M), rotation, Radiation parameter (R), Schmidt number (Sc), Heat Source parameter (Q), Chemical reaction (Kr), thermal Grashof number (Gr) and mass Grashof number (Gm)

A. Velocity profiles

Effects of radiation and rotation parameter on primary velocity; the primary velocity profiles of air for different values of the radiation parameter $(R = 0.5, 1.0, 1.5, 2.0)$ and rotation parameter $(\Omega = 0.1, 0.2, 0.3, 0.4)$ are shown in figure (2). It is observed that from figure 2 (a) can be observed in the primary velocity decreases with increasing the radiation parameter (R) and with increasing the rotation parameter Ω the primary velocity decreases in cooling of the plate observed in figure 2 (b). This shows that primary velocity decreases in the presence of high thermal rotation. Effects of magnetic field and rotation parameter on primary velocity the primary velocity profiles of air for different values of the magnetic parameter $(M = 0.5, 1.0, 1.5, 2.0)$ are shown in figure (3). It is observed that the primary velocity decreases with increasing magnetic parameter This shows that primary velocity decreases in the presence of high magnetic field and rotation. In fact rotation has more influence than magnetic field on primary velocity. Effects of thermal and mass Grashof number on primary velocity the primary velocity profiles for different thermal Grashof number $(Gr = 1.0, 2.0, 3.0, 4.0)$, mass Grashof number $(Gc = 1.0, 2.0, 3.0, 4.0)$ are shown in figure 4(a) and 4 (b). It is clear that the primary velocity increases with increasing thermal Grashof number or mass Grashof number. Figure (5) shows that primary velocity profiles for different values of chemical reaction parameter (Kr) it is clear that an increases chemical reaction parameter the velocity profiles decreases. The velocity profiles directed for various values of Schmidt number shown in figure (6), it is observed that increases in Schmidt number the result are decreases.

B. Temperature Distribution

The temperature profiles for different values of heat source parameter (Q = 0.5, 1, 1.5, 2) are shown in figure 7(a). The effect of thermal radiation parameter is important in temperature profiles. It is observed that the temperature decreases with increasing heat source parameter. The temperature profiles for air (Pr = 0.71) are calculated for different values of thermal radiation parameter (R = 0.2, 2, 10) are shown in figure 7(b). The effect of thermal radiation parameter is important in temperature profiles. It is observed that the temperature decreases with increasing radiation parameter.

©IJRASET: All Rights are Reserved

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 6.887
Volume 6 Issue V, May 2018- Available at www.ijraset.com

C. Concentration Profiles

Variations in the concentration profiles for different values of the chemical reaction parameter and Schmidt number are shown from figure 8(a) and 8(b); form these figures it is clear that an increasing values opf chemical reaction parameter and Schmdit number the concentration profiles decreases.

D. Skin friction

The skin-friction components for different parameter are calculated using Equation (16). The skin-friction components at the wall for different values of $Gc = 5.0,\ R = 2.0,\ M = 1.0,\ Kr = 1.0,\ n = 0.1,\ Q = 1.0$ and Ω are shown in table (1)

Table (1): Skin-friction (τ) for different values of Ω versus Gr

Ω	Gr	τ_x
0.2	1.0	4.69
0.4	2.0	4.29
0.6	3.0	3.89
0.8	4.0	3.49

The effect of rotation on skin-friction decreases the component τ_x as Ω increases. As time advances the component τ_x increases. Greater cooling of the plate, due to free-convection currents, lower rises τ_x.

Using equation (15) we get the following expression for Nusselet number

$$N_u = \left(\frac{\partial \theta}{\partial z}\right)_{z=0} = m_6$$

Table (2): Nusselt number for different values of R

R	t	N_u
0.2	0.2	1.1806
2.0		2.0693
5.0		3.0905
	0.4	2.9483

Nusselt number for different thermal radiation and time are calculated using equation (15), table (2) shows that as radiation increases Nusselt number increases, but the trend is reversed with time.

E. Sherwood Number

Using equation (14) we get the following expression for Sherwood number

$$S_h = \left(\frac{\partial C}{\partial z}\right)_{z=0} = m_2$$

Table (3): Sherwood number (Sh) for different values of Sc

Sc	Kr	Sh
0.5	1.0	0.00349300
1.0	2.0	0.00033550
1.5	3.0	$5.556\,e^{-005}$
2.0	4.0	$1.22\,e^{-005}$

©IJRASET: All Rights are Reserved

Sherwood numbers for different Schmidt number and time are calculated using equation (16), table (2) shows that as Schmidt number increases Sherwood number increases, but the trend is reversed with time.

III. CONCLUSIONS

Theoretical analysis is performed to study magneto hydrodynamic effects on flow past an impulsively started infinite isothermal vertical plate with uniform mass diffusion, in the presence of thermal radiation in a rotating fluid. The dimensionless governing equations are solved by Laplace-transform technique. The conclusions of the study are as follows:

A. Concentration falls with the increase in Schmidt number.
B. Temperature is enhanced with the decreasing radiation parameter and increasing time.
C. The thermal radiation or rotation parameter on primary flow has a retarding effect for cooling of the plate.
D. The influence of rotation on primary flow is significant than magnetic field, but both has a retarding effect for cooling of the plate.
E. The secondary velocity is enhanced with the raise in thermal radiation or magnetic field parameter and opposite phenomenon occurs with the rotation parameter.
F. The effect of thermal Grashof number or mass Grashof number enhanced the primary flow but decreased the secondary flow.
G. The skin-friction components falls with increasing radiation parameter, thermal Grashof number and mass Grashof number on τ_x rises on τ_y. But reverse phenomenon occurs with rotation parameter or magnetic parameter.
H. Sherwood number increases with increasing Schmidt number and the Nusselt number increases with increasing radiation parameter but both shows reverse trend with respect to time.

REFERENCES

[1] A K Singh (1984): Hydromagnetic free convection flow past an impulsively started vertical infinite plate in a rotating fluid, Int. Comm. Heat Mass Transfer, 11 (1984), pp. 399–406.
[2] A R Bestman and S K Adjepong (1988): Unsteady hydromagnetic free-convection flow with radiative heat transfer in a rotating fluid, Astrophys. Space Sci., Vol. 143, (1988), pp. 73–80.
[3] A Raptis and C Perdikis (1999): Radiation and free convection flow past a moving plate, Int. J. Appl. Mech. and Engg, 4 (1999), pp. 817–821.
[4] A Raptis and C Perdikis (2003): Thermal Radiation of an optically thin gray gas, Int. J. Appl. Mech. and Engg., Vol. 8 (2003), pp. 131–134.
[5] A Raptis and C V Massalas (1998): Magnetohydrodynamic flow past a plate by the presence of radiation, Int. J. Heat Mass Transfer., 34 (1998), pp.107–109.
[6] Basant Kumar Jha (1998): Effects of applied magnetic field on transient free-convective flow in a vertical channel, Indian J. Pure appl. Math., Vol. 29 (1998), pp. 441–445.
[7] H Cavus and A I Karafistan (2010): On the modeling of rotational effects in the lower convective region of the sun, Braz. J. Phys., Vol. 40 (2010) 2.
[8] H P Greenspan (1968): The theory of rotating fluids. Cambridge University Press, UK, 1968.
[9] L Debnath (1972): On unsteady magnetichydrodynamics boundary layers in a rotating flow, Z. Angew. Math. Mech, Vol. 52 (1972), pp. 623–626.
[10] L Debnath (1975): Exact solution of the unsteady hydrodynamic and hydromagnetic boundary layer equations in a rotating fluid system, Z. Angew. Math. Mech., Vol. 55 (1975), pp. 431–438.
[11] M Bhavana, D Chenna Kesavaiah and A Sudhakaraiah (2013): The Soret effect on free convective unsteady MHD flow over a vertical plate with heat source, International Journal of Innovative Research in Science, Engineering and Technology, Vol. 2 (5), pp. 1617-1628
[12] P Puri and P K Kulshrestha (1976): Unsteady hydro-magnetic boundary layer in a rotating medium, J. Appl. Mech. Trans. ASME., 98 (1976), pp. 205–208.
[13] R Cai and N Zhang (2003): Explicit analytical solutions of linear and non linear interior heat and mass transfer equation sets for drying process, ASME TRANS J. Heat Transfer, Vol. 125 (2003), pp.175–178.
[14] R Viskanta and R J Grosh (1962): Boundary layer in thermal radiation absorbing and emitting media, Int. J. Heat Mass Transfer, 5 (1962), pp.795–805.
[15] S Saravanan and P Kandaswamy (2000): Natural convection in low Prandtl number fluids with a vertical magnetic field, Trans. ASME J. Heat Transfer, 122 (2000), pp. 602–606.
[16] Srinathuni Lavanya and D Chenna Kesavaiah (2017): Heat transfer to MHD free convection flow of a viscoelastic dusty gas through a porous medium with chemical reaction, International Journal of Pure and Applied Researches, Vol. 3 (1), pp. 43 - 56
[17] Srinathuni Lavanya and D Chenna Kesavaiah (2017): Radiation effects on MHD natural convection heat transfer flow from spirally enhanced wavy channel through a porous medium, International Journal on Future Revolution in Computer Science & Communication Engineering ,Vol. 3(10), pp. 130-140
[18] U N Das, R K Deka and V M Soundalgekar (1996): Radiation effects on flow past an impulsively started vertical infinite plate, J. Theo. Mech, Vol. 1 (1996), pp. 111–115.
[19] V M Soundalgekar (1979): Effects of mass transfer and free-convection currents on the flow past an impulsively started vertical plate, Trans. ASME. J. Appl. Mech., 46 (1979), pp.757–760.
[20] V M Soundalgekar, S K Gupta and R N Aranake (1979): Free convection effects on MHD Stokes problem for a vertical plate, Nucl. Eng. Des., 51 (1979), pp. 403–407.
[21] V S Arpaci (1968): Effect of thermal radiation on the laminar free convection from a heated vertical plate, Int. Int. J. Heat Mass Transfer, Vol. 11 (1968), pp. 871–881.

©IJRASET: All Rights are Reserved

[22] W G England and A F Emery (1969): Thermal radiation effects on the laminar free convection boundary layer of an absorbing gas, J. Heat Transfer, Vol. 91, (1969), pp. 37–44.

Figures:




Figure 2(a): Primary velocity profiles for different values of R




Figure 2(b): Primary velocity profiles for different values of Ω




Figure 4 (a): Primary Velocity Profiles for different values of Gr




Figure 4 (b): Primary Velocity Profiles for different values of Gc




Figure (5): Primary Velocity profiles for different values of Kr




Figure (6): Primary velocity profiles for different values of Sc




Figure 7 (a): Temperature profiles for different values of Q




Figure 7(b): Temperature profiles for different values of R




Figure 8(a): Concentration profiles for different values of Kr





Figure 8(b): Concentration profiles for different values of Sc

©IJRASET: All Rights are Reserved



10.22214/IJRASET

45.98

IMPACT FACTOR: 7.129

IMPACT FACTOR: 7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)