



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 7 Issue: VII Month of publication: July 2019

DOI: <http://doi.org/10.22214/ijraset.2019.7127>

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

Three Connected Domination in a Graph

V. Praba

Assistant Professor, Department of Mathematics, SIGC, Trichy-620002, Tamilnadu, India.

Abstract: Claude Berge [1] introduced the concept of strong stable set S in a graph. These sets are independent and any vertex outside S can have at most one neighbour in S . This concept was generalized by E. Sampathkumar and L. Pushpalatha [5]. A maximal independent set is a minimal dominating set. What type of domination will result from maximal semi-strong sets? This new type of domination which we call us -Three-connected domination is initiated and studied in this paper.

Keywords: Strong stable set, Semi-strong set, Three-connected domination.

MSC: 05C69.

AMS Mathematics subject Classification (2010):11D09

I. INTRODUCTION

Let $G = (V, E)$ be a simple, finite, undirected graph. A subset S of $V(G)$ is called a strong stable set of G if $|N[v] \cap S| \leq 1$ for v in $V(G)$. It can be easily seen that such a set is independent and the distance between any two vertices of S greater than equal to three. That is, the strong stable set is a 2-packing. Generalising this concept, E. Sampathkumar and L. Pushpa Latha [5] introduced the concept of semi-strong sets. A subset S of $V(G)$ is called semi-strong stable if $|N(v) \cap S| \leq 1$ for every v in $V(G)$. A strong stable set is semi-strong stable but the converse is not true. For example, in C_5 , any two consecutive vertices is a semi-strong stable set. If S is a semi-strong stable set, then any component of S is either K_1 or K_2 and the distance between any two points of S is not equal to two. A maximal semi-strong stable set gives rise to a new type of domination and this is studied in this paper.

II. THREE-CONNECTED DOMINATING SET

- 1) **Definition 2.1:** Let S be a subset of $V(G)$. For any $u \in V - S$, if there exists $v \in V(G)$, $v \neq u$ such that v is adjacent with u and v is adjacent with a vertex of S , (that is, for any $u \in V(G)$ and $w \in S$ such that uvw is a path P_3), then S is called a 3-connected dominating set of G .
- 2) **Remark 2.2:** Any 3-connected dominating set S of G which is semi-strong is a maximal semi-strong set of G .
- 3) **Theorem 2.3:** Let S be a subset of $V(G)$ such that for any $u \in V - S$, there exists v and a vertex w in S such that uvw is a path. This property is super hereditary.

Proof

Let S be a subset of $V(G)$ satisfying the hypothesis. Let T be a proper super set of S . Let $u \in V - T$. Then $u \in V - S$. By hypothesis, there exists a vertex v and a vertex w in S such that uvw is a path.

- a) **Case 1:** $v \in V - T$. In this case, $u, v \in V - T$ and $w \in T$ (since $w \in S \subset T$). Moreover uvw is a path.
- b) **Case 2:** $v \in T - S$ and $u \in V - T$. There exist w in S such that uvw is a path. That is, $u \in V - T$, $v \in T$, $w \in T$ and uvw is a path.
- c) **Case 3:** $v \in S$ and $u \in V - T$. There exist $w \in S$ such that uvw is a path. That is, $v \in T$ and $w \in T$ and uvw is a path. In all the three cases, for any $u \in V - T$, there exist $v \in V(G)$, $v \neq u$ and $w \in T$ such that uvw is a path. Therefore the property for maximality of a semi-strong set S is super hereditary.
- 4) **Remark 2.4:** The above property is called a 3-connected dominating property.
- 5) **Theorem 2.5:** Any minimal 3-connected dominating set is a maximal semi-strong set.

Proof

Let S be a minimal 3-connected dominating set of G .

- a) **Case 1:** Let $u \in V - S$

- i) **Subcase 1:** There exists $v \in V - S$ and $w \in S$ such that uvw is a path. Suppose u has at least two neighbours in S . Let $x, y \in S$ such that u is adjacent with x and y .
 1. Consider $S - \{x\}$. For any u_1 in $V - (S - \{x\})$, $u_1 \neq x$, $u_1 \in V - S$. There exists v in $V(G)$, $v \neq u_1$ and w in S such that uvw is a path if $w = x$. Then u_1vw is a triangle and not a path, contradiction. Therefore $w \neq x$. Therefore $w \in S - \{x\}$. Therefore there exists $w \in (S - \{x\})$ such that u_1vw is a path.
 2. Suppose $u_1 = x$. Then $u \in V - S$ such that u is adjacent with x and adjacent with $y \in (S - \{x\})$. That is, u_1 is adjacent with u and u is adjacent with $y \in (S - \{x\})$. Therefore $S - \{x\}$ is a 3-connected dominating set of G , a contradiction (since S is minimal).

ii) Subcase 2: There exist $v, w \in S$ such that uvw is a path.

1. Suppose u has at least two neighbours say v, x in S . Let $u_1 \in V - (S - \{x\})$.

2. Suppose $u_1 \neq x$. Therefore $u_1 \in V - S$. Hence there exists v in $V(G)$ and w in S such that u_1vw is a path. If $w = x$, then u_1vx is a triangle, a contradiction. Therefore $w \neq x$. Therefore $w \in S - \{x\}$ and uvw is a path.

3. Suppose $u_1 = x$. In this case u_1 is adjacent with $u \in V - S$ and u is adjacent with $v \in (S - \{x\})$. Also u_1uv is a path. Therefore $S - \{x\}$ is a 3-connected dominating set, a contradiction since S is minimal. Therefore $|N(u) \cap S| \neq 1$.

b) Case 2: $u \in S$, Suppose u has at least two neighbours say x, y in S . Consider $S - \{x\}$. Then $x \in V - (S - \{x\})$. x is adjacent with $u \in V(G)$ and u is adjacent with $y \in S - \{x\}$. Therefore xuy is a path. Therefore $S - \{x\}$ is a 3-connected dominating set of $V(G)$, a contradiction. Therefore for any u in S , $|N(u) \cap S| \leq 1$. Hence S is a semi-strong set of G . Since S is a 3-connected dominating set of G and since S is semi-strong set of G , we get that S is a maximal semi-strong set of G .

6) Theorem 2.6: Any maximal semi-strong set of G is a minimal 3-connected dominating set of G .

Proof

Suppose S is a maximal semi-strong set of G . Then S is a 3-connected dominating set of G . Suppose S is not a minimal 3-connected dominating set of G . Therefore there exists a proper subset T of S such that T is a 3-connected dominating set of G . Since S is semi-strong, T is semi-strong. Therefore T is a maximal semi-strong set of G which satisfies 3-connected property. Therefore T is a maximal semi-strong set of G , a contradiction, since S is a proper superset of T and S is a semi-strong set of G . Therefore S is a minimal 3-connected dominating set of G .

7) Definition 2.7: The minimum (maximum) cardinality of a minimal 3-connected dominating set of G is called 3-connected domination number of G (upper 3-connected domination number of G) and is denoted by $\gamma_{3-c}(G)$ ($\Gamma_{3-c}(G)$).

8) Remark 2.8: Let S be a minimum cardinality of a maximal semi-strong set of G . Then S is a minimal 3-connected dominating set of G . Therefore $\gamma_{3-c}(G) \leq |S| = lss(G) \leq ss(G)$.

9) Remark 2.9: Let S be a maximum semi-strong set of G . Therefore S is a minimal 3-connected dominating set of G . Therefore $ss(G) = |S| \leq \Gamma_{3-c}(G)$. Therefore $\gamma_{3-c}(G) \leq lss(G) \leq ss(G) \leq \Gamma_{3-c}(G)$.

10) Illustration 2.10: Let G be the graph given in Figure 1:

In this graph, $S_1 = \{u_1, u_2, u_5, u_7, u_8, u_{11}\}$ is a ss -set of G . Hence $ss(G) = 6$. $S_2 = \{u_3, u_6, u_7, u_{11}\}$ is a maximal semi-strong set of G of minimum cardinality. Therefore $lss(G) = 4$. $S_3 = \{u_3, u_6, u_9\}$ is a minimum 3-connected dominating set of G .

Hence $\gamma_{3-c}(G) = 3 \leq lss(G) = 4$. That is, $\gamma_{3-c}(G) < lss(G)$.




Figure 1: An example graph G for $\gamma_{3-c}(G) < lss(G)$

11) *Theorem 2.11:* Let S be a 3-connected dominating set of G . S is minimal if and only if for any w in S there exists a vertex u in $V - S$ such that any 3-connected path from u to S ends in w .

Proof

Let S be a minimal 3-connected dominating set of G . Let $w \in S$. Then $S - \{w\}$ is not a 3-connected dominating set of G . Therefore there exists u in $V - (S - \{w\})$ such that there is no 3-connected path uv_1w_1 where $v_1 \in V(G)$ and $w_1 \in S - \{w\}$. Since S is a 3-connected dominating set of G , there exists $v_1 \in V(G)$ and $w_1 \in S$ such that uv_1w_1 is path. If $w_1 \neq w$, then there exists a 3-connected path uv_1w_1 from u to $S - \{w\}$, a contradiction. Therefore $w_1 = w$. Therefore any 3-connected path from u to S is of the form uvw . That is, there exists u in $V - S$ such that any 3-connected path from u to S ends in w .

Conversely, let S be a 3-connected dominating set of G such that for any w in S , there exists u in $V - S$ such that 3-connected path from u to S ends in w .

1) *Claim:* $S - \{w\}$ is not a 3-connected dominating set for any w in S .

Since S is a 3-connected dominating set of G satisfying the above property, there exists u in $V - S$ such that any 3-connected path from u to S must end in w . Therefore $u \in V - (S - \{w\})$, $u \neq v$. Suppose there exists a 3-connected path from u to $S - \{w\}$ say uvw_1 , where $w_1 \in S - \{w\}$. Then $w_1 \in S$ and uvw_1 is a path ending in w_1 in S , $w_1 \neq w$, a contradiction. Therefore $S - \{w\}$ is not a 3-connected dominating set of G . Hence the claim.

Therefore S is a minimal 3-connected dominating set of G .

III. THREE-CONNECTED PATH IRREDUNDANCE

1) *Definition 3.1:* Let S be a subset of $V(G)$ such that for any w in S , there exists a u in $V - S$ such that any 3-connected path from u to S ends in w . Then S is called a 3-connected path irredundant set of G .

2) *Theorem 3.2:* The above property of a set S is hereditary.

Proof

Let S be a subset of $V(G)$ satisfying the above property. Let T be a proper subset of S .

Let $w \in T$. Then $w \in S$. Therefore there exist $u \in V - S$ such that any 3-connected path from u to S ends in w . Therefore $u \in V - T$. Suppose there exists a 3-connected path such that $w_1 \in T$,

$w \neq w_1$. Then $w_1 \in S$. Therefore there exists a 3-connected path from u to w_1 in S , a contradiction. Therefore $w_1 = w$. Hence T is a subset of $V(G)$ satisfying the above property. Hence the theorem.

3) *Definition 3.3:* Let S be a 3-connected path set of G . The minimum (maximum) cardinality of a maximal 3-connected path irredundant set of G is called 3-connected path irredundant number of G (upper 3-connected path irredundant number of G) is denoted by $ir_{3-c}(G)$ ($IR_{3-c}(G)$).

4) *Remark 3.4:* Any 3-consecutive dominating set of G is minimal if and only if it is a 3-consecutive path irredundant set of G .

5) *Theorem 3.5:* Every minimal 3-connected dominating set of G is a maximal 3-connected path irredundant set of G .

Proof

Let S be a minimal 3-connected dominating set of G . Then S satisfies the property that for every w in S , there exists u in $V - S$ such that any 3-connected path from u to S ends in w . Therefore S is a 3-connected path irredundant set of G . Suppose S is not a maximal 3-connected path irredundant set of G .




Figure 2: An example graph G for which $ir_{3-c}(G) < \gamma_{3-c}(G)$

Since 3-connected path irredundant is hereditary, it is enough to consider 1-maximality. Since S is not maximal, there exists u in $(V - S)$ such that $S \cup \{u\}$ is 3-connected path irredundant set of G . Therefore for any x in $S \cup \{u\}$, there exist y in $V - (S \cup \{u\})$ such that any 3-connected path from y in $S \cup \{u\}$ ends in x . Take $x = u$. Then there exists y in $V - (S \cup \{u\})$ such that any 3-connected path from y in $S \cup \{u\}$ ends in u . That is, there exists y in $V - S$ such that any 3-connected path from y to S does not end in any vertex of S , that is, S does not satisfy 3-connected path irredundant condition, a contradiction. Therefore S is a maximal 3-connected path irredundant set of G .

6) *Remark 3.6:* For any graph G , $ir_{3-C}(G) \leq \gamma_{3-C}(G) \leq lss(G) \leq ss(G) \leq \Gamma_{3-C}(G) \leq IR_{3-C}(G)$.

7) *Remark 3.7:* In the following example, $ir_{3-C}(G) < \gamma_{3-C}(G)$. Let G be the graph given in Figure 2. The set

$S_1 = \{u_2, u_4, u_6\}$ is a minimum 3-connected dominating set of G . Therefore $\gamma_{3-C}(G) = 3$.

The set $S_2 = \{u_3, u_5\}$ is maximum 3-connected path irredundant set of G . $ir_{3-C}(G) = 2$.

Therefore $ir_{3-C}(G) < \gamma_{3-C}(G)$

REFERENCES

- [1] C. Berge, Graphs and Hyper graphs, North Holland, Amsterdam, 1973.
- [2] F. Harary, Graph Theory, Addison-Wesley Reading, MA (1969).
- [3] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, Fundamentals of Domination in Graphs, Marcel Dekker Inc., 1998.
- [4] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, Domination in Graphs: Advanced topics, Marcel Dekker Inc., 1998.
- [5] E. Sampathkumar and L. Pushpa Latha, Semi-Strong Chromatic Number of a Graph, Indian Journal of Pure and Applied Mathematics, 26(1): 35-40, 1995.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)