



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 7 Issue: VII Month of publication: July 2019

DOI: <http://doi.org/10.22214/ijraset.2019.7212>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

A Study on Diseases Affecting *Litopenaeus vannamei* Farming in Coastal Districts of Andhra Pradesh, India

V. Venkateswarlu¹, P.V. Seshiah², P. C. Behra³

¹Department of Marine Biology, Vikrama Simhapuri University, Nellore

^{1, 2, 3}PVS Group, Vijayawada, Andhra Pradesh, India

Abstract: Andhra Pradesh stands top in the aquaculture production in India and shrimp farming is one of the major sources of income. Now-a-days shrimp culture is facing many problems among them disease mitigation became major hurdle for the shrimp farmers. A study was done to evaluate the diseases affecting *L.vannamei* farming in coastal districts of Andhrapradesh, i.e., Gudur, Krishna, West Godavari and East Godavari. The summer crop was considered for the study during February – June 2019. The diseases that are responsible for huge losses in *L. vannamei* culture was found to be White Spot Syndrome Virus (WSSV), White Faecal Syndrome (WFS), Black Gill Disease (BGD), Loose Shell Syndrome (LSS), Running Mortality Syndrome (RMS), and Enterocytozoon Hepatopenaei (EHP) of all the selected 4 districts of the Andhra Pradesh. About 65% culture ponds of Guntur district, 69% in Krishna and 40% in West Godavari and 50% in East Godavari districts were affected by diseases. It was observed that, more disease related problems observed during the summer crop in all selected areas. The farms which implemented biosecurity measures, and maintained without any Dissolved Oxygen (DO) problems, were found to contain less diseases. Therefore it is suggested that by practicing better management practices (BMP) by following biosecurity measures, diseases can be prevented.

Keywords: Shrimp farming, *L.vannamei*, diseases, Andhra Pradesh, BMP practices.

I. INTRODUCTION

Shrimp farming become small scale business in south East Asia as a global industry than other aquaculture farming systems^[1]. The impressive growth of shrimp farming bed a major player on global shrimp Industry, it play a virtual role on offering quality nutrition, employment and rural development. Andhra Pradesh has the second largest brackish water area in India after West Bengal, covering an area of about 37,560 ha. Andhra Pradesh has the coastline of 972 km, spreading over nine districts namely Srikakulam, Vijayanagaram, Visakhapatnam, East Godavari, West Godavari, Krishna, Guntur, Prakasam and Sri Potti Sri Ramulu Nellore. Development of coastal aquaculture in Andhra Pradesh is centered on shrimp (*L. vannamei*) farming. The culture area increased from 264 Ha to 37,560 Ha during the period from 2009-10 to 2014-15. Andhra Pradesh (2, 76,077 MT) is the leading farmed shrimp producer with 78% and the rest of India production was 77, 336 MT (MPEDA, 2015). The exports production of *Litopenaeus vannamei* increased from 18247 to over 400000 MT in the country during 2010-11 to 2017-18 (MPEDA.2015)^[2]. The shrimp aquaculture in India suffered significantly due to disease infections. In India, the gross economic losses due to shrimp diseases were estimated at more than Rs.1,000 crores in 2006-2008 and loss continues even now^[3]. The threat of the diseases continued due to poor practicing of biosecurity measures like Pond preparations, water filtration and disinfection before and after pumping water in to the pond, not arranging bird and crab fencing, procuring the seed from unregistered hatcheries which supply non SPF seed. The diseases may be caused by various etiological agents such as viruses, bacteria, fungi, parasites, algal toxins, nutritional deficiency or the adverse environment. The frequent outbreaks of diseases such as White Spot Syndrome Virus (WSSV), Black Gill Disease (BGD), Running Mortality Syndrome (RMS), Loose Shell Syndrome (LSS), White Faecal Syndrome (WFS), White Muscle Disease (WMD) and Infectious Hypodermal and Haematopoietic Necrosis (IHHN) in shrimps causing economic loss to the aquaculture industry^[4]. The present work was under taken to evaluate the disease out breaks in *L.vannamei* farming in Andhra Pradesh.




Fig.1 Andhra Pradesh map (Study area districts)

II. METHODOLOGY

The present study was conducted in four districts, viz, Guntur, Krishna, West Godavari and East Godavari of Andhra Pradesh during February to June 2019. The four districts were selected based on the farming practices of the shrimp *L.vannamei*. Stratified random data sampling procedures was adopted for the selection of farms and shrimp farmers. A total of 80 farms in each district were considered for the study.

S. No	Name of the Study area	No. of Villages covered	No. of Farms covered	Total culture area covered (Ha)
1	Guntur	8	80	189
2	Krishna	12	80	239
3	West Godavari	9	80	205
4	East Godavari	8	80	201

Table.1 Study area in 4 districts of Andhra Pradesh

The primary data were collected by survey method with the help of pre-tested schedule through personal interview to assess the disease impact of the shrimp farming in the selected study area.

Survey Method	Description of the Units Covered
Sample Survey	Farms registered with CAA, Farmers in the 37 villages.

Table.2. Impact assessment methods adopted by the study

III. RESULTS AND DISCUSSION

Six major diseases viz. White Spot Syndrome Virus (WSSV), White Faecal Syndrome (WFS), Black Gill Disease (BGD), Loose Shell Syndrome (LSS), Running Mortality Syndrome (RMS), and Enterocytozoon Hepatopenaei (EHP) were found in *L. vannamei* culture ponds of all the 4 divisions of the Andhra Pradesh. About 65% culture ponds of Guntur district, 69% in Krishna and 40% in West Godavari and 50% in East Godavari districts were affected by diseases. It was observed that, more disease related problems observed during the summer crop in all selected areas. It was observed that, only 20% of farms implemented bio-security measures in Guntur and Krishna districts and more diseases were observed in those districts. Whereas in East and West Godavari districts 40 % of farms implemented biosecurity measures, and maintained their farms without any Dissolved Oxygen (DO) problems, therefore less infections were observed in those districts.

All farmed brackish water and marine (penaeid) shrimp species are highly susceptible to white spot virus disease, with mass mortalities commonly reaching 80–100% in ponds within a period of 3–10 days^[5,6]. White Spot Syndrome Virus (WSSV) has a wide host range in crustaceans^[7,8,9] and is potentially becoming cause to most of the commercially cultivating shrimp species (OIE 2003). Bright milky colour White spots appear on the carapace and other parts of the body. Due to lack of bio-security at the farm, pumping water without disinfection and filtration, cross contamination were observed to be main reasons for the outbreak of WSSV (Fig. 2).



Fig 2: White Spots on carapace of the *L. vannamei*

White Faeces Syndrome (WFS) is observed in all the districts of *L. vannamei* culture ponds (Fig. 3). Incidences of WFS were observed after 20 days of stocking of the PLs in culture ponds. WFS in shrimp arises with protozoan Gregarines and Hepatopancreas infection caused by the high vibrio sps. Bacteria the Lipid cells leaching into the gut. White faecal material float on the water surface in the culture ponds. Observed *Vibrio* species have been found in the pond water and fecal analysis from infected shrimps^[11, 12].



Fig 3: White faecal matter in *L. vannamei* culture pond.

Black Gill Disease (BGD) was observed in Guntur and Krishna districts majority of the ponds, observed less in West Godavari and East Godavari districts, Poor pond management and Poor water quality is the main reason for Black Gill Disease (BGD) disease (Fig. 4). This disease is normal in shrimps. It is commonly caused by to abiotic factors in the pond like heavy organic load in pond bottom, presence of pollutants, low DO problems and high stocking densities. Observed that disease infected ponds was not managed properly it will affect the growth and survival of the shrimps in the ponds^[13]. Black gill Disease. In the initial stages appear light yellow or brown in colour and finally it leads to change as black in color^[14]. The black gill diseases of shrimp causes damage in grow out ponds during the end of the culture.



Fig 4: Shrimps infected with black gill disease.

Loose Shell Syndrome (LSS) was observed in majority of ponds in all the four districts i.e., Guntur, Krishna, West Godavari and East Godavari districts (Fig. 5). Previously studies also reported in the cultured *Penaeus monodon* since 1998 in India ^[15]. The affected shrimp are with soft and thin exoskeleton, lethargic in movement, spongy while touch, feed intake was poor and also observed a gap between the muscle and the shell. Moribund, weak shrimp with soft shell swim near dikes of the ponds appear in shiny light pink colour and results in progressive mortality. The mineral deficiency, poor Nutrition and water quality and high loads of pathogenic bacteria and other disease causes of WFS may cause the loose shell syndrome ^[16,17].



Fig 5: Loose Shell Syndrome (LSS) in *L. vannamei*.

Running Mortality Syndrome (RMS) was observed more in Guntur, Krishna districts and less in West Godavari and East Godavari (Fig. 6). The farmed shrimp in the affected ponds show continuous mortality with no symptom to any other diseases. In the initial stages of the RMS conditions shrimp show antennae cut, uropod turn pink or red in colour and later the Hepatopancreas begins will turn reddish yellow; finally entire body turns into red colour. Continuous internal mortality was noticed in the pond. The dead shrimp settle at the bottom of the pond and sometimes observed in check tray and not observed near the dikes and pond surface. Several farmers have lost survival and productivity in their crops. In the beginning, the farmers managed the situation by immediately removing the dead shrimps and following strict biosecurity as well as Better Management Practices (BMPs) by reducing the pond pollution. Farmers observed stocked low stocking densities were able to harvest the crop successfully without Running Mortality Syndrome RMS.




Fig 6: Running Mortality Syndrome in *L. vannamei*.

Enterocytozoon hepatopenaei (EHP), observed in Mainly in Krishna district, less in Guntur than Krishna district also observed equal in West Godavari and East Godavari district, EHP is a microsporidian caused by parasite to be associated with slow (retarded or stunted) growth and also cause white feces syndrome (WFS) in cultured shrimp in many of the shrimp farms observed in the Guntur, Krishna districts (Fig.7). That the EHP could be detected from slow growing as well as WFS-affected animals ^[18]. EHP is also reported from slow growing shrimp. Thus, EHP is an emerging problem that is under urgent need of control ^[19]. It is caused by Microsporidian fungi infected from the soil in hatcheries infected poly chaetes giving as a feed for brood stock.



Fig 7: EHP infected shrimp *L. vannamei* stock.

IV. CONCLUSION

The disease problems are becoming serious threat to the sustainability of shrimp aquaculture in Andhra Pradesh coastal districts. Aquaculture farmers should be aware about these diseases prevention and management of the diseases of shrimp and Health management. Farmers have to follow the Better management Practices (BMPs) with strong Biosecurity measures in all stages of the culture from pond preparation to harvesting, then only *L.vannamei* culture can be achieved without and diseases problems. The Better Management Practices (BMPs) like proper pond preparation, water and soil treatment with adopted reservoir system for water filling and also should purchase Specific Pathogen Free (SPF) quality shrimp seed from Coastal Aquaculture Authority registered hatcheries. During culture should follow the biosecurity measures viz., Farm materials disinfection, Foot and Hand dips, Shrimp toilets or sludge drains mainly Bird fencing and crab fencing to avoid disease transmission. If disease is found in pond that should be treat properly like water, Pond bottom, Animals with CAA registered Aquaculture Farm and Animal care products also farmers can adopt the practice weekly testing of the pond water and animals in Aquaculture Water Quality and Disease diagnosis Laboratory for monitoring the water quality changes, diseases symptoms during the culture period. With this practice farmers can easily identify the changes in water and animals, disease also can be prevented and control the disease possibility and can reduce the risk or loss of the crop from diseases.

V. ACKNOWLEDGEMENTS

The authors express deep thanks to the supported shrimp farmers, Technicians and other stakeholders of Guntur, Krishna, West Godavari and East Godavari districts in Andhra Pradesh.

REFERENCES

- [1] Joseph Selvin AS, Ninawe Lipton AP. Disease management (prospective approaches), Ane Books Pvt. Ltd. Thomson Press, India, 2009. ISBN (10): 81-9083-229-8.
- [2] MPEDA, 2015 and 2017. Annual Report 2014-2015, & 2017 Marine Products Export Development Authority, Cochin.
- [3] Kalaimani, N., Ravisankar, T., Chakravarthy, N., Raja, S., Santiago, T.C. and Ponniah, A.G. 2013. Economic Losses due to Disease Incidences in Shrimp Farms of India. *Fish. Techn.* 50: 80-86.
- [4] Srinivas, D. and Venkatrayulu, Ch. 2016. Current Status and Prospects of Pacific White Shrimp *Litopenaeus vannamei* (Boone, 1931) Farming in Coastal Districts of Andhra Pradesh in India. *International Journal of Science and Research*. 5(3): 1211-1214.
- [5] Chou HY, Huang CY, Wang CH, Chiang HC, Lo CF. Pathogenicity of a baculovirus infection causing white spot syndrome in cultured penaeid shrimp in Taiwan. *Dis. Aquat. Org.* 1995; 23:165-173.
- [6] Lightner DV, Hasson KW, White BL, Redman RM. Experimental infection of western hemisphere penaeid shrimp with Asian white spot syndrome virus and Asian yellow head virus. *J. Aquat Anim Health*. 1998; 10:271-281.
- [7] Lo CF, Leu JH, Chen CH, Peng SE, Chen YT, Chou CM et al. Detection of baculovirus associated with White Spot Syndrome (WSBV) in penaeid shrimps using polymerase chain reaction. *Dis Aquat Org*, 1996; 25:133-141.
- [8] Flegel TW. Major viral diseases of the black tiger prawn (*Penaeus monodon*) in Thailand. *World J Microbial Biotechnol*. 1997; 13:433-442.
- [9] Flegel TW, Alday-Sanz V. The crisis in Asian Shrimp Aquaculture: Current Status and future needs. *Journal of Applied Ichthyology*. 1998; 14:269-273.
- [10] Sriurairatana S, Boonyawiwat V, Gangnonngiw W, Laosutthipong C, Hiranchan J. White Feces Syndrome of Shrimp Arises from Transformation, Sloughing and Aggregation of Hepatopancreatic Microvilli into Vermiform Bodies Superficially Resembling Gregarines. *PLoS ONE*, 2014; 9(6):e99170. doi:10.1371/journal.pone.0099170
- [11] Limsuwan C. White Feces Disease in Thailand. *Boletines nicovita magazine*, 2010, 2-4.
- [12] Durai V, Gulan B, Michel Johnson, Maheswari ML, Pravin kumar M. Effect on white gut and white feces disease in semi intensive *Litopenaeus vannamei* shrimp culture system in south Indian state of Tamilnadu. *International Journal of Marine sci.* 2015; 5(14):1-5.
- [13] Burgents JE, Burnett KG, Burnett LE. Disease resistance of Pacific white shrimp, *Litopenaeus vannamei*, following the dietary administration of a yeast culture food supplement. *Aquaculture* 2004; 231(1-4):1-8.
- [14] Rhoobunjongde W, Hatai K, Wada S, Kubota SS. *Fusarium moniliforme* (Sheldon) isolated from gills of kuruma prawn *Penaeus japonicus* (Bate) with black gill disease. *Nippon Suisan Gakkaishi* 1991; 57(4):629-35.
- [15] Gopalakrishnan A, Parida A. Incidence of loose shell syndrome disease of the shrimp *Penaeus monodon* and its impact in the grow-out culture. *Curr. Sci*, 2005; 88:1148-1154.
- [16] Mastan SA, Incidence of loose shell syndrome (LSS) in cultured *Litopenaeus vannamei* in A.P., *Indo American journal of pharmaceutical research* 2015; 5(7):2600-2604.
- [17] Raja K, Gopalakrishnan A, Singh R, Vijayakumar R. Loose Shell Syndrome (LSS) in *Litopenaeus vannamei* grow-out Ponds and its Effect on Growth and Production. *Fish Aquac J*. 2015; 6:151. Doi: 10.4172/2150-3508.1000151.
- [18] Ha, N.T.H., Ha, D.T., Thuy, N.T. and Lien, V.T.K., 2010. Enterocytozoon hepatopenaei has been detected parasitizing tiger shrimp (*Penaeus monodon*) cultured in Vietnam and showing white feces syndrome (In Vietnamese with English abstract). *Agric. Rural Dev.: Sci. Technol.* 12: 45-50 (translation from Vietnamese).
- [19] Sritunyalucksana, K., Sanguanrut, P., Salachan P. V., Thitamadee, S. and Flegel, T.W. 2014. Urgent appeal to control spread of the shrimp microsporidian parasite *Enterocytozoon hepatopenaei* (EHP).



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)