



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 7 Issue: VIII Month of publication: August 2019 DOI: http://doi.org/10.22214/ijraset.2019.8047

www.ijraset.com

Call: 🕥 08813907089 🔰 E-mail ID: ijraset@gmail.com



MNDO Analysis of Reactivity Parameters of Pyran and its Amido Derivatives

M. B. Kalhans¹, A. K. Singh²

¹Department of Chemistry, Bareilly College, Bareilly, (U.P.) 243005, India ²Department of Physics, Bareilly College, Bareilly, (U.P.) 243005, India

Abstract: MNDO method is widely used for the study of molecular electronic parameters. Analytical calculations of the gradient are key to efficient geometry optimization. The IP, EA and q at O in Pyran and its derivatives have been calculated with the help of MNDO calculations and by substituting these values in Klopmann equation the softness indicators have been evaluated. The core-core repulsion and softness parameters of the compounds indicate the comparative donor capacity of Pyran and its Amido derivatives.

Keywords: MNDO, Gradient, Geometry optimization, softness indicators.

I. INTRODUCTION

The MNDO calculation have recently been used for study of molecular electronic structure.¹⁻³ The reactivity of compounds have been mainly studied with the help of the knowledge of frontier orbitals. Klopman gave a quantitative approach for explaining the reactivity of nucleophiles and electrophiles, and developed an equation for evaluating the quantitative values of reactivity.⁴⁻⁷ Klopman calculations were based upon the perturbation theory of charge and frontier controlled reactions. The charge controlled reactions were used to explain the ionic interactions and frontier controlled reactions to explain the covalent interactions.⁸ A chemical system may be an atom, molecule, ion or radical or several such units in a state of interaction. The chemical potential must be constant everywhere in such a system and is defined by

$$\mu = \left(\frac{\delta E}{\delta N}\right)_{v} = \left(\frac{\delta E}{\delta P}\right)_{v}$$
(1)

Analytical calculations of the gradient are key to efficient geometry optimization. The SCF energy expression is given by

$$E_{\rm HF} = \sum_{i=1}^{N12} E_i + \frac{1}{2} \sum_{v=1}^{b} \sum_{v=1}^{b} P_{vs} H_{rs}^{\rm core} + V_{\rm N,N}$$
(2)

The geometry optimization schemes are those of Mclver and Komornicki and force method due to Pulay.⁹⁻¹²

The optimization method developed by Pulay is based on Newton Raphson minimization algorithms. The method required some approximation to force constant matrix k, which is assumed to diagonal. Force constant calculation requires the determination of forces in the neighborhood of the equilibrium. Ki value is used to quantify ligand receptor interaction¹³⁻¹⁵. If the ith force F_i is changed by ΔF_i arising from internal coordinate change Δq_i then

$$k_{ij} = -\frac{\Delta F_i}{\Delta q_i} \tag{3}$$

II. METHODOLOGY

The LCAO approximation, from Hartee-Fock orbitals to Roothm equation served as theoretical basis for approximate molecular orbital theories.¹⁶⁻¹⁸ The molecular properties in semi empirical methods are not only derived from the principles of quantum mechanics but are also used to interpret and correlate the theoretical data with the experimental data. MNDO method involves the use of atomic parameters instead of pair parameters. The data and statistics obtained by using MNDO approximation for heat of formation, molecular geometries, ionization potential, dipole moment were quite close to experimental observation. The data obtained for MNDO method about bond length, bond angle and core-core repulsion are very accurate when compared to experimental values. The negative orbital energies are the calculated values of ionization potentials i.e. the validity of Koopman theorem which neglects the electronic reorganization and the charge in correlation of energies upon ionization. The mean absolute error for first ionization energies is reduced from 0.71 ev in MINDO/3 to 0.48 ev in MNDO. To identify difficult problems recent studies have introduced data modelability¹⁹⁻²¹. The limitation of MNDO method is an over estimation of the ionization energies corresponding to molecular orbitals with high coefficient.



International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.177 Volume 7 Issue VIII, Aug 2019- Available at www.ijraset.com

The ionization potential, election affinity and q at O in Pyran. Pyran 2-amide, Pyran 3-amide and Pyran 4-amide have been calculated with the help of MNDO calculations. Substituting these values in Klopman equation the softness parameters have been evaluated in Table-1.

The values of bond lengths between various atoms in Pyran and its derivatives are given in the Table 2. The sequence of bond lengths are given in Table 3. The bond orders of different bonds present in Pyran and the three isomeric amido Pyrans as calculated by MNDO method is given in Table 4.

The molecular valence electronic energy (Uval) including nuclear repulsion is taken as the sum of the purely electronic energy (Eel.val) of the valence electrons and the core-core repulsion energy Vcc^{22} .

$$U_{val} = E_{el.val} - V_{cc}$$
(4)

Although this form of Vcc is used in CNDO and INDO, it is more consistent with the approximation used to evaluate the electron core integrals in Dewar type theories to take –

$$V_{cc} = \sum_{B>A} \sum_{A} [C_A C_B (S_A S_A / S_B S_B) + f_{AB}]$$
(5)

In MNDO, the core – core repulsion is given by

$$f_{AB}^{MNDO} = C_A C_O \left(S_A S_A / S_B S_B \right) \left(e^{-\alpha_A R_{AB}} + e^{-\alpha_B R_{AB}} \right)$$
(6)

Where \propto_A and \propto_B are the parameters for atoms A and B. Core-core repulsion is calculated on the basis of above equation and given in Table 5.

III. RESULTS AND DISCUSSION

Although the stability or strength of individual bonds can be predicted in Pyran and the three isomers but these data do not furnish any comprehensive information which could be related to their comparative activity. However, it can be emphatically said that measurement of bond length has provided the information about the distance of amido group from ring oxygen in case of three isomers i.e. ortho, meta and para amido pyrans. Since the bond order and bond length of a particular bond is inversely proportional to each other the numerical values obtained for the two parameters by MNDO technique support the fact.

Core-Core repulsion in Pyran and three isomeric amido Pyrans have the sequence Pyran 2-admide (5375.48511) Pyran 3-admide (5341.42977) > Pyran 4-amide (5331.36629) > Pyran (2719.24801). In Pyran 2-amide the atomic cores of substituents – CONH₂ are nearest from ring oxygen this gives maximum core-core repulsion. In case of Pyran 3-amide the atomic cores of substituents – CONH₂ is at intermediate distance from O, this gives intermediate value of core-core repulsion and in Pyran 4 amide, the atomic cores of the substituent are at maximum distance from atom O so it has minimum core-core repulsion except Pyran which has not atomic cores of an additional substituent group thereby giving least value of core-core repulsion.

IV. CONCLUSION

Core-core repulsion gives a measure of the comfortability of the various atomic nuclei constituting the 3D molecular structure. The core-core repulsion order of the three isomeric amido pyrans with pyran itself is; Pyran 2 amide > Pyran 3-amide > Pyran 4 amide > Pyran. Analysis of table-1 reveals that $E_m^{\#}$ values of Pyran and three isomeric Amido Pyrans are in the sequence Pyran (1.793441) > 4-Amido Pyran (1.012191) > 3-Amido Pyran (0.9439262) > 2-Amido Pyran (0.599695). So the tendency to act as legand is maximum in Pyran and minimum in 2-Amido Pyran. So the study may be helpful in predicting the donor capacity of a particular site in larges molecules to make an easy route for synthesising useful derivatives.

REFERENCES

- [1] M.J.S. Dewar, W.J. Theil, J. Am. Chem. Soc., 99, 4899, 4707 (1977).
- [2] M.J.S. Dewar, M.L. Makee, J. Am. Chem. Soc., 99, 5231(1977).
- [3] M.J.S. Dewar, H.S. Rzepa, J. Am. Chem. Soc., 100, 58777, 784 (1978).
- [4] A.D. Daniels, G.E. Scuseria, J. Chem. Phys, 110, 1321 (1999).
- [5] H.H. Wadleigh, I.V. Ionova, E.A. Carter, J. Chem. Phys., 110, 4152 (1999).
- [6] J.F. Rico, r. Lopoz, G. Ramirez, J. Chem. Phys. 110, 4213 (1999).
- [7] G. Klopmann, J. Am. Chem. Soc., 90, 223 (1968).
- [8] G. Klopmann (Ed.) Chemical Reactivity and Reaction Paths, Wiley Inter science, New York (1974).
- [9] M.C. Flanigan, A.Konornicki, J.W. McIver, Modern Theoretical Chemistry Vol. 8, Plenum Press, New York (1977).
- [10] M.J.S. Dewar, H.S. Rzepa, J. Am. Chem. Soc. 100, 777 (1978).
- [11] P. Pulay, Mol. Phys., 17, 197 (1969).
- [12] P. Pulay, Modern Theoretical Chemistry, Vol. 3, Applications of electronic structure theory, Plenum Press, New York p. 153 (1977).
- [13] A. Cortes, M. Cascante, M.L. Cavdenas, A.B. Cornish, Biochem. J. 357; 263-268 (2001).





ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.177 Volume 7 Issue VIII, Aug 2019- Available at www.ijraset.com

- [14] R.R. Neubig, M. Spedding, T. Kenakin, A. Christopous Tos; Updates on terms and symbols in quantitative pharmacology. Pharm col. Rev. 554, 597-600 (2003).
- [15] C.L. Brace, J.L. Melville, D. Picketts, J.D. Hirst, J. Inf. model 47, 1,219-227 (2007).
- [16] D. R. Hartec, Proc. Cambridge Phil. Soc. 24, 89 (1978).
- [17] R.P. Iczkowski, J.L. Margrave, J. Am. Chem. Soc. 83, 3547 (1961).
- [18] M. Misono, E. Ochiavi Y. Saito and Y. Yaneda J. Inor. Nucl. Chem., 29, 2685 (1967).
- [19] D.P. Marks and R.S. Drago, J. Am. Chem. Soc., 97, 3324, (1975).
- [20] A Golbraikh, E. Muratov, D. Fourches, A. Tropsha, Chem. Inf. Model, 54, 1, 1-4 (2014).
- [21] A. Golbraikh, D. Fourches, A. Sedijkh, E. Muratov, I. Liepina, A. Tropsha; Practical Aspects of Computational Chem.-II, Springer, Boston, pp. 187-230 (2014).
- [22] Charlie Harper, 'Introductors to Mathematical Physics', Prentice Hall of India Pvt. Ltd. (1995).

Parameter	Pyran	Pyran 2-amdie	Pyran 3-amide	Pyran 4-amide
	12 02051		1 yran 5 annae	
Lonisation	13.92951	17.17413	17.88749	18.04708
potential				
Electron Affinity	-4.966315	-7.679936	-8.527449	-8.635807
Charge	-0.013952	-0.160161	-0.192525	-0.199776
Atom electron	1.82356	1.83843	1.83798	1.84038
density				
Electronegativity	4.481595	4.747095	4.680021	4.705635
Radius of atom	0.74	0.74	0.74	0.74
Fukui function	0.230035	0.2298038	0.2297475	0.2300475
Density	12.5536	12.65596	12.65287	12.63939
distribution fn				
Softness of atom	1.793441	0.599695	0.9439262	1.012191
$E_m^{\#}$				

Table.1. Values of different parameters of Pyran and three isomeric Amido pyrans at (O)

Table-2 Bond lengths between various atoms in Pyran and its amido derivatives

	6			
Type of Bond	Pyran	Pyran 2-amdie	Pyran 3-amide	Pyran 4-amide
$C_1 = C_2$	1.377347	1.339642	1.338281	1.345744
$C_2 = C_3$	1.475088	1.488549	1.487898	1.488597
C ₃ =O ₄	1.463994	1.425947	1.431119	1.427021
C ₄ =C ₅	1.256592	1.368518	1.350099	1.359912
$C_5 = C_6$	1.449241	1.352891	1.355030	1.345871
C ₆ =C ₇	-	1.501648	1.481309	1.491130 C ₁ =C ₇
C ₇ =O ₈	-	1.223713	1.227424	1.226641
C ₇ =N ₉	-	1.39777	1.3996	1.397141
N ₉ =H ₁₄	-	0.993452	0.991887	0.990522
N ₉ =H ₁₅	-	0.99041	0.990871	0.993145 (N ₉ =H ₁₃)
C1=H10	1.095756	1.093806	1.09625	1.092723 (C ₃ =H ₁₇)
C ₂ =H ₁₁	1.094006	1.094127	1.094173	1.098487
C ₃ =H ₁₂	1.105382	1.106241	1.105691	1.106223
C ₃ =H ₁₆	1.105412	1.106332	1.105752	1.106290
C ₆ =H ₁₃	1.092492	1.097218	1.096420	1.095101 (C ₆ =H ₁₈)



International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.177

Volume 7 Issue VIII, Aug 2019- Available at www.ijraset.com

Sequence of bon	1 bond lengths in Pyran, Pyran 2-amide, Pyran 5-amide and Pyran 4-amide			
Bonds	Sequence			
$C_1 = C_2$	Pyran > Pyran 4-amide > Pyran 2-amide > Pyran 3-amide			
	1.377347	1.345744	1.339642	1.338281
$C_2 = C_3$	Pyran 4-amid	e > Pyran 2-amide> Pyran	3-amide > Pyran	
	1.488597	1.488549	1.487898	1.475088
$C_3=O_4$	Pyran > Pyran	n 3-amide > Pyran 4-amid	e>Pyran 2-amide	
	1.463994	1.431119	1.427021	1.425947
$C_4=C_5$	Pyran 2-amid	e > Pyran 4-amide> Pyran	a 3-amide > Pyran	
	1.368518	1.359912	1.350099	1.256592
$C_5 = C_6$	Pyran > Pyran 3-amide > Pyran 2-amide > Pyran 2-amide			
	1.449241	1.355030	1.352891	1.345871
$C_2 = H_{11}$	Pyran 4-amide > Pyran > Pyran 3-amide > Pyran 4-amide			
	1.098407	1.094006	1.094173	1.094127
$C_3 = H_{12}$	Pyran 2-amide > Pyran 4-amide > Pyran 3-amide > Pyran			
	1.106241	1.106223	1.105691	1.105382
$C_7=O_8$	 Pyran 3-amide > Pyran 4-amide > Pyran 3-amide 			nide
		1.227424	1.226641	1.22373
$C_7=N_9$	 Pyran 3-amide > Pyran 2-amide > Pyran 4-amide 			nide
		1.399600	1.397770	1.397141
$N_9 = H_{14}$	 Pyran 2-amide > Pyran 3-amide > Pyran 3-amide 			nide
		0.993452	0.991887	0.990522
C ₃ =H ₁₆	Pyran 2-amid	e > Pyran 4-amide> Pyran	3-amide > Pyran	
	1.106332	1.106290	1.105752	1.105412

Table-3	
Sequence of bond lengths in Pyran, Pyran 2-amide, Pyran 3-amide and Pyran 4-amid	le

Table-4

Bond order between	various atom	s in Pyran ar	nd its amide	derivatives
Dona oraci between	various atom	is min yrun u	na no unnac	activatives

		5		
Type of Bond	Pyran	Pyran 2-amdie	Pyran 3-amide	Pyran 4-amide
$C_1 = C_2$	1.529575	1.870828	1.879819	1.824243
$C_2 = C_3$	1.048076	0.994502	0.995467	0.996301
C ₃ =O ₄	0.877255	0.963798	0.949721	0.962350
$C_4 = C_5$	1.56671	1.063188	1.098160	1.061889
$C_5 = C_6$	1.124637	1.755712	1.723492	1.797104
$C_6 = C_1$	1.418485	1.062583	1.044245	1.054638
$C_5 = C_7$	-	0.920321	0.955540	0.940691
C7=O8	-	1.762047	1.736440	1.742967
$C_1 = N_9$	-	1.150037	1.143943	1.154905
N9=H14	-	0.958922	0.961371	0.958451
N ₉ =H ₁₅	-	0.958247	0.958394	0.959930
C ₆ =H ₁₃	0.976995	0.955427	0.943138	0.955126
$C_1 = H_{10}$	0.979284	0.964261	0.956515	0.953398
C ₂ =H ₁₁	0.978237	0.962018	0.961099	0.948261
C3=H16	0.956437	0.963003	0.964602	0.962675
C ₃ =H ₁₂	0.956143	0.963984	0.964593	0.962662

Table-5 Core-core repulsion values

Molecule	Core-Core Repulsion in (Ev)
Pyran	2719.24801
Pyran 2-amide	5375.48511
Pyran 3-amide	5341.42977
Pyran 4-amide	5331.36629











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24*7 Support on Whatsapp)