iJRASET
International Journal For Research in Applied Science and Engineering Technology

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 7 **Issue:** VIII **Month of publication:** August 2019

DOI: http://doi.org/10.22214/ijraset.2019.8136

www.ijraset.com

Call: 08813907089 | E-mail ID: ijraset@gmail.com

International Journal for Research in Applied Science & Engineering Technology (IJRASET)
ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.177
Volume 7 Issue VIII, Aug 2019- Available at www.ijraset.com

An Advance Class of Analytic Functions with Fekete-Szegö Inequality using subordination Method

S. K. Gandhi[1], Gurmeet Singh[2], Preeti Kumawat[3], G. S. Rathore[4], Lokendra Kumawat[5]
[1, 3, 4, 5]*Department of Mathematics and Statistics, Mohanlal Sukhadia University, Udaipur (Raj) 313001*
[2]*Department of Mathematics, GSSDGS Khalsa College, Patiala, India*

Abstract: ***In this Paper we have introduced an advance class of analytic functions along with its subclasses by using principle of subordination and as so obtained sharp upper Bound of the function*** $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ ***belonging to these classes. Extremal functions are also investigated.***

Keywords: ***Bounded functions, Close to convex function, extremal function, Inverse Starlike functions, Starlike functions, Univalent functions.***

I. INTRODUCTION

Let $\mathcal{A}$ denote the class of analytic function

$f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ (1.1)

In the unit disc $= \{z: |z| < 1|\}$, $\mathcal{S}$ be the class of analytic univalent functions in $\mathbb{E}$.

Bieber Bach [7] proved that $|a_2| \leq 2$ for the functions $f(z) \in \mathcal{S}$. and Löwner [5] proved that $|a_3| \leq 3$ for the functions $f(z) \in \mathcal{S}$..

With the above known estimates this inequality plays an important role to determining estimates of higher coefficients for some sub classes $\mathcal{S}$ {Chhichra [11], Babalola [6]}

Using Löwner's method [5] In 1933, Fekete and szego investigated a well known relation between a_3 and a_2^2 for the class $\mathcal{S}$

$$|a_3 - \mu a_2^2| \leq \begin{cases} 3 - 4\mu & , if\ \mu \leq 0 \\ 1 + 2e^{\left(\frac{-2\mu}{1-\mu}\right)} & , if\ 0 \leq \mu \leq 1 \\ 4\mu - 3 & , if\ \mu \geq 1 \end{cases} \quad (1.2)$$

Let us define some subclasses of $\mathcal{S}$

Let $\mathcal{K}$ denotes subclasses of $\mathcal{S}$ of univalent convex functions $h(z) = z + \sum_{n=2}^{\infty} c_n z^n \in \mathcal{A}$ satisfying the condition

$Re\frac{((zh'(z))}{h'(z)} > 0, z \in \mathbb{E}.$ (1.3)

A function $f(z) \in \mathcal{A}$ is said to be close to convex if there exist $g(z) \in S^*$ such that

$Re\frac{((zf'(z))}{g(z)} > 0, z \in \mathbb{E}.$ (1.4)

The class of close to convex functions introduced by Kaplan [17],and he proved that close to convex functions are univalent.

$S^*(A,B) = \{ f(z) \in \mathcal{A} ; \frac{((zf'(z))}{g(z)} \prec \frac{1+Az}{1+Bz}, -1 \leq B \leq A \leq 1, z \in \mathbb{E} \}$ (1.5)

Where $S^*(A,B)$ is a subclass of S^*.

For strongly alpha quasi-convex functions Fekete-Szegö problem was studied by Abdel-Gawad [3]. The upper bound of $|a_3 - \mu a_2^2|$ for different functions in the class S has been investigated by many authors including Goel and Mehrok [13] and recently by Al-Shaqsi and Darus [4] Hayami and Owa [16], Al-Abbadi and Darus [10].

Gurmeet singh et al. [3] introduced the class of inverse Starlike functions

$g(z) = z + \sum_{n=2}^{\infty} b_n z^n \in A$ satisfying the condition

$\text{Re}\left(\frac{zf(z)}{2\int_0^z f(z)dz}\right) > 0, z \in E$ i.e. $\frac{zf(z)}{2\int_0^z f(z)dz} < \frac{1+z}{1-z}$ (1.6)

Gandhi et al. [14] established a new class of analytic functions with Fekete-szego inequality using subordination method.

Here introduce the class $\mathcal{A}$, of Univalent starlike functions $g(z) = z + \sum_{n=2}^{\infty} b_n z^n \in \mathcal{A}$ satisfying the condition

©IJRASET: All Rights are Reserved

$\left[\frac{z\{zf(z)\}'}{2f(z)}\right] \prec \left(\frac{1+z}{1-z}\right)^{\alpha}; \alpha > 0$ (1.7)

And subclass consisting of the functions $g(z) = z + \sum_{n=2}^{\infty} b_n z^n \in \mathcal{A}$

satisfying the condition

$\left[\frac{z\{zf(z)\}'}{2f(z)}\right] \prec \left(\frac{1+Az}{1+Bz}\right)^{\alpha}; -1 \le B \le A \le 1; \alpha > 0$ (1.8)

Here, Symbol $\prec$ stands for subordination, defined as follows:

A. Principle of Subordination

If $f(z)$ and $F(z)$ are two functions which are analytic in $\mathbb{E}$, then $f(z)$ is called a subordinate to $F(z)$ in $\mathbb{E}$, if there exists a function $w(z)$ which is analytic in $\mathbb{E}$ satisfying the conditions

(i) $w(0) = 0$ and (ii) $|w(z)| < 1$

such that $f(z) = F(w(z))$, where $z \in \mathbb{E}$ and we denote it as $f(z) \prec F(z)$.

Let $\mathcal{U}$ denote the class of analytic bounded functions of the form

$w(z) = \sum_{n=1}^{\infty} d_n z^n, w(0) = 0, |w(z)| < 1$ (1.9)

With $|d_1| \le 1, |d_2| \le 1 - |d_1|^2$.

II. RESULTS AND DISCUSSION

1) Theorem 1: If $f(z) \in \mathcal{A}$, then the result

$$|a_3 - \mu a_2^2| \le \begin{cases} 10\alpha^2 - 16\mu\alpha^2 & , if\ \mu \le \frac{5\alpha-1}{8\alpha} \quad (1.10) \\ 2\alpha & , if\ \frac{5\alpha-1}{8\alpha} \le \mu \le \frac{5\alpha+1}{8\alpha} \quad (1.11) \\ 16\mu\alpha^2 - 10\alpha^2 & , if\ \mu \ge \frac{5\alpha+1}{8\alpha} \quad (1.12) \end{cases}$$

is sharp.

Proof: By using expantion method (1.7) leads to

$1 + \frac{1}{2}a_2 z + (a_3 - \frac{1}{2}a_2^2) z^2 + \text{------} = 1 + 2\alpha c_1 z + 2\alpha(c_2 + \alpha c_1^2) z^2 + \text{-----}$ (1.13)

After Identifying the terms we have

$|a_3 - \mu a_2^2| \le |2\alpha c_2 + 10\alpha^2 c_1^2 - 16\mu\alpha^2 c_1^2|$

This leads to

$|a_3 - \mu a_2^2| \le 2\alpha + [|10\alpha^2 - 16\mu\alpha^2| - 2\alpha]|c_1|^2$ (1.14)

Case I: If $\mu \le \frac{5}{8}$, then (1.14) leads to

$|a_3 - \mu a_2^2| \le 2\alpha + [(10\alpha^2 - 2\alpha) - 16\mu\alpha^2|c_1|^2$ (1.15)

Subcase I(a): If $\mu \le \frac{5\alpha-1}{8\alpha}$, then (1.15) leads to

$|a_3 - \mu a_2^2| \le 10\alpha^2 - 16\mu\alpha^2$ (1.16)

Subcase I(b) : If $\mu \ge \frac{5\alpha-1}{8\alpha}$, then (1.15) leads to

$|a_3 - \mu a_2^2| \le 2\alpha$ (1.17)

Case II : If $\mu \ge \frac{5}{8}$, then (1.14) leads to

$|a_3 - \mu a_2^2| \le 2\alpha + [16\mu\alpha^2 - (10\alpha^2 + 2\alpha)]|c_1|^2$ (1.18)

Subcase II(a): If $\mu \le \frac{5\alpha+1}{8\alpha}$, then (1.18) leads to

$|a_3 - \mu a_2^2| \le 2\alpha$ (1.19)

Subcase II(b): If $\mu \ge \frac{5\alpha+1}{8\alpha}$, then (1.18) leads to

$|a_3 - \mu a_2^2| \le 16\mu\alpha^2 - 10\alpha^2$ (1.20)

Combining subcase II(a) and subcase I(b), we get

$|a_3 - \mu a_2^2| \le 2\alpha$, $if\ \frac{5\alpha-1}{8\alpha} \le \mu \le \frac{5\alpha+1}{8\alpha}$ (1.21)

This completes the theorem, therefore the result is sharp.

Extremal function for the first and third inequality is given by

©IJRASET: All Rights are Reserved

$$f_1(z) = \frac{z}{(1-\alpha z)^4} \quad (1.22a)$$

And Extremal function for the second inequality is given by

$$f_2(z) = \frac{z}{(1-\alpha z^2)^2} \quad (1.22b)$$

2) Theorem 2 : If $\boldsymbol{f(z) \in \mathcal{A}}$, then the result

$|\boldsymbol{a_3} - \mu \boldsymbol{a_2^2}| \leq$

$$\begin{cases} (A-B)(2A-3B)\alpha^2 - 4\mu\alpha^2(A-B)^2 & ,if\ \mu \leq \dfrac{(2A-3B)\alpha - 1}{4(A-B)\alpha} \quad (1.23a) \\ (A-B)\alpha & ,if\ \dfrac{(2A-3B)\alpha - 1}{4(A-B)\alpha} \leq \mu \leq \dfrac{(2A-3B)\alpha + 1}{4(A-B)\alpha} \quad (1.23b) \\ 4\mu\alpha^2(A-B)^2 - (A-B)(2A-3B)\alpha^2 & ,if\ \mu \leq \dfrac{(2A-3B)\alpha + 1}{4(A-B)\alpha} \quad (1.23c) \end{cases}$$

is sharp.

Proof: By using expantion method (1.8) leads to

$$1 + \frac{1}{2}a_2 z + (a_3 - \frac{1}{2}a_2^2) z^2 + \text{------} = 1 + (A-B)\alpha c_1 z + (A-B)\alpha(c_2 - B\alpha c_1^2) z^2 + \text{----} \quad (1.24)$$

After Identifying the terms in (1.24) we have

$$|a_3 - \mu a_2^2| \leq |(A-B)\,\alpha(c_2 - B\alpha c_1^2) + 2(A-B)^2\alpha^2 c_1^2 - 4\mu(A-B)^2\alpha^2 c_1^2|$$

This leads to

$$|a_3 - \mu a_2^2| \leq (A-B)\alpha + \{|2(A-B)^2\alpha^2 - B(A-B)\alpha^2 - 4\mu(A-B)^2\alpha^2| - (A-B)\alpha\}|c_1|^2 \quad (1.25)$$

here two cases arise:

Case I: If $\mu \leq \frac{2A-3B}{4(A-B)}$, then (1.25) leads to

$$|a_3 - \mu a_2^2| \leq (A-B)\alpha + [(A-B)\{(2A-3B)\alpha - 1\}\alpha - 4\mu(A-B)^2\alpha^2]\,|c_1|^2 \quad (1.26)$$

Under this case (1.26) two subcases arise:

Subcase I(a): If $\mu \leq \frac{(2A-3B)\alpha-1}{4(A-B)\alpha}$, then (1.26) leads to

$$|a_3 - \mu a_2^2| \leq \{(A-B)(2A-3B)\alpha - 4\mu(A-B)^2\,\alpha^2 \quad (1.27)$$

Subcase I(b): If $\mu \geq \frac{(2A-3B)\alpha-1}{4(A-B)\alpha}$, then (1.26) leads to

$$|a_3 - \mu a_2^2| \leq (A-B)\alpha \quad (1.28)$$

Case II: If $\mu \geq \frac{2A-3B}{4(A-B)}$, then (1.25) leads to

$$|a_3 - \mu a_2^2| \leq (A-B) + [4\mu(A-B)^2\alpha^2 - (A-B)\{(2A-3B)\alpha + 1\}\alpha]\,|c_1|^2 \quad (1.29)$$

Under this case again two subcases arise:

Subcase II(a): If $\mu \leq \frac{(2A-3B)\alpha+1}{4(A-B)\alpha}$, then (1.29) leads to

$$|a_3 - \mu a_2^2| \leq (A-B)\alpha \quad (1.30)$$

Subcase II(B): $\mu \geq \frac{(2A-3B)\alpha+1}{4(A-B)\alpha}$, then (1.29) leads to

$$|a_3 - \mu a_2^2| \leq \{4\mu(A-B)^2\alpha^2 - (A-B)(2A-3B)\alpha^2 \quad (1.31)$$

Combining subcase II(a) and subcase I(b), we get

$$|a_3 - \mu a_2^2| \leq (A-B)\alpha \quad ,\ if\ \frac{(2A-3B)\alpha - 1}{4(A-B)\alpha} \leq \mu \leq \frac{(2A-3B)\alpha + 1}{4(A-B)\alpha} \quad (1.32)$$

This completes the theorem therefore, the result is sharp.

Extremal function for the first and third inequality is given by

$$f_1(z) = z\left\{1 + \left(\frac{2A-2B}{2A-3B}\right)z\alpha\right\}^{2A-3B} \quad (1.33a)$$

And for the for second inequality is given by

$$f_2(z) = \frac{z}{(1-\alpha z^2)^{A-B}} \quad (1.33b)$$

©IJRASET: All Rights are Reserved

III. CONCLUDING REMARKS AND COROLLARIES

1) Corollary 1.1: Taking $\alpha = 1$ in the theorem 1, we get

$$|a_3 - \mu a_2^2| \leq \begin{cases} 10 - 16\mu & , if\ \mu \leq \frac{1}{2} \quad (1.34) \\ 2 & , if\ \frac{1}{2} \leq \mu \leq \frac{3}{4} \quad (1.35) \\ 16\mu - 10 & , if\ \mu \geq \frac{3}{4} \quad (1.36) \end{cases}$$

These estimates were derived by Keogh and Merkes [1] and the results are for the class of univalent convex functions.
Further if we take A = 1 and B = -1 $(-1 \leq B \leq A \leq 1)$ in the result of theorem 2 , we get the result of theorem 1, therefore our result for the theorem 2 reduces to the result of the theorem1. Hence theorem 2 is the generalization of theorem 1. And the results are sharp and also if we put A = 1 and B = -1 in extremal function of theorem 2, we get the extremal function of theorem 1. The extremal function given by [(1.22a),(1.22b)] increases as α increases and decreases as α decreases and the extremal function given by [(1.33a),(1.33b)] also increases and decreases as α increases and decreases respectively. Hence extremal function is an increasing function.

REFERENCES

[1] F. R. Keogh, E. R.Merkes, "A coefficient inequality for certain classes of analytic functions", Procedure of American Mathematical Society, 20, 8 (1989)
[2] Gurmeet Singh, M. S. Saroa and B. S.Mehrok, "Fekete-szegö inequality for a new class of analytic functions", Elsevier, Proc. of International conference on Information and Mathematical Sciences, 90 (2013)
[3] H. R.Abdel-Gawad, D.K. Thomas ,"The Fekete-Szegö problem for strongly close-to-convex functions" Proceedings of the Amercian Mathematical Society, vol. 144, no.2 345(1992)
[4] K. Al-Shaqsi, M.Darus , "On Fekete-Szegö problems for certain subclass of analyticFunctions", Applied Mathematical Sciences, vol. 2 no. 9-12,pp. 431(2008)
[5] K. Löwner, "Uber monotone Matrixfunktionen", Math. Z 38, 177 (1934).
[6] Kunle Oladeji Babalola, "The fifth and Sixth coefficient of α – close- to –convex function", Kragujevac J. Math. 32, 5 (2009)
[7] L. Bieberbach, "Uber einige Extremal Probleme im Gebiete der Konformen Abbildung" Math. 77, 153 (1916)
[8] L. Bieberbach, "Uberdie Koeffizientem derjenigem Potenzreihen, welche eine Schlithe Abbildung des Einheitskrises Vermittelen", Preuss. AKad. Wiss Sitzungsb. 940 (1916)
[9] M.Fekete, G.Szego, "Eine Bemerkung Uber ungerade Schlichte Funktionen" , J. London Math. Soc. 8 , 85 (1933)
[10] M. H. Al-Abbadi, M.Darus , "Fekete-Sezegö theorem for a certain class of analytic functions", Sanis Malaysiana,vol. 40, no. 4, 385(2011)
[11] P. N. Chhichra, "New Subclasses of the class of close to convex functions", Procedure of American Mathematical Society , 62 , 37 (1977)
[12] R.M. Goel , B. S. Mehrok, "A Subclass of Univalent functions", Houston Journal of Mathematics 8, 343 (1982)
[13] R.M. Goel , B. S. Mehrok, "A coefficient inequality for certain classes of analytic function", Tamkang Journal of Mathematics 22, 153 (1990)
[14] S. K. Gandhi, Gurmeet Singh, Preeti Kumawat, G. S. Rathore and Lokendra Kumawat, "A new class of analytic functions with Fekete-Szego Inequality using subordination Method",International journal of research in advent technology, vol. 6, no. 9,(2018)
[15] S. R. Keogh, E. R. Merkes, "A coefficient inequality for certain classes of analytic functions", Proceedings of the American Mathematical Society. Vol. 20, 8 (1969)
[16] T. Hayami, S. Owa and H. M. Srivastava, "Coefficient inequalities for certain classes of analytic and univalent functions" , J. Ineq. Pure and Appl. Math. 8(4), 1 (2007)
[17] W. Kaplan, "Close –to-convex schlicht functions", Michigan Mathematical Journal 1, 169 (1952)

©IJRASET: All Rights are Reserved



10.22214/IJRASET

45.98

IMPACT FACTOR: 7.129

IMPACT FACTOR: 7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)