iJRASET
International Journal For Research in Applied Science and Engineering Technology

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 7 **Issue:** IX **Month of publication:** September 2019

DOI: http://doi.org/10.22214/ijraset.2019.9129

www.ijraset.com

Call: 08813907089 | E-mail ID: ijraset@gmail.com

International Journal for Research in Applied Science & Engineering Technology (IJRASET)
ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.177
Volume 7 Issue IX, Sep 2019- Available at www.ijraset.com

Integrals of Four Variables with Statistical Distribution associated with hyper geometric Function of Matrix Argument

Dr. Pooja Singh[1], Ms Nidhi [2]
[1, 2]*Maharaja Surajmal Institute of Technology, Affliated to GGSIPU, Delhi*

Abstract: ***In this chapter, we evaluated ten integrals associated with hypergeometric function of four variables of matrix argument with their statistical distribution. All the matrix involved were real positive definite symmetric of order m x m.***
Keywords: Hypergeometric function, Matrix argument

I. INTRODUCTION

In this paper, we have evaluated forty integrals associated with hypergeometric function of four variables of matrix argument with their statistical distribution. All the matrices involved are real positive definite symmetric of order m x m. We will start with introducing matrix sequences, matrix series, and concepts analogous to convergence of series in scales variable. A matrix series is obtained by adding up the matrices in a matrix sequence. For example if A_0, A_1, A_2 …… is matrix series given by

$$F(A) = \sum_{k=0}^{\infty} A_k \qquad (1.1)$$

If the matrix series is a power series than we will be considering powers of matrices and hence in this case the series will be define only for n x n matrices,for an n x n matrix A. consider the series.

$$g(A) = \sum_{k=0}^{\infty} a_k A_k \text{ ……..} \qquad (1.2)$$

where $a_0, a_1, \ldots\ldots a_k$ are scalars.
As in the case of scalars series, convergence of a matrix will be defined in terms of the sums.
A general hypergeometric series ${}_pF_q(.)$ in a real scalar variable X is defined as following :

$${}_pF_q(a_1, \ldots\ldots a_p ; b_1, \ldots\ldots b_q; X) = \sum_{r=0}^{\infty} \frac{(a_1)_r \ldots\ldots (a_p)_r \, X^r}{(b_1)_r \ldots\ldots (b_q)_r \, r!} \qquad (1.3)$$

For $(a)_m = a(a+1) \ldots\ldots (a+m-1)$
$(a)_0 = 1 \qquad a \neq 0$

For example: ${}_0F_0(\,;\,;X) = e^x$
${}_1F_0(\alpha;\,;X) = (I-x)^{-\alpha}$ for $|X| < 1$

In (1.3) there are p upper parameters $a_1, \ldots\ldots, a_p$ and q lower parameters $b_1, \ldots\ldots, b_q$. The series in (1.3) is convergent for all $|X| < \infty$ ifq $\geq$p, convergent for $|X|<I$ if p = q+1, divergent for all X, $X \neq 0$,if p> q+1 and the convergence. Condition for X = 1 and X = -1 can also be worked out. A matrix series is in n x n matrix. A corresponding to the right side in (1.3) is obtained by replacing X by A, thus we may define a hypergeometric series in an n x n matrix. A as follows.

$${}_pF_q(a_1, \ldots\ldots a_p ; b_1, \ldots\ldots b_q; A) = q \geq p, \sum_{r=0}^{\infty} \frac{(a_1)_r \ldots\ldots (a_p)_r \, X^r}{(b_1)_r \ldots\ldots (b_q)_r \, r!} \qquad (1.4)$$

where $a_1, \ldots\ldots a_p, b_1 \ldots\ldots b_q$ are scalars.
The series in (1.4) is convergent for all A is q $\geq$ p convergent for p = q + 1 when the Eigen values of A are all less than 1 in absolute value and divergent when p > q + 1.
Similarly, it may be defined for two, three and four variables. Other definition involving in this chapter for four variables of m x m matrix. Exactly similar analogous in scalar variable due to Exton (1985). In what follows we shall take p, q, r, and s to be positive

©IJRASET: All Rights are Reserved

integers of the symbols X and Δ (n, a) stand the sequence of parameters of square positive definite matrices $X_1, X_2, \ldots\ldots X_r$ of order n x n and $\frac{\alpha}{n}, \frac{\alpha+1}{n}, \ldots\ldots \frac{\alpha+n-1}{n}$ respectively.

Also, in all be established here after, proper conditions of convergence of the involved are assumed. In which follows X, Y, Z, T, U etc. matrices are positive definite symmetric of same order m x m.

II. INTEGRALS OF FOUR VARIABLES WITH STATISTICAL DISTRIBUTION ASSOCIATED WITH HYPERGEOMETRIC FUNCTION OF MATRIX ARGUMENT

The formulae to be established are as following :

1) Result 1: $\int_0^I U^{a-\frac{m+1}{2}} (I-U)^{b-\frac{m+1}{2}} {}_1F_1(a, 1-a; c; \frac{U}{2})$.

$$F_1^4\left[a_1, a_1, a_1, a_2, b_1, b_1, b_1, b_1, c_1, c_2, c_3, c_4; X(I-U)^n, Y(I-U)^n, Z(I-U)^n, T(I-U)^n\right] dU$$

$$= LF_1^4\left[\begin{matrix} a_1, a_1, a_1, a_2, b_1, b_1, b_1, b_1, c_1, c_2, c_3, c_4; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}), \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{matrix}\right] \quad (2.1)$$

where

$$F_1^4\left[a_1, a_1, a_1, a_2, b_1, b_1, b_1, b_1, c_1, c_2, c_3, c_4; (I-U)^n X, (I-U)^n Y, (I-U)^n Z, (I-U)^n T\right]$$

$$= \sum_{p,q,r,s=0}^{\infty} \frac{(a_1)_{p+q+r} (a_2)_s (b_1)_{p+q+r+s}}{p!q!r!s!(c_1)_p (c_2)_q (c_3)_r (c_4)_s} (I-U)^{np} X^p (I-U)^{nq} Y^q (I-U)^{nr} Z^r (I-U)^{ns} T^s$$

so that

$$F_1^4\left[a_1, a_1, a_1, a_2, b_1, b_1, b_1, b_1, c_1, c_2, c_3, c_4; (I-U)^n X, (I-U)^n Y, (I-U)^n Z, (I-U)^n T\right]$$

$$= F_1^4\left[(I-U)^n X, (I-U)^n Y, (I-U)^n Z, (I-U)^n T\right]$$

Then a probability density function (p.d.f.) of (3.2.1.) is given by: $F(U) = \frac{U^{a-\frac{m+1}{2}} (I-U)^{b-\frac{m+1}{2}} F_1(\chi_1) F_1^4(\chi_2)}{LF_1^4(\chi_3)}$

= 0 else where

where $\chi_1 = (a, a-1; c; \frac{U}{2})$

$$\chi_2 = \left[(I-U)^n X, (I-U)^n Y, (I-U)^n Z, (I-U)^n T\right]$$

$$\chi_1 = \left[\begin{matrix} a_1, a_1, a_2, a_1, b_1, b_1, b_1, b_1, c_1, c_2, c_3, c_4; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}), \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{matrix}\right]$$

2) Result 2: $\int_0^I U^{a-\frac{m+1}{2}} (I-U)^{b-\frac{m+1}{2}} {}_1F_1(a, 1-a; c; \frac{U}{2}) . F_2^4\left[(I-U)^n X, (I-U)^n Y, (I-U)^n Z, (I-U)^n T\right] dU$

$$= LF_2^4\left[\begin{matrix} a_1, a_1, a_2, a_2, b_1, b_1, b_1, b_1, c_1, c_2, c_3, c_4; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}), \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{matrix}\right] \quad (2.2)$$

©IJRASET: All Rights are Reserved

where

$$F_2^4\left(a_1, a_1, a_2, a_2, b_1, b_1, b_1, b_1, c_1, c_2, c_3, c_4; (I-U)^n X, (I-U)^n Y, (I-U)^n Z, (I-U)^n T\right)$$

$$= \sum_{p,q,r,s=0}^{\infty} \frac{(a_1)_{p+q}(a_2)_{r+s}(b_1)_{p+q+r+s}}{p!q!r!s!(c_1)_p(c_2)_q(c_3)_r(c_4)_s}(I-U)^{np}X^p(I-U)^{nq}Y^q(I-U)^{nr}Z^r(I-U)^{ns}T^s$$

Then a probability density function (p.d.f.) of (3.2.2) is given by: $F(U) = \dfrac{U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}}F_1(\chi_1)F_2^4(\chi_2)}{LF_2^4(\chi_4)}$

= 0 else where

where $\chi_4 = \begin{bmatrix} a_1, a_1, a_2, a_2, b_1, b_1, b_1, b_1, c_1, c_2, c_3, c_4; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}) \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{bmatrix}$

3) Result 3: $\int_0^I U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}} F_1(a, 1-a; c; \frac{U}{2}).F_3^4\left[(I-U)^n X, (I-U)^n Y, (I-U)^n Z, (I-U)^n T\right]dU$

$$= LF_3^4\begin{bmatrix} a_1, a_1, a_2, a_3, b_1, b_1, b_1, b_1, c_1, c_2, c_3, c_4; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}), \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{bmatrix} \quad (2.3)$$

where

$$F_3^4\left(a_1, a_1, a_2, a_3, b_1, b_1, b_1, b_1, c_1, c_2, c_3, c_4; (I-U)^n X, (I-U)^n Y, (I-U)^n Z, (I-U)^n T\right)$$

$$= \sum_{p,q,r,s=0}^{\infty} \frac{(a_1)_{p+q}(a_2)_{r+s}(b_1)_{p+q+r+s}}{p!q!r!s!(c_1)_p(c_2)_q(c_3)_r(c_4)_s}(I-U)^{np}X^p(I-U)^{nq}Y^q(I-U)^{nr}Z^r(I-U)^{ns}T^s$$

Then a (p.d.f.) of (2.3) is given by : $F(U) = \dfrac{U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}}F_1(\chi_1)F_3^4(\chi_2)}{LF_3^4(\chi_5)}$

= 0 else where

where $\chi_5 = \begin{bmatrix} a_1, a_1, a_2, a_3, b_1, b_1, b_1, b_1, c_1, c_2, c_3, c_4; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}) \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{bmatrix}$

4) Result 4: $\int_0^I U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}} F_1(a, 1-a; c; \frac{U}{2}).F_4^4\left[(I-U)^n X, (I-U)^n Y, (I-U)^n Z, (I-U)^n T\right]dU$

$$= LF_4^4\begin{bmatrix} a_1, a_1, a_1, a_1, b_1, b_1, b_1, b_2, c_1, c_2, c_3, c_4; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}), \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{bmatrix} \quad (2.4)$$

©IJRASET: All Rights are Reserved

where $F_4^4\left(a_1, a_1, a_1, a_1, b_1, b_1, b_2, b_1, c_1, c_2, c_3, c_4; X, Y, Z, T\right)$

$$= \sum_{p, q, r, s = 0}^{\infty} \frac{(a_1)_{p+q+r+s}(b_1)_{p+q+s}(b_2)_r}{p!q!r!s!(c_1)_p(c_2)_q(c_3)_r(c_4)_s} X^p Y^q Z^r T^s$$

Then a (p.d.f.) of (2.4) is given by : $F(U) = \dfrac{U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}} F_1(\chi_1) F_4^4(\chi_2)}{LF_4^4(\chi_6)}$

= 0 else where

where $\chi_6 = \left[\begin{matrix} a_1, a_1, a_1, a_1, b_1, b_1, b_2, b_1, c_1, c_2, c_3, c_4; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}) \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{matrix}\right]$

5) Result5: $\int_0^I U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}} F_1(a, 1-a; c; \frac{U}{2}).F_5^4\left[(I-U)^n X, (I-U)^n Y, (I-U)^n Z, (I-U)^n T\right] dU$ =

$$LF_5^4\left[\begin{matrix} a_1, a_1, a_1, a_2, b_1, b_1, b_2, b_2, c_1, c_2, c_3, c_4; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}), \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{matrix}\right] \quad (2.5)$$

where $F_5^4\left(a_1, a_1, a_1, a_2, b_1, b_1, b_2, b_2, c_1, c_2, c_3, c_4; X, Y, Z, T\right)$

$$= \sum_{p, q, r, s = 0}^{\infty} \frac{(a_1)_{p+q+r}(a_2)_s(b_1)_{p+q}(b_2)_{r+s}}{p!q!r!s!(c_1)_p(c_2)_q(c_3)_r(c_4)_s} X^p Y^q Z^r T^s$$

Then a (p.d.f.) of (2.5) is given by : $F(U) = \dfrac{U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}} F_1(\chi_1) F_5^4(\chi_2)}{LF_5^4(\chi_7)}$

= 0 else where

where $\chi_7 = \left[\begin{matrix} a_1, a_1, a_1, a_2, b_1, b_1, b_2, b_2, c_1, c_2, c_3, c_4; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}) \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{matrix}\right]$

6) Result 6: $\int_0^I U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}} F_1(a, 1-a; c; \frac{U}{2}).F_6^4\left[(I-U)^n X, (I-U)^n Y, (I-U)^n Z, (I-U)^n T\right] dU$

$$= LF_6^4\left[\begin{matrix} a_1, a_1, a_1, a_2, b_1, b_2, b_3, b_1, c_1, c_2, c_3, c_4; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}), \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{matrix}\right] \quad (2.6)$$

where $F_6^4\left(a_1, a_1, a_1, a_2, b_1, b_2, b_3, b_1, c_1, c_2, c_3, c_4; X, Y, Z, T\right)$

©IJRASET: All Rights are Reserved

$$= \sum_{p,q,r,s=0}^{\infty} \frac{(a_1)_{p+q+r}(a_2)_s(b_1)_{p+s}(b_2)_q(b_3)_s}{p!q!r!s!(c_1)_p(c_2)_q(c_3)_r(c_4)_s} X^p Y^q Z^r T^s$$

Then a (p.d.f.) of (3.2.6) is given by : $F(U) = \dfrac{U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}} F_1(\chi_1) F_6^4(\chi_2)}{LF_6^4(\chi_8)}$

$= 0$ else where

where $\chi_8 = \left[\begin{matrix} a_1, a_1, a_1, a_2, b_1, b_2, b_3, b_1, c_1, c_2, c_3, c_4; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}) \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{matrix} \right]$

7) Result 7: $\int_0^I U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}} F_1(a, 1-a; c; \frac{U}{2}).F_7^4\left[(I-U)^n X, (I-U)^n Y, (I-U)^n Z, (I-U)^n T\right] dU$

$$= LF_7^4 \left[\begin{matrix} a_1, a_1, a_2, a_2, b_1, b_2, b_1, b_2, c_1, c_2, c_3, c_4; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}), \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{matrix} \right] \quad (2.7)$$

where $F_7^4(a_1, a_1, a_2, a_2, b_1, b_2, b_1, b_2, c_1, c_2, c_3, c_4; X, Y, Z, T)$

$$= \sum_{p,q,r,s=0}^{\infty} \frac{(a_1)_{p+q}(a_2)_{r+s}(b_1)_{p+r}(b_2)_q(b_3)_s}{p!q!r!s!(c_1)_p(c_2)_q(c_3)_{r+s}(c_4)_s} X^p Y^q Z^r T^s$$

Then a (p.d.f.) of (3.2.7) is given by : $F(U) = \dfrac{U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}} F_1(\chi_1) F_7^4(\chi_2)}{LF_7^4(\chi_9)}$

$= 0$ else where

where $\chi_9 = \left[\begin{matrix} a_1, a_1, a_2, a_2, b_1, b_2, b_1, b_2, c_1, c_2, c_3, c_4; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}) \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{matrix} \right]$

8) Result 8: $\int_0^I U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}} F_1(a, 1-a; c; \frac{U}{2}).F_8^4\left[(I-U)^n X, (I-U)^n Y, (I-U)^n Z, (I-U)^n T\right] dU$

$$= LF_8^4 \left[\begin{matrix} a_1, a_1, a_2, a_2, b_1, b_2, b_1, b_3, c_1, c_2, c_3, c_4; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}), \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{matrix} \right] \quad (2.8)$$

where $F_8^4(a_1, a_1, a_2, a_2, b_1, b_2, b_1, b_3, c_1, c_2, c_3, c_4; X, Y, Z, T)$

$$= \sum_{p,q,r,s=0}^{\infty} \frac{(a_1)_{p+q}(a_2)_{r+s}(b_1)_{p+r}(b_2)_q(b_3)_s}{p!q!r!s!(c_1)_p(c_2)_q(c_3)_r(c_4)_s} X^p Y^q Z^r T^s$$

©IJRASET: All Rights are Reserved

Then a (p.d.f.) of (2.8) is given by : $F(U) = \dfrac{U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}} F_1(\chi_1) F_8^4(\chi_2)}{L F_8^4(\chi_{10})}$

= 0 else where

where $\chi_{10} = \begin{bmatrix} a_1, a_1, a_2, a_2, b_1, b_2, b_3, b_4, c_1, c_2, c_3, c_4; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}), \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{bmatrix}$

9) Result 9: $\int_0^I U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}} F_1(a, 1-a; c; \frac{U}{2}). F_9^4\left[(I-U)^n X, (I-U)^n Y, (I-U)^n Z, (I-U)^n T\right] dU$

$$= L F_9^4 \begin{bmatrix} a_1, a_1, a_1, a_2, b_1, b_1, b_1, b_1, c_1, c_2, c_3, c_1; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}), \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{bmatrix} \quad (2.9)$$

where $F_9^4(a_1, a_1, a_1, a_2, b_1, b_1, b_1, b_1, c_1, c_2, c_3, c_1; X, Y, Z, T)$

$$= \sum_{p,q,r,s=0}^{\infty} \frac{(a_1)_{p+q+r}(a_2)_s (b_1)_{p+q+r+s}}{p!q!r!s!(c_1)_{p+s}(c_2)_q(c_3)_r} X^p Y^q Z^r T^s$$

Then a (p.d.f.) of (2.9) is given by : $F(U) = \dfrac{U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}} F_1(\chi_1) F_9^4(\chi_2)}{L F_9^4(\chi_{11})}$

= 0 else where

where $\chi_{11} = \begin{bmatrix} a_1, a_1, a_1, a_2, b_1, b_1, b_1, b_1, c_1, c_2, c_3, c_1; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}), \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{bmatrix}$

10) Result 10: $\int_0^I U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}} F_1(a, 1-a; c; \frac{U}{2}). F_{10}^4\left[(I-U)^n X, (I-U)^n Y, (I-U)^n Z, (I-U)^n T\right] dU$

$$= L F_{10}^4 \begin{bmatrix} a_1, a_1, a_2, a_2, b_1, b_1, b_1, b_1, c_1, c_2, c_1, c_3; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}), \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{bmatrix} \quad (2.10)$$

where $F_{10}^4(a_1, a_1, a_2, a_2, b_1, b_1, b_1, b_1, c_1, c_2, c_1, c_3; X, Y, Z, T)$

$$= \sum_{p,q,r,s=0}^{\infty} \frac{(a_1)_{p+q}(a_2)_{r+s}(b_1)_{p+q+r+s}}{p!q!r!s!(c_1)_{p+r}(c_2)_q(c_3)_s} X^p Y^q Z^r T^s$$

Then a p.d.f. $F(U) = \dfrac{U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}} F_1[\chi_1] F_{10}^4[\chi_2]}{L F_{10}^4[\chi_{12}]}$

= 0 else where

where $\chi_{12} = \begin{bmatrix} a_1, a_1, a_2, a_2, b_1, b_1, b_1, b_1, c_1, c_2, c_1, c_3; \Delta(n, b), \Delta(\frac{n}{2}, \frac{c+b}{2}), \\ \Delta(\frac{n}{2}, \frac{1+c+b}{2}), \Delta(n, c+b), \Delta(\frac{n}{2}, c+b+a), \Delta(\frac{n}{2}, \frac{1+c+b-a}{2}); X, Y, Z, T \end{bmatrix}$

©IJRASET: All Rights are Reserved

International Journal for Research in Applied Science & Engineering Technology (IJRASET)
ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.177
Volume 7 Issue IX, Sep 2019- Available at www.ijraset.com

III. SOLUTION OF INTEGRALS

One of the proofs is expressing the quadruple hypergeometric function in terms of equivalent series, in the integrand of the (2.1). We find that the integral becomes

$$\int_0^I U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}}F_1(a,1-a;c;\frac{U}{2})$$

$$F_1^4\left[a_1,a_1,a_1,a_2,b_1,b_1,b_1,b_1,c_1,c_2,c_3,c_4;(I-U)^nX,(I-U)^nY,(I-U)^nZ,(I-U)^nT\right]dU$$

or

$$\int_0^I U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}}F_1(a,1-a;c;\frac{U}{2})\sum_{p,q,r,s=0}^{\infty}A_{r,s}^{p,q}\left[(I-U)^nX,(I-U)^nY,(I-U)^nZ,(I-U)^nT\right]dU \tag{3.1}$$

where $A_{r,s}^{p,q}$ stands for the expression

$$\sum_{p,q,r,s=0}^{\infty}\frac{(a_1)_{p+q+r}(a_2)_s(b_1)_{p+q+r+s}}{p!q!r!s!(c_1)_p(c_2)_q(c_3)_r(c_4)_s}\left[(I-U)X\right]^p\left[(I-U)Y\right]^q\left[(I-U)Z\right]^r\left[(I-U)T\right]^s$$

We assume that the series is uniformly convergent in the region of integration, the inversion of integration and summation is infinite, then integral

$$\int_0^I U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}}F_1(a,1-a;c;\frac{U}{2})\,A_{r,s}^{p,q}\left[X^p(I-U)^{p\ n}Y^q(I-U)^{q\ n}Z^r(I-U)^mT^s(I-U)^{sn}\right]dU \tag{3.2}$$

$$=\sum_{p,q,r,s=0}^{\infty}A_{r,s}^{p,q}X^pY^qZ^rT^s\int_0^I U^{a-\frac{m+1}{2}}(I-U)^{b-\frac{m+1}{2}}F_1(a,1-a;c;\frac{U}{2}).\left[(I-U)^{n(p+q+r+s)}\right]dU$$

$$=\sum_{p,q,r,s=0}^{\infty}A_{r,s}^{p,q}X^pY^qZ^rT^s\int_0^I U^{a-\frac{m+1}{2}}(I-U)^{b+n(p+q+r+s)\frac{m+1}{2}}F_1(a,1-a;c;\frac{U}{2})dU$$

$$=\sum_{p,q,r,s=0}^{\infty}A_{r,s}^{p,q}X^pY^qZ^rT^s\frac{\Gamma_m(c)\Gamma_m(b+n(p+q+r+s))\,\Gamma_m\left(\frac{c+b+n+(p+q+r+s)}{2}\right)}{\Gamma_m(c+b+n(p+q+r+s))\,\Gamma_m\left[\frac{(c+b+n(p+q+r+s))-a}{2}\right]}$$

$$\cdot\frac{\Gamma_m\left(\frac{1+c+b+n(p+q+r+s)}{2}\right)}{\Gamma_m\left[\frac{(1+c+b+n(p+q+r+s))-a}{2}\right]} \tag{3.3}$$

$$=L\sum_{p,q,r,s=0}^{\infty}\frac{(a_1)_{p+q+r}(a_2)_s(b_1)_{p+q+r+s}}{p!q!r!s!(c_1)_p(c_2)_q(c_3)_r(c_4)_s}$$

$$\frac{(b)_{n(p+q+r+s)}\left(\frac{c+b}{2}\right)_{n(p+q+r+s)}\left(\frac{1+c+b}{2}\right)_{n(p+q+r+s)}}{(c+b)_{n(p+q+r+s)}\left[\frac{(c+b+a)}{2}\right]_{\frac{n}{2}(p+q+r+s)}\left[\frac{(1+c+b-a)}{2}\right]_{\frac{n}{2}(p+q+r+s)}}X^pY^qZ^rT^s$$

By applying the below mentioned result, we get

$$(a)k_1=k^{k1}\prod_{j=1}^{k}\left\{\frac{(\alpha+j-1)}{k}\right\}$$

©IJRASET: All Rights are Reserved

where k is a positive integral and non-negative, and finally after calculations, we arrive at the result (1).Similarly, other integrals from (2.2) to (2.10)are all proved. Therefore, ten direct results have been quoted and proved in a similar manner.

IV. CONCLUSION

This paper deals with classical special function and generalized hypergeometric function of matrix argument (for positive definite symmetric matrix and hermitian positive definite matrix in complex case). We derived various important formulas which have a wide range of applications in the field of Mathematical Sciences especially multivariable distribution theory.

REFERENCES

[1] M. Abramowitz, I.A Stegun, Handbook of mathematical functions, Dover publications, New York (1972).
[2] T.W. Anderson, An introduction to multivariate statistical analysis (2nd ed.), John Wiley & sons, New York (1984).
[3] H. Exton, Some Integral representation and transformations of hypergeometric function of four variables, Bull. Amer. Math. Soc., 14 (1973),132-140.
[4] B.Usha, A. Shoukat, An Integral containing Hypergeometric function, Advances in Computational Mathematics and its Applications, 2, No.2 (2012), 263-266.
[5] Rao S, Prajapati J, Shukla A. Wright Type Hypergeometric Function and Its Properties. Advances in Pure Mathematics. 2013; 3(3): 335-342.
[6] Sethi PL, Sidhu BS. Multivariate integral representation suggested by Laguerre and Jacobi polynomials of matrix argument. J. Vihnana Parshad Anusandhan Patrika. 2005; 48(2): 171-219.

©IJRASET: All Rights are Reserved



10.22214/IJRASET

45.98

IMPACT FACTOR: 7.129

IMPACT FACTOR: 7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)