



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 8 Issue: II Month of publication: February 2020

DOI: <http://doi.org/10.22214/ijraset.2020.2029>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

HPTLC Fingerprinting in the Standardization of Arthrum Plus Capsule: A Poly Herbal Formulation

Mrs. Parul Vasava¹, Dr. Nidhi N Chauhan², Mrs. Neha Chauhan³, Dr. Kishor Dholvani⁴

^{1, 2, 3, 4}Department of Pharmacognosy, Laxminarayan Dev college of Pharmacy, Bholav, Bharuch, Gujarat, India

Abstract: Polyherbal formulations are potential for their safety, cost, effectiveness and better acceptability than allopathic drugs. Quality assurance of herbal products may be ensured by proper quality control of herbal ingredients and by means of good manufacturing practice. Therapeutic efficacy of herbal medicine is exerted due to the bioactive constituents of plants. We have developed simple scheme for standardization of (ACP).

Prepared polyherbal formulation was standardized by TLC and HPTLC fingerprinting for deciding identity, purity and strength of polyherbal formulation. The present study was aimed to standardize a polyherbal formulation (Arthrum plus capsule) including traditional drug extracts such as vitex negundo, Boswellia serata, Pluchea lanceolata, yograj guggul. HPTLC fingerprinting profile showed different band patterns at different wavelength under short UV, at 254nm and 366nm, after derivatization with vanillin sulphuric acid spraying reagent unique Rf patterns were recorded. Conclusion: APC was authenticated according to pharmacopeial standards as its analysis was important to ensure the quality of drug.

Keywords: Arthrum plus capsule, Polyherbal formulation, HPTLC fingerprinting.

I. INTRODUCTION

Polyherbal formulations (PHFs) are now influential drug to traditional medicine practices in India. Quality assurance of Ayurvedic formulations is an important step to study chemical profile and safety of drug, as the standardisation of drug is emerging aspect in Ayurvedic drug preparation. [1,2] Leading population of world is now relies on phytotherapy for health care as per World Health Organization (WHO). WHO consider phytotherapy is safe, cost effective and more significant without any side effect [3,4]. Phytotherapy is now emerging with issues regarding their quality, safety and efficacy.

WHO has given specific guidelines, for the assessment of safety, effectiveness and quality of herbal drug [5]. Ayurvedic Pharmacopoeia of India and AYUSH also provides monograph on the preparation of Ayurvedic PHF, that aided in standardisation of drug. A polyherbal Ayurvedic formulation (Arthrum plus capsule) was prepared by four ingredients such as vitex negundo extract, Boswellia serata extract, Pluchea lanceolata extract, yograj guggul powder. The formulation will be used in the treatment of Rheumatoid Arthritis condition. polyherbal formulation was subjected for standardisation as per pharmacopeial procedures [6,7,8,9]

II. MATERIALS AND METHODS

Arthrum Plus Capsule consists of four ingredients viz., Boswellia serrata (Shallaki), Yograj Guggula, Pluchea lanceolata (Raasna), Vitex negundo (Nirgundi) [10]

The extract of Boswellia serrata (Shallaki), Pluchea lanceolata (Raasna), Vitex negundo (Nirgundi) and powder mixture of Yograj guggul, were procured from Prashant Pharmaceuticals, Rajpipala, India. While finished formulation Arthrum Plus capsule were formulated by Vital care Pvt. Ltd., Vadodara, India. All the extracts used in Arthrum Plus capsule were vacuum dried powder form.

All the reagents and instruments used in standardization and evaluation of Anti-Rheumatoid Arthritis activity of Arthrum Plus capsule were facilitated by Shri B.M. Shah College of Pharmaceutical Education and Research, Modasa.

All the reagents and instruments used in standardization and evaluation of Anti-Rheumatoid Arthritis activity of Arthrum Plus capsule were facilitated by Shri B.M. Shah College of Pharmaceutical Education and Research, Modasa.

A. HPTLC finger-printing of raw Material and Finished Product [11,12,15]

HPTLC is the most simple separation technique available today which gives better precision and accuracy with extreme flexibility for various steps (stationary phase, mobile phase, development technique and detection). The HPTLC was carried out using a Hamilton 100 µl HPTLC syringe, Camag Linomat V automatic spotting device, Camag twin trough chamber, Camag TLC Scanner-3, WINCAT integration software, aluminium sheet precoated with Silica Gel 60F254 (Merck), 0.2 mm thickness. HPTLC fingerprinting technique is useful to identify and to check the purity of raw herbal extracts as well as finished product. Henceforth it is very useful tool in standardizing process of raw herbal extracts and finished products.

B. Steps Involved in HPTLC Analysis

- 1) **Selection Of Plate And Adsorbent:** Precoated aluminium plates with Silica Gel 60F254 (E. Merck, India) of 10 x 10 cm and 0.2 mm thickness, were used for the detection. The plates were prewashed by methanol and activated at 60°C for 5 min prior to chromatography.
- 2) **Sample Solution:** Accurately weighed 50 mg of methanol extract of raw material was taken, dissolved in methanol and transferred to a 10 ml volumetric flask. This solution was further used for HPTLC finger-printing.
- 3) **Application Of Sample:** Sample application is the most critical step for obtaining good resolution for quantification in HPTLC. The automatic application devices are preferable. The most recent automatic device “CAMAG LINOMAT V” was used to apply 1 band of 6 mm width with different concentration of all of extract solution viz. 4,6,8,10,12 µl.
- 4) **Development:** The plate was developed in CAMAG glass twin-through chamber (10-10 cm) previously saturated with the solvent for 60 min (temperature 25.2 °C, relative humidity 40%). The development distance was 8 cm. Subsequently scanning was done. The mobile phase or solvent system for all the ingredients as well as Arthrum Plus Capsule which is given in the Table 1
- 5) **Detection:** The plate was scanned at UV 366 nm and 254 nm using CAMAG TLC Scanner-3 and LINOMAT-V. Rf value of each compound which were separated on plate and data of peak area of each band was recorded

Table 1: Solvent system for plant extract, and Arthrum Plus Capsule^[13,14,15]

Sample	Solvent system
Boswellia serrata.	Ethyl acetate: Hexane (1:1)
	Toluene: Ethyl acetate (9:1)
Vitex negundo	Toluene: Ethyl acetate (9:1)
	Toluene: Ethyl acetate (7:3)
	Toluene: Ethyl acetate: Formic Acid: Methanol (6:6:1.6:0.4)
	Benzene: Ethyl acetate (9:1)
Plueacha lanceolata	Toluene: Ethyl acetate: Formic Acid (5:5:1)
	Ethyl acetate: Hexane (1:1)
	Toluene: Ethyl acetate (9:1)
Yograj Guggula	Benzene: Ethyl acetate (9:1)
	Ethyl acetate: Hexane (1:1)
	Toluene: Ethyl acetate (9:1)
	Toluene: Ethyl acetate (7:3)
	Toluene: Ethyl acetate: Formic Acid: Methanol (6:6:1.6:0.4)

III. RESULT AND DISCUSSION

A. HPTLC Fingerprinting

Rf values and colour of the spots in chromatogram developed in different mobile system for Arthrum plus capsule were recorded (Table 2). TLC photo-documentation revealed presence of many phytoconstituents with different Rf values and HPTLC densitometric scan of the plates showed numerous bands under short UV. Densitometric scan at 254 nm revealed different peaks corresponding to different compounds in the APC, Compound with Rf 0.4(0.05 micro gm/ml) 0.24(1.5micro gm/ml), 0.74(4.45micro gm/ml), 0.34(0.04 micro gm/ml), 0.45(0.03micro gm/ml). It indicates the concentration of 11-keto-boswellic acid, Quercetin, piperine, gingerol, gallic acid were found in the different extract of Arthrum plus capsule by HPTLC method respectively.

(Table 2). Rf value of the sample

Track	Peak	Max. Rf	Peak Area	Concentration (µg/ml)	
Formulation	B. serrata .Ext	1	0.74	2632.9	3.8
	<i>V. negundo extract</i>		0.57	595.5	-
	Vitexin		0.56	555.4	-
	Yograj guggul Piperine	1	0.33	359.5	0.06
	Yograj guggul Gingerol	1	0.33	552.6	1.8
	Yograj guggul Guggulosterol		0.82	438.4	-
	Yograj guggul Gallic acid	1	0.4	423.1	0.05
B. serrata .Ext	1	0.24	6112.6	1.5	
<i>P.lanceolata</i>	1	0.74	2956.5	4.4	
<i>V. negundo extract</i>		0.56	105.4	-	
Vitexin		0.57	796.8	-	
Yograj guggul Piperine	1	0.34	333.7	0.04	
Yograj guggul Gingerol	1	0.33	929.3	5.6	
Yograj guggul Guggulosterol		0.82	602.4	-	
Yograj guggul Gallic acid	1	0.45	363	0.03	

B. Quantitative Estimation of 11-keto-boswellic acid in *B. serrata* Roxb by HPTLC method

1) Calibration curve for 11-keto-boswellic acid (STANDER)




Figure: 1A HPTLC plats of 11-keto-boswellic acid




Figure: 2 HPTLC Chromatogram and 3 Dimage of 11-keto-boswellic acid






Fig: 3 calibration curve of standard 11-keto-boswellic acid

2) *B. serrata* Roxb(11-keto-boswellic acid)(in formulation)



UV 254 nm UV 366nm 3D image

Fig.4 HPTLC plats and 3D image of Formulation, *B. serrata* extract and 11- keto boswellic acid




Formulation Extract of *B. Serrat* 11- keto-boswellic acid

Fig: 5 HPTLC Chromatogram of Formulation, *B. Serrat* and 11- keto-boswellic acid

C. *Quantitative estimation of quercetin P.lanceolata* by HPTLC method

1) *Calibration of Quercetin (Stander)*



Standard quercetin 3 D- image of the quercetin HPTLC chromatogram

Fig. 6 HPTLC plate, chromatogram and 3 D- image of the quercetin




Fig. 7 Calibration curve of Quercetin

2) *P.lanceolata* (Quercetin)(formulation)




Fig:8 HPTLC plate and 3D chromatogram of formulation, *P. Lanceolata* extracts with quercetin




Fig. 9 HPTLC Chromatogram of *P. lanceolata* extract, formulation and quercetin

D. Estimation of vitexin in vitex negundo extract and Formulation by HPTLC




Fig. 10 HPTLC plats and 3D HPTLC chromatogram of *vitex negundo*





Fig. 11 HPTLC Chromatogram of *v. negundo* extract and formulation


E. Quantitative estimation of Yograj Guggul powder by HPTLC method.

Quantitative estimation of piperin in Yograj Guggul powder by HPTLC method.

1) Calibration of Piperin



UV 254 nm UV 366 nm HPTLC chromatogram
Fig. 12 HPTLC Chromatogram and 3D image of Piperine



1 µL Piperine 2 µL Piperine 3 µL Piperine 4 µL Piperine 5 µL Piperine 6µLPiperine

Figure: 13HPTLC Chromatogram of Piperine






Fig. 14 Calibration curve of piperine

2) Yograj Guggul (Piperine)



UV 254 nm UV 366nm 3D HPTLC chromatograph

Fig. 15 HPTLC and 3D image of formulation, yograj guggul and piperine




Formulation yograj guggul Piperine

Fig. 16 Chromatogram of Yograj Guggul and formulation


F. Quantitative estimation of Gingerolin Yograj Guggul powder by HPTLC method.

1) Calibration of Gingerol



UV254nm UV366nm 3D image


Fig. 17 HPTLC plats and 3D chromatogram of gingerol



HPTLC Chromatograph of gingerol calibration curve of gingerol

Fig.18. HPTLC Chromatograph and calibration curve of gingerol

2) Zingiber officinales (gingerol)



UV254nm UV366nm HPTLC chromatogram

Fig: 19HPTLC plats,3 D chromatogram of Yograj guggul




Fig. 20 HPTLC Chromatogram of gingerol, Yograj guggul and formulation

G. Estimation of Guggulosterol in Extract of *Commiphora mukul* and Formulation by HPTLC




Fig. 21 HPTLC plats and 3D chromatogram of Guggulosterol




Fig 22 HPTLC chromatogram of Yograj Guggul extract, guggulosterol and formulation

H. Quantitative estimation of gallic acid in Yograj guggul powder by HPTLC method.

1) Calibration of Gallic acid






Fig. 23 HPTLC plates and 3-D Chromatograph of gallic acid



3 µL Gallic acid 4 µL Gallic acid 5 µL Gallic acid Calibration curve of Gallic acid


Fig. 24 HPTLC Chromatogram and calibration curve of standard Gallic acid

2) Amla, Baheda and Harde (Gallic acid)



UV 254nm UV 366nm 3D HPTLC chromatograph

Figure: 25 HPTLC plats and chromatogram of Yograj Guggul




Formulation

Triphala

Gallic acid

Fig. 26 HPTLC Chromatogram of Gallic acid, Formulation, Extract of Yograj Guggul



UV 254 nm

UV 366nm

3D HPTLC chromatograph

Fig. 27 HPTLC and 3D image of formulation, yograj guggul and piperine




Fig .28 Chromatogram of Yograj Guggul and formulation

I. Quantitative Estimation of Gingerolin Yograj Guggul powder by HPTLC method.

1) Calibration of gingerol





Fig.29 HPTLC plats and 3D chromatogram of gingerol



HPTLC Chromatograph of gingerol calibration curve of gingerol
 Fig. 30 HPTLC Chromatograph and calibration curve of gingerol

2) Zingiber Officinales (gingerol)




Fig: 31 HPTLC plats 3 D chromatogram of Yograj guggul




Fig. 32 HPTLC Chromatogram of gingerol, Yograj guggul and formulation

J. Estimation of Guggulosterol in Extract of Commiphora mukul and Formulation by HPTLC




Fig.33 HPTLC plats and 3D chromatogram of Guggulosterol




Fig: 34 HPTLC chromatogram of Yograj Guggul extract, guggulosterol and formulation

K. Quantitative Estimation of gallic acid in Yograj guggul powder by HPTLC method.

1) Calibration of Gallic acid






Fig.35 HPTLC plates and 3-D Chromatograph of gallic acid




4 μ L Gallic acid 5 μ L Gallic acid Calibration curve of Gallic acid
 Fig. 36 HPTLC Chromatogram and calibration curve of standard Gallic acid

2) Amla, Baheda and Harde (Gallic acid)



UV 254nm UV 366nm 3D HPTLC chromatogram
 Figure: 37 HPTLC plats and chromatogram of Yograj Guggul



Formulation Triphala Gallic acid
 Fig. 38 HPTLC Chromatogram of Gallic acid, Formulation, Extract of Yograj Guggul

IV. SUMMARY AND CONCLUSION

Standardization of polyherbal formulation is specified level to assure the quality and efficacy of drug. HPTLC fingerprinting also revealed active constituents at different Rf values. Result indicate the concentration of 11-keto- boswellic acid content were found in the extract of *B.serrata* and Arthrum plus capsule by HPTLC method was 0.33mcg/ml and 1.5 mcg/ml with max Rf value 0.24, in extract of *p.lanceolata* concentration of Quercetin was 3.8,4.4 mcg/ml with 0.74 Rf ,in extract of yograj guggul concentration of piperine 0.06 and 0,04 mcg/mlwith Rf 0.3, gingerol 1.8 and 5.6mcg/ml with Rf 0.33,gallic acid 0.05 and 0.03mcg/ml with 0.45 Rf respectively.

A. Conclusion

in conclusion Data suggested that Arthrum plus capsule and its ingredient were consistent with various quality and purity parameter such like HPTLC Analysis.

REFERANCE

- [1] Subramani Parasuraman, Gan Siaw Thing, and Sokkalingam Arumugam Dhanara Polyherbal formulation: Concept of ayurveda. *Pharmacogn Rev.* 2014 JulDec; 8(16): 73–80.
- [2] Arun Raj GR, Shailaja U, Prasanna N Rao, Ajayan S, Nivya P Thomas. Review on the contribution of Ura- Marunnu, a traditional baby care practice in southern India. *The Pharma Innovation* 2014; 2(11):42-70.
- [3] Kimmatkar N, Thawani V, Hingorani L, Khiyani R. Efficacy and tolerability of *Boswellia serrata* extract in treatment of osteoarthritis of knee-a randomized double blind placebo controlled trial. *Phytomed* 2003; 10:3-7.
- [4] Waddar S, Gopi KBJ, Rao PN, Raj AGR, Waddar S. Standardisation of Mulaka (*Raphenus sativus* Linn.) Kshara: a herbal alkaline preparation. *Journal of Pharmacognosy and Phytochemistry* 2014; 3(1):108-110.
- [5] Chaudhary RR. Herbal medicine for human health. World Health Organization, Geneva, CBS Publishers and distributors LTD; New Delhi, 1999. 5.
- [6] N.Thangarathinam, N.Jayshree, A.Vijay Metha, L. Ramanathan. Development, Standardization and Evaluation of a Polyherbal Syrup *Int. J. Pharm. Sci. Rev. Res.*, 20(2), May – Jun 2013; n° 25, 149-154
- [7] World Health Organization. WHO guidelines for assessing quality of herbal medicine with reference to contaminants and residue, Geneva, WHO Press; 2007.
- [8] Mukherjee PK,GMP for Indian system of Medicine, Business Horizons,New Delhi 2003,page:39-40.
- [9] Anonyms, WHO guidelines on safety monitoring of herbal medicines in pharmacovigilance system, 2004;5-6.
- [10] http://www.vitalcare.co.in/arthrum_plus_capsule.htm.
- [11] Sethi PD. High Performance Thin Layer Chromatography. 1st ed. New Delhi, CBS Publishers and Distributors; 1996.
- [12] Wager H, Baldt S, Zganski EM: In Plant drug analysis; A thin layer chromatography Atlas. Berlin Springer. 1984, pg 299.
- [13] Stahl I. Thin layer chromatography. A Laboratory Hand Book (student edition), Berlin, Springer-Verlag 1969; 52-86.
- [14] The Ayurvedic pharmacopoeia of India part-I ,vol III,First edition page :142-144.
- [15] Chandravansi Lowkesh, V.K Chetan kumar, " Quality characterisation and HPTLC fingerprinting of Vachyadi syrup: poly herbalformulation."The journal of phytopharmacology 2016;5(6):234-237.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)