



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 8 Issue: VI Month of publication: June 2020

DOI: http://doi.org/10.22214/ijraset.2020.6146

www.ijraset.com

Call: 🕥 08813907089 🔰 E-mail ID: ijraset@gmail.com

Developed Ratio Estimator for Estimating Population Mean from a Finite Population using Sample Size and Correlation

Shamsuddeen Ahmad Sabo¹, Ibrahim Zakariyya Musa², Yakubu Isyaku Kibiya³

^{1, 2, 3}Post Graduate Research Scholar, Mathematics Department, Mewar University, Chittorgarh, Rajasthan, India

Abstract: Estimation of the population mean is a crucial matter in our world activities today. In this paper a new improved ratio estimator for estimation of population mean from a finite population using SRSWOR by utilizing information on sample size and correlation of the auxiliary variable was proposed. The derivation of the Bias and mean square error (MSE) was done to the first order of approximation. The expression for the optimum Bias and minimum mean square error was evaluated and presented for the minimum value of a constant τ . A mathematical efficiency comparison was performed and shows that our proposed estimator is more efficient than the other estimators. The empirical study performed shows that our estimator has a smaller bias and MSE compared to the other estimators. The PRE of our proposed estimator over other estimators compared to the usual estimator is higher than all the other modified estimators. Hence our proposed estimator is better than the other estimators and should be given preference over the modified estimators when estimating a population means. Keywords: SRSWOR, Auxiliary variable, study variable, Bias, MSE

I. INTRODUCTION

There is no argument that the significance attached to the vital role played by sampling theory in statistics cannot be overemphasized considering its vast number of applications in different spheres of life. The application of sampling theory is not only limited to the sample selection, but also to the information about the population of its parameters. The main objective of sampling theory is to estimate the parameter of interest using some statistical properties as a bedrock. In many situations, it happens that information regarding some variables apart from the variables of interest (study variables) can easily be collected. This information is called auxiliary information which is formed by auxiliary variable(s). In survey sampling, if the auxiliary variable(s) are highly correlated and suitably used, the precision of estimators of the mean of the study variable is increased by reducing the variance of the estimators of the population parameters [4]. In estimating population parameters one can propose a different number of estimators. One can have used a sample mean or sample median or any other sample statistic [1]. To estimate a population mean, the most suitable estimator is the sample mean since the best estimator of every population parameter is its corresponding statistic. But some sample statistic tends to yield estimate with a large sampling variance when estimating its corresponding parameter. Hence one has to look for a better estimator that will perform better than the other estimator.

Different techniques for estimation of population mean to exist in the literature, but the most widely used ones are ratio, product, and regression estimators. When Regression line of study variable and auxiliary variable(s) pass through the origin and their correlation is positively high then Ratio estimators are applied. Likewise, Product estimators can be applicable when the regression line of the two variables doesn't pass through the origin and their correlation is negatively low. Situations may arise when there exists a correlation between the study variable and auxiliary variable(s) but nothing can be said about the direction of the correlation (either positive or negative) and the regression line is linear in this case regression estimator is used.

Let us consider a variable under study as Y which is correlated with its auxiliary variable X. We obtain the observations x_i and y_i corresponding to X and Y respectively for each of the sampling unit using simple random sampling without replacement (SRSWOR). The usual estimator for Y is its corresponding sample mean $\bar{y}_r = \frac{1}{n} \sum_{i=1}^{N} \bar{y}_i$ which is unbiased but has a large sampling variance, with a variance $V(\bar{y}_r) = \theta \bar{Y}^2 C_y^2$. [7] Cochran made his first contribution by coming up with a ratio estimator of the form $\bar{Y}_r = \bar{y}_r \frac{\bar{X}}{\bar{x}}$ in order to reduce the effect of having a large sampling variance. The estimator given by Cochran is biased and its bias and mean square error (MSE) are:

A lot of researchers have contributed to the literature by proposing an estimator which will be more efficient in estimating the population means thereby modifying the estimator given by Cochran using some parameters like the coefficient of variation, skewness, kurtosis, median, correlation coefficient, number of sample size, minimum, maximum, and their combinations [13] [14]



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 8 Issue VI June 2020- Available at www.ijraset.com

[10] [8] [12] [2] [1] [9] [11]. To the best of our knowledge, no previous research used the information from sample size and correlation coefficient together. Table (1) below gives a summary of some estimators with their constants, bias, and MSE which we called modified estimators throughout this paper unless otherwise stated. The modified estimators in the table (1), were used in making a comparison with our proposed estimator.

| ESTIMATORS | CONSTANTS | BIAS | MSE |
|---|--|---|--|
| | | | |
| $\widehat{Y}_1 = \overline{y} \left(\frac{\overline{X} + C_x}{\overline{x} + C_x} \right)$ | $\gamma_1 = \frac{\overline{X}}{\overline{X} + C_x}$ | $\theta \bar{Y}(\gamma_1^2 C_x^2 - \gamma_1 \rho C_x C_y)$ | $\theta \bar{Y}^2 (\mathcal{C}_y^2 + \gamma_1^2 \mathcal{C}_x^2 - 2\gamma_1 \rho \mathcal{C}_x \mathcal{C}_y)$ |
| $\widehat{Y}_2 = \overline{y} (\frac{\overline{X}C_x + \beta_2}{\overline{x}C_x + \beta_2})$ | $\gamma_2 = \frac{\overline{X}C_x}{\overline{X}C_x + \beta_2}$ | $\theta \bar{Y}(\gamma_2^2 C_x^2 - \gamma_2 \rho C_x C_y)$ | $\theta \bar{Y}^2 (\mathcal{C}_y^2 + \gamma_2^2 \mathcal{C}_x^2 - 2\gamma_2 \rho \mathcal{C}_x \mathcal{C}_y)$ |
| $\widehat{Y}_3 = \overline{y} (\frac{\overline{X} + \rho}{\overline{x} + \rho})$ | $\gamma_3 = \frac{\bar{X}}{\bar{X} + \rho}$ | $\theta \tilde{Y}(\gamma_3^2 C_x^2 - \gamma_3 \rho C_x C_y)$ | $\theta \tilde{Y}^2 (C_y^2 + \gamma_3^2 C_x^2 - 2\gamma_3 \rho C_x C_y)$ |
| $\widehat{\overline{Y}}_4 = \overline{y}(\frac{\overline{X} + \beta_2}{\overline{x} + \beta_2})$ | $\gamma_4 = \frac{\overline{X}}{\overline{X} + \beta_3}$ | $\theta \tilde{Y}(\gamma_4^2 C_x^2 - \gamma_4 \rho C_x C_y)$ | $\theta \tilde{Y}^2 (C_y^2 + \gamma_4^2 C_x^2 - 2\gamma_4 \rho C_x C_y)$ |
| $\widehat{Y}_5 = \overline{y}(\frac{\overline{X} + \beta_1}{\overline{x} + \beta_1})$ | $\gamma_5 = \frac{\overline{X}}{\overline{X} + \beta_1}$ | $\theta \tilde{Y}(\gamma_5^2 C_x^2 - \gamma_5 \rho C_x C_y)$ | $\theta \tilde{Y}^2 (C_y^2 + \gamma_5^2 C_x^2 - 2\gamma_5 \rho C_x C_y)$ |
| $\widehat{\widehat{Y}_6} = \overline{y}(\frac{\overline{X} + M_d}{\overline{x} + M_d})$ | $\gamma_6 = \frac{\overline{X}}{\overline{X} + M_d}$ | $\theta \overline{Y}(\gamma_b^2 C_x^2 - \gamma_b \rho C_x C_y)$ | $\theta \tilde{Y}^2 (C_y^2 + \gamma_b^2 C_x^2 - 2\gamma_b \rho C_x C_y)$ |
| $\widehat{\overline{Y}_7} = \overline{y}(\frac{\overline{X} + n}{\overline{x} + n})$ | $\gamma_7 = \frac{\overline{X}}{\overline{X} + n}$ | $\theta \overline{Y}(\gamma_7^2 C_x^2 - \gamma_7 \rho C_x C_y)$ | $\theta \tilde{Y}^2 (\mathcal{C}_y^2 + \gamma_7^2 \mathcal{C}_x^2 - 2\gamma_7 \rho \mathcal{C}_x \mathcal{C}_y)$ |
| $\widehat{Y}_{\alpha} = \overline{v} \{ \alpha + (1 - \alpha) \left(\frac{\overline{X} + n}{n} \right) \}$ | $v_{+} = \frac{\overline{X}}{\overline{X}}$ | $\theta \overline{Y} (\gamma_8^2 C_x^2 - \gamma_8 \rho C_x C_y$ | $\theta \overline{Y}^2 (C_y^2 + \gamma_8^2 C_x^2 - 2\gamma_8 \rho C_x C_y + \alpha^2 \gamma_8^2 C_x^2$ |
| $x_8 = y(u + (1 - u))(\bar{x} + n)^3$ | $\sqrt{X} = \sqrt{X} + n$ | $+ \alpha \gamma_8^2 C_x^2$) | $+ 2\alpha\gamma_8\rho C_x C_y$ |
| | | | $-2\alpha\gamma_8\rho\gamma_8^2C_x^2)$ |

Table (1) Modified estimators with their corresponding Constants, Bias, and MSE

Where: $\theta = \frac{1-f}{n}$ and $f = \frac{n}{N}$

II. METHODOLOGY OF OUR PROPOSED ESTIMATOR

Motivated by [1] [2] and [13] We proposed the following estimator:

$$\widehat{Y}_{p} = \overline{y} \left\{ \alpha + (1 - \alpha) \left(\frac{X\rho + n}{\overline{x}\rho + n} \right) \right\} - - - - - (3)$$
Define; $\varepsilon_{0} = \frac{\overline{y}}{\overline{y}} - 1$, $\varepsilon_{1} = \frac{\overline{x}}{\overline{x}} - 1$, $\varepsilon_{2} = \frac{s_{y}^{2}}{s_{y}^{2}} - 1$, $\varepsilon_{3} = \frac{s_{x}^{2}}{s_{y}^{2}} - 1$, $\varepsilon_{4} = \frac{s_{xy}}{s_{xy}} - 1$
Such that;

And

$$E[\varepsilon_0] = [\varepsilon_1] = E[\varepsilon_2] = E[\varepsilon_3] = E[\varepsilon_4] = 0$$

$$E[\varepsilon_0^2] = \theta C_y^2; E[\varepsilon_1^2] = \theta C_x^2; E[\varepsilon_0 \varepsilon_1] = \theta \rho C_x C_y$$

Where $\theta = \frac{1-f}{n}$ It implies that;

$$\overline{y} = \overline{Y}(1 + \varepsilon_0)$$
 and $\overline{x} = \overline{X}(1 + \varepsilon_1)$

Putting these values in the propose estimator in (1) above gives;

$$\widehat{Y}_{p} = \overline{Y}(1-\varepsilon_{0})\left\{\alpha + (1-\alpha)\left(\frac{X\rho + n}{\overline{X}(1-\varepsilon_{1})\rho + n}\right)\right\}$$

 $\widehat{Y}_p = \overline{Y}(1 - \varepsilon_0)\{\alpha + (1 - \alpha)(1 + \gamma \varepsilon_1)^{-1}\}, \quad \text{Where } \gamma = \frac{\overline{X}\rho}{\overline{x}\rho + n}$ Using the fact that; $(1 + \gamma \varepsilon_1)^{-1} = 1 - x + x^2 - x^3 + \dots + \infty$

$$\widehat{Y}_p = \overline{Y}(1 - \varepsilon_0)\{\alpha + (1 - \alpha)(1 - \gamma \varepsilon_1 + \gamma^2 \varepsilon_1^2 - \gamma^3 \varepsilon_1^3 + \cdots)\}$$

Taking the expansion up to the first order of approximation and expanding through gives;

$$\widehat{Y_p} = \overline{Y}\{(1 - \gamma\varepsilon_1 + \alpha\gamma\varepsilon_1 + \gamma^2\varepsilon_1^2 - \alpha\gamma^2\varepsilon_1^2 + \varepsilon_0 - \gamma\varepsilon_0\varepsilon_1 + \alpha\gamma\varepsilon_0\varepsilon_1)\}$$

Taking the expectation gives;

$$E[\bar{Y}_p] = E[\bar{Y}\{(1 - \gamma\varepsilon_1 + \alpha\gamma\varepsilon_1 + \gamma^2\varepsilon_1^2 - \alpha\gamma^2\varepsilon_1^2 + \varepsilon_0 - \gamma\varepsilon_0\varepsilon_1 + \alpha\gamma\varepsilon_0\varepsilon_1)\}]$$
$$E[\bar{Y}_p] = \bar{Y}\{(1 - \gamma\theta\rho C_x C_y + \gamma^2\theta C_x^2 + \alpha\gamma\theta\rho C_x C_y - \alpha\gamma^2\theta C_x^2)\}$$

 $\operatorname{Bias}\left(\widehat{Y_p}\right) = E\left[\widehat{Y_p}\right] - \overline{Y}$



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429

Volume 8 Issue VI June 2020- Available at www.ijraset.com

 $MSE(\widehat{Y_p}) =$

Expanding and taking the expansion up to the first order approximation yield;

$$\theta \overline{Y}^2 \Big(C_y^2 + \gamma^2 C_x^2 - 2\gamma \rho C_x C_y + \alpha^2 \gamma^2 C_x^2 + 2\alpha \gamma \rho C_x C_y - 2\alpha \gamma^2 C_x^2 \Big) - - - - - (5)$$

The Min. $MSE(\widehat{Y_p})$ is found by finding the optimum value of alpha (α), which is found by differentiating equation (5) w.r.t. α and equating to zero. Which yield;

 $\frac{\partial MSE(\widehat{Y_p})}{\partial \alpha} = 0$

$$\alpha \gamma^2 C_x^2 = \gamma^2 C_x^2 - \gamma \rho C_x C_y$$

Which is minimum for;

$$\alpha = \frac{\gamma^2 C_x^2 - \gamma \rho C_x C_y}{\gamma^2 C_x^2}$$

To find the Min. $MSE(\widehat{Y}_p)$ we substitute the value of α into equation (5) to yield;

Where $\tau = \left(\frac{\gamma^2 c_x^2 - \gamma \rho c_x c_y}{\gamma c_x}\right)^2 \ge 0$

To find the optimum Bias, we substitute the value of α into equation (4) to yield;

$$Bias_{opt}\left(\widehat{Y}_{p}\right) = \theta \overline{Y}\left(\gamma^{2}C_{x}^{2} - \gamma\rho C_{x}C_{y} - \tau\right) - -(7)$$

Where $\tau = \left(\frac{\gamma^2 C_x^2 - \gamma \rho C_x C_y}{\gamma C_x}\right)^2 \ge 0$

III. MATHEMATICAL EFFICIENCY COMPARISON BETWEEN OUR PROPOSED ESTIMATOR AND OTHER ESTIMATORS

The efficiency of our proposed estimator in (1) above is compared with that of existing and modified estimators

1) Comparison with the usual sample mean(\bar{y}_r)

Our proposed estimator is better than the sample mean (\bar{y}_r) if :

$$\begin{split} MSE\left(\widehat{Y_p}\right) &\leq V(\overline{y}_r)\\ \text{That is: } V(\overline{y}_r) - MSE\left(\widehat{Y_p}\right) &\geq 0 \end{split}$$

$$\begin{split} \theta \bar{Y}^2 C_y^2 &- \theta \bar{Y}^2 \big(C_y^2 + \gamma^2 C_x^2 - 2\gamma \rho C_x C_y - \tau \big) \\ &- \theta \bar{Y}^2 \big(\gamma^2 C_x^2 - (2\gamma \rho C_x C_y + \tau \big) > 0 \end{split}$$

If $\gamma^2 C_x^2 < 2\gamma \rho C_x C_y + \tau$

2) Comparison with Cochran estimator($\widehat{Y_r}$)

Our proposed estimator is better than the estimator given by Cochran if:

$$\begin{split} \operatorname{MSE}(\widehat{Y}_{r}) - \operatorname{MSE}(\widehat{Y}_{p}) &= \theta \overline{Y}^{2} \left(C_{y}^{2} + C_{x}^{2} - 2\rho C_{x} C_{y} \right) - \theta \overline{Y}^{2} \left(C_{y}^{2} + \gamma^{2} C_{x}^{2} - 2\gamma \rho C_{x} C_{y} - \tau \right) \\ &= \theta \overline{Y}^{2} \left((C_{x} - \rho C_{y})^{2} - (\gamma C_{x} - \rho C_{y})^{2} + \tau \right) > 0 \\ \operatorname{If} C_{x} - \rho C_{y})^{2} > (\gamma C_{x} - \rho C_{y})^{2} \end{split}$$

3) Comparison with other modified estimators $\widehat{(Y_j)} \forall j = 1, 2, 3, 4, 5, 6, and 7$ Our proposed estimator performs better than the modified estimators if: $MSE(\widehat{Y_j}) - Min.MSE(\widehat{Y_p}) = \theta \overline{Y}^2 (C_y^2 + \gamma_j^2 C_x^2 - 2\gamma_j \rho C_x C_y) - \theta \overline{Y}^2 (C_y^2 + \gamma_p^2 C_x^2 - 2\gamma_p \rho C_x C_y - \tau)$ $= \theta \overline{Y}^2 [C_x^2 (\gamma_j^2 - \gamma_p^2) - 2\rho C_x C_y (\gamma_j - \gamma_p) + \tau] > 0$ If $C_x^2 (\gamma_j^2 - \gamma_p^2) - 2\rho C_x C_y (\gamma_j - \gamma_p) > 0$ If: $\rho \leq \frac{\gamma_j + \gamma_p}{2} \cdot \frac{C_x}{C_y}$





4)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 8 Issue VI June 2020- Available at www.ijraset.com

Comparison with \widehat{Y}_8 estimator our proposed estimator is better than \widehat{Y}_8 if $MSE(\widehat{Y}_8) - MSE(\widehat{Y}_p)$ $\theta \overline{Y}^2 (C_y^2 + \gamma_8^2 C_x^2 - 2\gamma_8 \rho C_x C_y - \tau) - \theta \overline{Y}^2 (C_y^2 + \gamma_p^2 C_x^2 - 2\gamma_p \rho C_x C_y - \tau)$ $\theta \overline{Y}^2 [C_x^2 (\gamma_8^2 - \gamma_p^2) - 2\gamma \rho C_x C_y (\gamma_8 - \gamma_p) - (\tau_8 - \tau_p)] > 0$ If:

 $C_x^2 \big(\gamma_8^2 - \gamma_p^2 \big) - 2 \gamma \rho C_x C_y \big(\gamma_8 - \gamma_p \big) > \big(\tau_8 - \tau_p \big)$

IV. RESULT AND DISCUSSION

To buttress our point that our proposed estimator performs better than the modified estimators, in addition to the mathematical efficiency comparison an empirical study was performed. We used data from [5] [15] and [16]. The summary of the data is given in table (2). Using the data in the table (2), the value of the constants, bias, and MSE were obtained and presented in table (3&4). Table (5) gives the summary of the efficiency of our proposed estimator relative to the modified estimators for population 1, 2, 3 &4. The Summary of Percentage relative efficiency of our proposed estimator over modified estimators compared to the usual estimator $\overline{y_r}$ for population 1,2,3, and 4 is presented in table (6). For clear vision, we use figures 1, 2, 3 & 4 to show the representation of the numerical values obtained in table 3 while fig 5, 6 7 & 8 was used to represent the numerical values obtained in the table (6).

| POPULATION | MURTHY | | MUKHOP | ADHYAY | COCHRAN | | |
|----------------|-----------------------|-----------------------|-----------------------|-----------------------|---------|-----------------------|--|
| PARAMETERS | <i>P</i> ₁ | P ₂ | P ₃ | P ₄ | P_5 | P ₆ | |
| Ν | 80 | 80 | 40 | 40 | 49 | 10 | |
| n | 20 | 20 | 8 | 8 | 5 | 4 | |
| F | 0.25 | 0.25 | 0.2 | 0.2 | 0.102 | 0.102 | |
| \overline{Y} | 51.8264 | 51.8264 | 50.7858 | 50.7858 | 127.796 | 101.1 | |
| \overline{X} | 11.2646 | 2.8513 | 2.3033 | 9.4543 | 103.143 | 58.8 | |
| ρ | 0.9413 | 0.9150 | 0.8006 | 0.8349 | 0.982 | 0.6515 | |
| Sy | 18.3566 | 18.3566 | 16.7352 | 16.7352 | 123.121 | 14.6523 | |
| S _x | 8.4561 | 2.7043 | 1.9360 | 6.3869 | 104.405 | 7.5339 | |
| Cy | 0.3542 | 0.3542 | 0.3295 | 0.3295 | 0.963 | 0.1449 | |
| C _x | 0.7507 | 0.9485 | 0.8406 | 0.6756 | 1.012 | 0.1281 | |
| C_y^2 | 0.1255 | 0.1255 | 0.1086 | 0.1086 | 0.927 | 0.0210 | |
| C_x^2 | 0.5635 | 0.8996 | 0.7065 | 0.4564 | 1.024 | 0.0164 | |
| β_1 | 1.0500 | 1.3006 | 0.9740 | 0.8799 | 4.777 | 0.5764 | |
| β_2 | -0.0634 | 0.6977 | -0.5344 | -0.4622 | 7.511 | 0.3814 | |
| M_d | 7.5750 | 1.4800 | 1.250 | 7.0700 | 64 | 58 | |

Table (2): summary of the data used in assessing our comparison

| Table (3): Constants of the modified Est | mators |
|--|--------|
|--|--------|

| $\widehat{Y_j}$ | <i>P</i> ₁ | <i>P</i> ₂ | <i>P</i> ₃ | P ₄ | P ₅ | P ₆ | Constants |
|---|-----------------------|-----------------------|-----------------------|----------------|----------------|----------------|--|
| $\widehat{Y_1} = \overline{y}(\frac{\overline{X} + C_x}{\overline{x} + C_x})$ | 0.9375 | 0.7504 | 0.7326 | 0.9333 | 0.9903 | 0.9978 | $\gamma_1 = \frac{\bar{X}}{\bar{X} + C_x}$ |
| $\widehat{Y}_2 = \overline{y}(\frac{\overline{X}C_x + \beta_2}{\overline{X}C_x + \beta_2})$ | 1.0075 | 0.7949 | 1.3813 | 1.0780 | 0.9329 | 0.9518 | $\gamma_2 = \frac{\overline{X}C_x}{\overline{X}C_x + \beta_2}$ |
| $\widehat{Y}_3 = \overline{y} (\frac{\overline{X} + \rho}{\overline{x} + \rho})$ | 0.9229 | 0.7571 | 0.7421 | 0.9189 | 0.9906 | 0.9890 | $\gamma_3 = \frac{\bar{X}}{\bar{X} + \rho}$ |
| $\widehat{Y}_4 = \overline{y}(\frac{\overline{X} + \beta_2}{\overline{x} + \beta_2})$ | 1.0057 | 0.8034 | 1.3021 | 1.0514 | 0.9321 | 0.9936 | $\gamma_4 = \frac{\overline{X}}{\overline{X} + \beta_3}$ |
| $\widehat{Y}_5 = \overline{y}(\frac{\overline{X} + \beta_1}{\overline{x} + \beta_1})$ | 0.9147 | 0.6868 | 0.7028 | 0.9149 | 0.9557 | 0.9903 | $\gamma_5 = \frac{\bar{X}}{\bar{X} + \beta_1}$ |
| $\widehat{Y}_6 = \overline{y}(\frac{\overline{X} + M_d}{\overline{x} + M_d})$ | 0.5979 | 0.6583 | 0.6482 | 0.5722 | 0.6171 | 0.5034 | $\gamma_6 = \frac{\bar{X}}{\bar{X} + M_d}$ |
| $\widehat{Y_7} = \overline{y}(\frac{\overline{X} + n}{\overline{x} + n})$ | 0.3606 | 0.1248 | 0.2235 | 0.5417 | 0.9538 | 0.9363 | $\gamma_7 = \frac{\bar{X}}{\bar{X} + n}$ |
| $\widehat{Y}_8 = \overline{y} \{ \alpha + (1 - \alpha) \left(\frac{\overline{X} + n}{\overline{x} + n} \right) \}$ | 0.3603 | 0.1248 | 0.2235 | 0.5417 | 0.9538 | 0.9363 | $\gamma_8 = \frac{\overline{X}}{\overline{X} + n}$ |
| $\widehat{Y_p} = \overline{y}\{\alpha + (1 - \alpha)\left(\frac{\overline{X}\rho + n}{\overline{x}\rho + n}\right)\}$ | 0.3465 | 0.1955 | 0.1813 | 0.4966 | 0.9530 | 0.9055 | $\gamma_p = \frac{\bar{X}\rho}{\bar{X}\rho + n}$ |



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 8 Issue VI June 2020- Available at www.ijraset.com

| Estimator | Population 1 | | Population 2 | | Population 3 | | Population 4 | |
|--------------------------|--------------|---------|--------------|---------|--------------|----------|--------------|---------|
| | BIAS | MSE | BIAS | MSE | BIAS | MSE | BIAS | MSE |
| $\overline{\mathcal{Y}}$ | 0 | 12.6409 | 0 | 12.6409 | 0 | 28.0101 | 0 | 28.0101 |
| \widehat{Y}_r | 0.6087 | 18.9764 | 1.1509 | 41.3171 | 2.4621 | 95.8438 | 1.3740 | 49.8521 |
| \widehat{Y}_1 | 0.5065 | 15.2496 | 0.5362 | 17.1936 | 1.1010 | 42.0409 | 1.1376 | 41.0305 |
| \widehat{Y}_2 | 0.6215 | 19.4498 | 0.6299 | 20.6686 | 5.2909 | 217.7358 | 1.6759 | 61.4365 |
| $\widehat{Y_3}$ | 0.4837 | 14.4438 | 0.5500 | 17.7073 | 1.1407 | 43.5111 | 1.0899 | 39.3070 |
| $\widehat{Y_4}$ | 0.6184 | 19.3390 | 0.6485 | 21.3737 | 4.6175 | 188.0493 | 1.5693 | 57.2840 |
| \widehat{Y}_5 | 0.4715 | 14.0208 | 0.4144 | 12.5502 | 0.9807 | 37.6305 | 1.0762 | 38.7911 |
| $\widehat{Y_6}$ | 0.1007 | 2.7800 | 0.3644 | 11.1401 | 0.7780 | 30.4603 | 0.2184 | 11.6580 |
| $\widehat{Y_7}$ | -0.0332 | 1.8332 | -0.0474 | 6.3154 | -0.0721 | 11.5806 | 0.1686 | 10.6005 |
| $\widehat{Y_8}$ | -0.0410 | 1.4303 | -0.1300 | 2.0377 | -0.1011 | 10.1074 | 0.1270 | 8.4858 |
| $\widehat{Y_p}$ | -0.0472 | 1.4001 | -0.1353 | 1.6619 | -0.1422 | 9.8792 | -0.2031 | 8.3824 |

Table (4): Summary of the Bias and MSE of the proposed and usual estimators

Table (5): Summary of Relative efficiency of the proposed estimator compared to another estimator

| Estimators($\widehat{\overline{Y}_{j}}$) | $Efficiency \ e \ (\widehat{Y_j}/\widehat{Y_p}) = MSE(\widehat{Y_p})/MSE(\widehat{Y_j})$ | | | | |
|--|--|--------------|--------------|--------------|--|
| | Population 1 | Population 2 | Population 3 | Population 4 | |
| $\widehat{Y_r}$ | 0.0738 | 0.0402 | 0.1031 | 0.1681 | |
| $\widehat{Y_{1}}$ | 0.0918 | 0.0967 | 0.2350 | 0.20430 | |
| \widehat{Y}_2 | 0.0720 | 0.0804 | 0.0454 | 0.1364 | |
| $\widehat{Y_3}$ | 0.0969 | 0.0939 | 0.2271 | 0.2126 | |
| $\widehat{Y_4}$ | 0.0724 | 0.0778 | 0.0525 | 0.1463 | |
| \widehat{Y}_5 | 0.0999 | 0.1324 | 0.2625 | 0.2161 | |
| $\widehat{Y_6}$ | 0.5036 | 0.1492 | 0.3243 | 0.7190 | |
| $\widehat{Y_7}$ | 0.7637 | 0.2632 | 0.8531 | 0.7865 | |
| $\widehat{Y_8}$ | 0.9789 | 0.8156 | 0.9774 | 0.9878 | |

Table (6): Summary of Percentage relative efficiency of proposed estimators over modified estimators compared to the estimator \bar{y} for population 1,2,3, and 4

| | | 1 1 | , , , , | | | | |
|-------------------------------|---|--------------|--------------|--------------|---|--|--|
| Estimators($\widehat{Y_j}$) | $PRE = V(\overline{y}) / MSE(\widehat{Y_{J}}) \times 100$ | | | | | | |
| | Population 1 | Population 2 | Population 3 | Population 4 | - | | |
| $\widehat{Y_r}$ | 66.6138 | 30.5874 | 29.2247 | 56.1864 | | | |
| \widehat{Y}_1 | 82.9129 | 73.5210 | 66.6258 | 68.2590 | | | |
| $\widehat{Y_2}$ | 64.9924 | 61.1599 | 12.8643 | 45.5920 | | | |
| \widehat{Y}_3 | 87.5178 | 71.3881 | 64.3746 | 71.2598 | | | |
| $\widehat{Y_4}$ | 65.1999 | 59.1423 | 14.8951 | 48.8969 | | | |
| \widehat{Y}_5 | 90.1582 | 100.7227 | 74.4346 | 72.2075 | | | |
| $\widehat{Y_6}$ | 454.7086 | 113.4721 | 91.9561 | 240.2651 | | | |
| $\widehat{Y_7}$ | 689.5528 | 200.1599 | 277.1247 | 330.0820 | | | |
| \widehat{Y}_8 | 883.7936 | 620.3515 | 277.1247 | 330.0820 | | | |
| $\widehat{Y_p}$ | 902.8569 | 760.6294 | 283.5260 | 334.1537 | - | | |
| | | | | | | | |



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 8 Issue VI June 2020- Available at www.ijraset.com

Fig1

MSE of modified estimators for popn 1

















ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 8 Issue VI June 2020- Available at www.ijraset.com

MSE of modified estimators for popn 4



Fig 4: MSE of modified and existed estimators corresponding to population 1



PRE of modified estimators for popn 1





PRE of modified estimators for popn 2





PRE of modified estimators for popn 3





ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 8 Issue VI June 2020- Available at www.ijraset.com

Fig 8

W Q

PRE of modified estimators for popn 4

V. CONCLUSION

In this paper, we proposed a new modified ratio class of estimators for the estimation of population mean under simple random sampling without replacement (SRSWOR). The derivation of the Bias and mean square error (MSE) was done to the first order of approximation. The expression for the optimum Bias and minimum mean square error was evaluated and presented for the minimum value of a constant τ . The mathematical efficiency comparison shows that our proposed estimator is more efficient than the other estimators. The empirical study performed shows that our estimator has a smaller bias and MSE compared to the other estimators. The PRE of our proposed estimator over other estimators compared to the usual estimator is higher than all the other modified estimators. Hence our proposed estimator is better than the other estimators and should be given preference over the modified estimators when estimating the population mean.

REFERENCES

- Rajesh Kumar Gupta, S.K Yadav (2018) "improved estimation of population mean using information on size of sample". American journal of mathematics and Statistics 8(2):27-35
- [2] Mohammad Jerajuddin, Jai Kishun (2016) "Modified Ratio Estimation for Population Mean using size of sample, selected from population". IJSRSET 2,2,10-16
- [3] Arijit Chaudhuri, Horst Stenger (2005) "Survey Sampling and Methods" 2nd edition, Chapman & Hall/CRC
- [4] Sarjindav Singh(2003) "Advanced sampling theory with applications; How Michael selected Amy " Springer. Science +Business media, B.V vol I-I
- [5] William G. Chocran (1977) "Sampling Techniques" 3rd edition, John Wiley and Sons
- [6] P.V. Sukhatme, B.V. Sukhatme, S.Sukhatme and C.Asok, "Sampling theory of Surveys with applications".
- [7] Chocran W.G(1940) "The estimation of the yields of the cereal experiments by sampling for the ratio of Grain to total produce", The Journal of Agric science, 30(2) 262-275
- [8] Kadillar C and Cingi H. (2006) "New Ratio estimations using correlation coefficients". Interstat 4, PP.1-11
- [9] Gupta S. and Shabbir J. (2007) "on the use of transformed auxiliary variables in estimating population mean by using two auxiliary variables" Journal of Statistical planning and inference 137: 1606-1611
- [10] Sisodia B.V.S. and Divivedi V.K. (1981) "A modified ratio estimator using coefficient of variation of auxiliary variable" Journal of Indian society of Agricultural statistics. 33,13-18
- [11] Subramani J. and kumar Pandiyan G. (2013) "New modified Ratio estimation of population mean when median of auxiliary variable is known" Rak. Journal of statistical operations research vol IX number 2, 2013, PP 137-145
- [12] Yan Z. & Tian B. (2010) "Ratio method to the mean estimation using coefficient of skewness of auxiliary variable" ICICA, Part-II, CCIS 106, pp.103-110.
- [13] Singh H.P. & Tailor R. (2003) "Use of known correlation coefficient in estimating the finite population Means" Statistics in Transition 6 (4), 555-560.
- [14] Upadhyaya L.N. & Singh H.P. (1999) "Use of transformed auxiliary variable in estimating the finite population means" Biometrical Journal 41(5), 627-636.
- [15] Murthy M.N. (1967) "Sampling theory and Methods" Statistical publishing society, Calcutta.
- [16] Mukhopadhyay P. (2009) "Theory and Methods of Survey sampling" 2nd edition. PHI Learning pvt. Ltd., New Delhi.











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24*7 Support on Whatsapp)