



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 8 Issue: XII Month of publication: December 2020 DOI: https://doi.org/10.22214/ijraset.2020.32694

www.ijraset.com

Call: 🕥 08813907089 🔰 E-mail ID: ijraset@gmail.com



Regeneration in *Brassica Juncea L.Crez.* (Indian Mustard)

Raj kumar¹, Aniket Kumar¹, Gaurav Krishna², Lavi Jaiswal³

¹Naini Agricultural Institute, Sam Higginbottom University of Agriculture Technology and Sciences 211007. ²Proprietor, Allahabad Advance Agri Solutions, Daragang, Allahabad -211006. ³Department of genetics and plant breeding, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi, 221005.

Abstract: An attempt to regeneration of the Indian mustard (Brassica juncea L.Crez.) in which hypocotyls and cotyledon are used. A quality seed are standardized by surface sterilization using 1 to 2% Nacl solution for 10 min. Sterilized seed are transfer to nutrient medium (MS media ,Cytokinin), for germination in presence of aseptic condition. We get 5 to 12 cm seedlings in bottle under controlled condition. Select the healthy seedling and separate hypocotyls and cotyledon under laminar air flow (LAR). Again the piece of hypocotyls and cotyledons are transferred to cytokinin known 0.2% medium derived from the MS(Murashige and Skoog) medium for regeneration. And put into controlled condition temperature 18 to 28 degree celcious in suitable light. After 8 day explants proliferated into mass of callus. This indicate the regeneration capacity of Brassica juncea species crop and variation of regeneration varies from hypocotyls to cotyledons. Which further use for genetic study , genetic transformation and crop improvement purpose.

Keywords: Regeneration, totipotancy, Embryogenesis.Sterilization.proliferation.MS medium. Transformation.

Abbreviations: MS media- murashige and Skoog (1962), NaCl-Sodium chloride, ABA-Abscisic acid, NAA-1-Napthalene acetic acid, PGR-Plant growth regulator, BAP-6-benzylaminopurine, LAR-Laminar air flow.

I. INTRODUCTION

Brassica juncea belongs to the family Brassicaceae (Cruciferae) is an amphidiploids species with the genome (AA=20) of B.compestris L. and genome (BB=16) of the B.nigra L. Koch. (AABB=36)¹. In India, the mustard is most important oilseed crop after groundnut accounting around 25% of total oil seed production. Indian mustard (Rai) cultivation has occupied about 85 to 90% of total area under cultivation of rapeseed and mustard ².Curently India account for about 12 to 13% of world oilseed area, 6 to 7 % of world oil meal production. In India rapeseed – mustard produced in area 21.1% and production 24.2% in among all the oilseed crops.³. The origin of B.juncea conflicting middle east seems to be the place of origin since the putative parents specie ,B.nigra and B.compestris would have been crossed (Olsson 1960, Mizushima and Tsunoda 1967).⁴. Rapeseed – mustard group of crop is grown in more than 70 countries globally on an area 31.68 mha with a production (mton 7.67, 2010) and productivity (kg/ha 1,179, 2010)⁵.

Oil content in brown sarson is 43%. Protein % in cake 38.52 to 42.2 % ,Ash % in cake 6.0 to 7.8 %. Iodine value 100.2 to 101.5 . Its oil generally rich in erucic acid (37.9 to 57%), it contan linolenic acid (4.7 to 13 %), the other important fatty acid present are linoleic acid (14%) and oleic acid (13%), Carbohydrate present in B. juncea are : total carbohydrate -(36.7%), free sugar (12.6%), Polysaccharide (2.2%), Composition of sugar as percentage are Stachyose 2.1%, Raffinose 1.2%, Melibiose 0.68%, Sucrose1.68%, Galactose 1.80%, Glucose 0.96%, fructose 2.10%, and other compound Pectic substances 6.5% and Cellulose 2.8%.⁶.

II. MATERIALS AND METHODS

A. Sterilization of Seed Surface

The seed material of B.juncea taken from the lab of proprietor (Allahabad Advance Agri Solution, Allahabad), 10gm. The first step to standardized our seed by surface sterilization. The seed must be surface sterilized to eliminate bacterial and fungal spore present on their surface. We achieved surface sterilization by washing them seed with distilled water properly for 20 min, and then with 1 to 2 % solution of sodium or calcium hypochloride for 10 min, then 0.1% of solution of mercuric chloride for-2 min, then the seed is rinsed several time with sterilized water to remove the disinfestant.⁷. This and the subsequent handling of explants or cultured cells and organ has to be done under aseptic condition, i.e in an environment free from bacteria and fungal spores through laminar air flow .⁸.



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429

Volume 8 Issue XII Dec 2020- Available at www.ijraset.com

B. M.S Media and Inoculation

For inoculation of seed in a MS media, we have already prepared MS media stock solution by following the composition of MS tissue culture medium (Murashige and skoog, 1962, Physiol plant 15:473-497). Take MS media in each sterilized cotton bed prepared bottle (bottle are sterilized by autoclaving at 121 degree celcious at 15 psi for 15 min. The pH is adjusted within a range of 5.6 to 5.8.⁹. Now inoculated the surface sterilized seed into MS medium under the aseptic condition and bottle closed by paraffin tape.

C. Callus Formation and Shoot Regeneration

The inoculated seed are maintained under a controlled environment, particularly in term of temperature and light. The temperature varies from 18 to 28 degree celcious. light is essential for this the culture room or the incubator should be kept as clean as possible it minimized contamination.



Fig. (A) Germinated seed under controlled condition (B) Sub culturing of seedling in form of cotyledon and hypocotyl under suitable medium.

D. Sub Culturing

After a period of time 8 day the germinated seed take out and transfer to fresh basal MS medium which contain the concentration of 2,4-D (1-5mg/l), and sucrose (0.5 - 3%). Only cutting hypocotyl and cotyledon are used for sub culturing. The embryogenic calli , obtained from hypocotyl and cotyledon , were refered as HEC (Hypocotyl-derived embryogenic calli) and CEC (cotyledon-derived embryogenic calli), respectively.⁷ In general callus culture are sub cultured every 4 to 6 week.

III. RESULT AND DISCUSSION

The Indian mustard B.juncea seed start germination after inoculation in the MS medium. The matured seedling with two leaf stage were arisen at 8 to 10 days after germinating of the seed . In this attempt the hypocotyls and cotyledon of the germinating seed used for the sub culturing. After the regeneration we get normal diploid 2n plantlet from hypocotyls and from cotyledon. The present study reveled that MS medium supplemented with 0.5 to 2.5 mg /litre of BAP along with 0.5 to 1.0 mg/l of NAA gives good response to maximum callus induction and shoot regeneration. Genetic modification can be done in the enzyme which involve in phytohormone production pathway, which can change the architecture of plant, as earlier genetic modification in gibbrelin oxidase has been done to change plant architecture . Finally it was concluded that the regeneration protocal develop in the present investigation is reliable and it would be effectively utilized for the genetic transformation of Brassica species using different genetic trait of crop improvement .¹⁰.



International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 8 Issue XII Dec 2020- Available at www.ijraset.com

IV. CONCLUSION

After performing this experiment it may be concluded that the regeneration protocal develop in the present investigation for Brassica juncea is reliable and can be effectively utilized for genetic transformation .

V. ACKNOWLEDGEMENT

Authors extent their thanks to Dr. Gaurav Krishna (Proprietor, Allahabad Advance Agri Solutions, Allahabad-211006) for providing seed, necessary facilities, guidance with their valuable suggestions, and continuous encouragement.

REFERENCES

- [1] Singh.A.K. (2008). Sustainable production of oil seed 15. 185.
- [2] Reddy.S.R., Reddy.Y .(2017). Agronomy of Field Crop. Rapeseed and mustard. 10.377-396.
- [3] Hegde.D.M. (2008).Sustainable production of oilseed. Rapeseed, Mustard technology. (editor Kumar.A., Chauhan.J.S., Chattopadhyay.C.).1.1-13.
- [4] Misra.A.K. (2008) .Sustainable production of oilseed, Taxonomy and botany of rapeseed-mustard .60-74.
- [5] Prasad .R.(2014).Textbook of field Crop Production (Commercial Crop 2nd). Kumar.A.,Rana.D.S.,Rapeseed –Mustard.3.80,81,82,-118.
- [6] Nagraj.G.(2009).Oilseed properties, processing, product and procedure.2, 29, 30, 32, 38,.
- [7] Akmal.M., Nafis.T., Mirja.K.J., Alam.p., Mohammad.A., Mujib.A., Abdin.M.Z.,(2011). High frequencies somatic embryogenesis in mustard crop (Brassica juncea L. ev. Pusa Jai kisan): Microscopic and histological analyses. Australian Journal of Crop Science. 5(13). 5,6.
- [8] Singh .B.D.(2015) Plant breeding principles and methods.Biotechology in crop improvements.(10th).35.660,662.
- [9] Dubey .S.K., Gupta.A.K., (2014). Callus Induction And Shoot Proliferation From Seedling Explants of Different Mustard Genotypes. International Journal of Current Microbiology and Applied Sciences.3(7). 2
- [10] Trivedi.N., Dubey.A.(2014). Efficient callus Regeneration and multiple shoot induction in brassica juncea var. Pusa Jaikisan.Research journal of recent science.3 (IVC -2014). 16-19.











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24*7 Support on Whatsapp)