



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 9 Issue: I Month of publication: January 2021

DOI: <https://doi.org/10.22214/ijraset.2021.32754>

www.ijraset.com

Call:  08813907089

E-mail ID: ijraset@gmail.com

A Comparative study of Physico-chemical Analysis of Ashwagandha Ghrita and Sapta Avarthita Ashwagandha Ghrita

Dr. Manjiri N Patil¹, Dr. Nitin B Patil²

¹BAMS, MD (Ayu), Assistant Professor, Dept of Panchakarma, Shri Acharya Deshbhushan Ayurvedic Medical College, Bedkihal, Belgaum.

²Associate Professor, Dept of Agada Tantra, Mai Bhago ayurvedic medical college, Muktsar, India

Abstract: Ghrita is one of the Ayurvedic drugs that contain ghee as the base to dissolve or extract or hold the active therapeutic principles from the ingredients. Ashwagandha is the herb used for rejuvenation of whole body and shows immunomodulatory, adaptogenic and several other activities.

Here in this study Ashwagandha ghrita and Sapata avarthita Ashwagandha ghrita were prepared and its physico-chemical analysis was done. By repeated processing of the same ghrita the final product gets potentified and useful for therapeutic purposes. The physico chemical properties reveals that Saponification value and Acid value were slightly reduced from ashwagandha ghrita to Sapta avarthana process which indicates an increase in free fatty acid content in Sapta avarthana. Decrease in iodine value indicates the degree of unsaturation was reduced to a minimal level and conversion of unsaturated fatty acids to saturated fatty acids occurred. There is no significant change in Refractive index. Specific gravity of Sapta avarthita ghrita was reduced which indicates the increase of density. To standardize the preparation TLC was carried out and details of sapta avrthita Ashwagandha ghrita preparation and its physico-chemical analysis is been discussed in this study.

Keywords: Immunomodulatory, Avarthana, Saponification Value, Acid value, Iodine value, TLC

I. INTRODUCTION

Ghrita is one of the Ayurvedic drugs that contain ghee as the base to dissolve or extract or hold the active therapeutic principles from the ingredients. Ghritas are medicated ghee preparations containing the fat-soluble components of the ingredients used in these preparations. The principle of preparation is the protracted boiling of ghee with prescribed *kashayas* (decoctions) and *kalkas* (a fine paste of the drug/drugs) to dehydration or near dehydration thereby effecting the transference of the fat-soluble principles to the ghrita, from the drug ingredients or *kashayas* or *swarasas* as the case may be according to the formulation.¹

Ghrita's lipophilic property facilitates drug delivery to mitochondrion microsome and nuclear membrane of the cell and are also used as *rasayana*. Ashwagandha is *brumhaniya*, *balya*, its active component is alkaloids and withanoloids which has medicinal characters and helps in maintain general health and wellness.²

Various studies carried out on different formulations of ashwagandha such as ashwagandha granules, ashwagandha arishta and its standardization. Here in this study Ashwagandha ghrita and Sapta avarthita Ashwagandha ghrita were prepared and its physico-chemical study was carried out. Ashwagandha ghrita is best *rasayana* and useful in *vata vyadhis* and in *sapta avarthana* it gets more potentified. Ashwagandha is one of the reputed herbs in Ayurveda and has many actions on body like anti-ageing, adaptogenic, immunomodulatory, anxiety, depression, stress, cardiovascular protection, hypothyroidism to name a few.³ It contains alkaloids and steroidal lactones, many bio-chemical heterogeneous alkaloids, including choline, tropanol, pseudotropanol, cuscoygrene, 3-tigloyloxytropana, isopelletierine and several other steroidal lactones, withanolides and several sitosterols.^{4,5}

II. METHOD OF PREPARATION

In preparation of Ashwagandha ghrita and Sapta avarthita Ashwagandha ghrita, Ashwagandha, Triphala, Musta, Haridra, Goghrita, Godugdha were used. These formulations prepared in KLEU's GMP certified Pharmacy, Belgaum.

Ashwagandha ghrita was prepared by *snehapaka vidhi*. Murchana of ghrita was done before preparing Ashwagandha ghrita to remove impurities of ghrita and to make it fragrance and colour full.⁶

Preparation of Murchita Ghrita⁷:

In the ratio of 1:4:16, *kalka*, *Ghrita* and *drava dravyas* are taken and subjected to *snehapaka* till the appearance of *sneha siddhi lakshanas*. *Kalka dravyas* added are *Tripahala*, *Haridra*, *Musta* and *Nimbu swarasa*. The quantity of ingredients are *Kalka*- 3.150kg in coarse form, *ghrita*- 12 kg, *Drava dravya*- water 44 lit.

These ingredients were taken into a big sufficient vessel and subjected to medium flame. Continuous stirring was done to prevent the sticking of *kalka* to the bottom of the vessel.

Preparation of *Ashwagandha ghrita*⁸:

The ingredients are *Ashwagandha kalka* 2.570 kg, *Murchita ghrita* 10.300kg, *Ashwagandha Kwatha* 40lit and *ksheera*(milk) 11 lit.

The above ingredients were taken into a big vessel (15 kg) and subjected to mild flame and continuous stirring was done till getting *sneha siddhi lakshanas*. After the evaporating of water only ghee and *kalka* remained, which became like brown mud paste. This was the stage to observe intensely, after this slowly ghee separates from the *kalka* and good quality smell generates. Once the *kalka* becomes bolus form, it should be taken into the hand and rolled in between the fingers to make *varti* (roll) and subjected to fire (candle light) to assess *madhyama sneha siddhi lakshana*.

After *samyak paka*, the ghee was taken off from the fire and filtered when it is in liquid form. The final product was measured and subjected for further process.

Preparation of *Sapta avarthita Ashwagandha ghrita*

The ingredients are *Ashwagandha kalka* 1.300 gm, 6 *avarthita ashwagandha ghrita* 6.500gm, *Ashwagandha Kwatha* 24lit and milk 6.5 lit.

The *sneha paka vidhi* mentioned earlier in *Ashwagandha ghrita* was carried out during *sapta avarhtana*.

A. Analytical Study

Authentication of all ingredients and Physico-chemical analysis, TLC of *Ashwagandha ghrita* and *Sapta avarthita Ashwagandha ghrita* was done at AYUSH approved drug testing laboratory, KLEU's Shri BMK Ayurveda Mahavidyalaya, Belgaum.

The following analysis were done by adopting standard protocols.⁹

B. Refractive Index

The refractive index is measured at 25°C (±0.5) with reference to the wavelength of the D line of sodium (λ 589.3 nm). The temperature should be carefully adjusted and maintained since the refractive index varies significantly with temperature.

The Abbe's refractometer is convenient for most measurements of refractive index. To achieve accuracy, the apparatus should be calibrated against distilled water which has a refractive index of 1.3325 at 25°C. After every step part was cleaned with Diethyl ether, which was evaporated by own. Thin layered of *Ashwagandha ghrita* and *Sapta avarthita Ashwagandha ghrita* samples were applied by a cotton swab.

C. Specific Gravity

Empty Specific Gravity bottle was weighed and the bottle filled with distilled water and again weighed, the same bottle was then filled with 1% of sample and weighed. All these 3 weights were noted at 25°C temperature, the Specific Gravity of sample was calculated.

D. Determination of Saponification Value

2g of *ghrita* taken into a 250 ml RB flask fitted with a reflux condenser. Added 25 ml of 0.5 M Alcoholic potash and reflux on a water bath for 30 minutes. Cool, add 1 ml of Phenolphthalein solution and titrate immediately with 0.5 Hydrochloric acid. Repeat the operation omitting the substance being examined. The values of samples were calculated.

E. Determination of Acid Value

10 g of *ghrita* sample taken in a conical flask and added 50 ml of acid free alcohol ether mixture (25 +25 ml) previously neutralized with the 0.1 M potassium hydroxide solution. Shaked well. Added one ml of Phenolphthalein solution and titrate against 0.1 M Potassium hydroxide solution. End point is the appearance of pale pink colour. The values were calculated.

F. Determination of Moisture Content (Loss on Drying)

5.140 g of *ghrita* in a previously weighed petri dish was taken and kept in an oven at 105°C for 5 hours. Cooled in desiccators and weighed. the percentage of loss in weight of the sample was calculated.

G. Determination of Iodine Value

Ghrita 2gm taken in dry 500 ml iodine flask, added 10 ml of carbon tetrachloride and dissolved. Added 20 ml of iodine monochloride solution, inserted the stopper and allowed to stand in the dark at a temperature between 15°C and 25°C for 30 minutes. Placed 15 ml of potassium iodide solution in the cup top, carefully removed the stopper, rinsed the stopper and the sides of the flask with 100 ml of water, shake and titrate with 0.1 M sodium thiosulphate using starch solution, added towards the end of the titration, as indicator. Noted the number of ml required. Repeated the procedure omitting the substance being examined and noted the number of ml required.

H. Qualitative Analysis by Thin Layer Chromatography⁹

The T.L.C is the important and simple analytical tools for the qualitative analysis of the raw materials.

I. Thin Layer Chromatography

T.L.C is one of the most widely used techniques for rapid identification of drugs and its formulations. It is equally applicable to the drugs as raw material state and pure state.

Chromatographic conditions:

The Alcoholic extract of *Ashwagandha ghrita* and *Sapta avarthita Ashwagandha ghrita* was subjected for thin layer chromatography as follows,

- 1) *Preparation of TLC*: Pre coated Silica Plate was used.
- 2) *Sample Preparation*: The extracts obtained after Alcoholic extraction were used for TLC. Plate/Stationary Phase-Silica gel G. Solvent front run up to 8 cm. Applicator – Capillary tube. Solvent/Mobile phase – Toluene: Ethyl acetate (7: 3) through trial-and-error method. This solvent system holds good and clear appearances of bands (spots) were seen. Detection was done by keeping plate in Iodine vapour chamber.

III. RESULTS

Table No 1: Showing the results of The Physico-chemical analysis of *Ashwagandha ghrita* and *Sapta avarthita Ashwagandha ghrita*

Samples	Refractive Index	Specific Gravity	Saponification Value	Acid Value	LOD	Iodine Value
Ashwagandha ghrita	1.46	1.0904	218.02	1.600	0.460	26.80
Sapta avarthita Ashwagandha ghrita	1.45	0.901	217.62	1.455	0.408	25.82

Table No 2: Showing the Rf value of *Ashwagandha ghrita* and *Sapta avarthita Ashwagandha ghrita*

Sample	Extract	Solvent system	Spots at 254nm	Spots at 366nm
Ashwagandha ghrita	Methanol Extract	Toulene: Ethyl Acetate (7:3)	0.07, 0.12, 0.42	0.05, 0.11, 0.15
Sapta avarthita Ashwagandha ghrita			0.08, 0.13, 0.52	0.05, 0.11, 0.15

IV. DISCUSSION

Ghrita is one among best *Ajasrika rasayana*, it is *Ayurvedhaka balavardhaka ojavardhaka*, *vayasthapaka dhatuposhaka* and best among the *sneha dravyas*.¹⁰

Ashwagandha is an Ayurvedic herb and many studies have been done on its therapeutic potential and is very reputed drug in immunomodulation, ant-ageing and tonic for the body.

In the present study *Ashwagandha ghrita* was prepared by traditional method and was processed seven times to get *Sapta avarthita ashwagandha ghrita*. The Physico-chemical analysis was performed and the values are discussed here.

Refractive index was not changed significantly from *Ashwagandha ghrita* to *sapta avarthana*, which indicates there is no change in length of fatty acids⁹. Specific gravity indicates the solute content in solvent. It was decreased from *ashwagandha ghrita* to *sapta avarthana* process, may be due to the fortification by *Ashwagandha*, it may have less specific gravity and due to continuous boiling the guru guna might become laghu in this parameter. Saponification value was not changed significantly from *ashwagandha ghrita* to *sapta avarthana* process it indicates that an increase in free fatty acid content in *sapta avrthana*.

When compared to *Ashwagandha ghrita* the acid value was slightly decreased in *Sapta avarthita Ashwagandha ghrita*. It is due to the drug effect which is added to the ghrita, due to the breaking in fatty acid chains acid value may decrease. Moisture content/Loss on drying was gradually reduced in *sapta avarthana* process. It indicates that the water content was present in *ashwagandha ghrita*, when it is processed further water content was evaporated completely.

Iodine value was slightly decreased from *ashwagandha ghrita* and *sapta avarthita ghrita*. Decrease in iodine value indicates that degree of unsaturation was reduced to a minimal level and conversion of unsaturated fatty acids to saturated fatty acids occurred¹¹. The above quality parameters/standardization parameters of *Sapta avarthita Ashwagandha ghrita* were shown minimal changes, in which average values can be taken as standards. TLC of both the samples were developed and Rf values at 254nm UV rays were noted at 0.07, 0.12, 0.42 in *Ashwagandha ghrita*, 0.08, 0.13, 0.52 in *Sapta avarthita Ashwagandha ghrita*, and Rf values at 366nm UV rays were noted at 0.05, 0.11, 0.15 in *Ashwagandha ghrita*, 0.05, 0.11, 0.15 in *Sapta avarthita Ashwagandha ghrita* (Table No.2) (Fig No.1) However, this formulation should be standardized by HPTLC, HPLC and pharmacokinetic profiling methods by using markers. These studies are suggested for future.


V. ACKNOWLEDGEMENT

The author would like to thank Dr. B.S. Prasad sir, Principal, KLEU'S Shri BMK Ayurved College of Belgaum, for providing the facilities for Analysis of the *ghrita*. The author would like to acknowledge helpful discussions on the manuscript with Dr.Srinath Vaidya, Dr.Ajit Lingayat, Dr.R.V.Sanapeti, Belgaum and Dr. Nitin Patil for helping to get TLC of samples.

REFERENCES

- [1] Dholakia M V, Vachharajani Y R, Shah D C, Bahadur M V and Subrata De J R A S 1992 13(3-4) 173
- [2] Rasayana sarvaswam-Dr.Santosh Belawadi, 1st edition, Nov 2011.
- [3] Unknown., *Withania somnifera* Monograph, *Alternative Medicine Review.*, 2004,9,210- 213
- [4] Farzana Akhtar, Isolation And Structural Studies On The Chemical Constituents Of *Withania Somnifera*, *Fumaria Flabellata* And X-Ray Diffraction Studies, *Pakistan research repository.*, 1997,275.
- [5] Mohammad H.M, Elisabeth M., Steroidal Lactones from *Withania somnifera*, an Ancient Plant for Novel Medicine, *Molecules.*, 2009, 14, 2373-2393.
- [6] Kinnari Dhruve and Anand K.Chaudhary, *Snehakalpana-A probable Pharmaceutical Explanation*, *Aryavaidyan*, Vol.XX., No.3, Feb-Apr.2007.p.181-189.
- [7] Govindadas Sen, Bhaishjya Ratnavali, Edited by prof.Siddhi Nandan Mishra, *Jwara chikitsa* 1266, Varanasi:Chaukhamba SurabharatiPrakashan, Reprint.p.206.
- [8] Chakrapanidatta, Chakradatta, Edited by Indradev Tripathi, Ramanath Dwivedy, *Vatavyadhi chikitsa* 90. Varanasi: Chaukhamba Sanskrit Bhawan, 2011.p.141.
- [9] Quality control manual for ASUdrugs, Govt. of India, Dept of Ayush, Ministry of family welfare, Pharmacopeia laboratory, for Indian Medicine, GhaZiabad, 2008.
- [10] Sushruta. *Sushuta samhita Dalhana*, Ed.By Yadavji T.Chikitsa sthana31/3. Varanasi: Chaukhamba Surabharati prakashan; 2010.p.509.
- [11] Dhurvey Y.R., Kawatikwar P.S, Sarkarkar P.M, Evaluation of Physico chemical properties of cow ghee before and after Hydrogenation, *IJCRGG*, Jan-Mar 2012.

Fig.No.1. Showing the TLC of both formulations.



A-Ashwagandha ghrita and Sapta avarthita Ashwagandha ghrita TLC at 254nm
B- Ashwagandha ghrita and Sapta avarthita Ashwagandha ghrita TLC at 366nm



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)