



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 9 Issue: II Month of publication: February 2021 DOI: https://doi.org/10.22214/ijraset.2021.32987

www.ijraset.com

Call: 🕥 08813907089 🔰 E-mail ID: ijraset@gmail.com



Total Lipid Content of Ascardia Galli from its Host Gallus Domesticus

R. M. Khadap

Department of Zoology, Nutan College Sailu. Dist. Parbhani-431503 M.S. India.

Abstract: Lipid are important constituents for an animal body as they serve both in structural makeup and in energy yielding, but in parasites there is a good deal difference in the mode of procurement of these impartment metabolites. They may be available to them in digested or semi digested state. In helminthes the fatty acid synthesis is very limited as per the studies of Ginger and Fairbairn, 1966. In nematode most of lipids located in the hypodermis especially in the lateral cords, muscle cells, intestinal cells and ovaries. Lipid is the chief food reserve in free living nematode and plant parasitic nematodes. Keywords: Ascardia galli, Gallus domesticus, total lipid content.

I. INTRODUCTION

In helminth worms are the biological importance of lipids as group of certain related hydrocarbons, various attempts have been made to have a data on total lipid content of several helminth parasites. The total lipid content of *Ascaris lumbriciies* have been worked out in fresh tissue in both male and female worms by Cavier et.al,1958. The lipid content has been worked out in dry tissue of *Nippostrongylus brasiliensis* by Roberts and Fairbaim,1965. The lipid content in *Ascaridia galli* has been worked out in fresh tissue of male and female worms by Shorb and Shorb,1966, hence the author in his present study made an attempt to estimate the total lipid content in male and female worms of *Ascaridia galli*.

II. MATERIAL AND METHOD

The nematodes were collected from intestine of *Gallus domesticus*. Required quantity of these sample dried at $48-50^{\circ}$ c for 48 hrs separately and preserved powdered samples weighted for quantitative estimation of lipid,% between male and female by methanol, chloroform estimation method (Barner H. and Block Stok, 1973).

III. RESULTS

The results obtained on the total lipid content male and female worms is 12.35 and 14.80 respectively. The male to female ratio of lipid content is 0.83 indicating that its content is more in female by 16.55%.

SEX	CONTENT	Male to female ratio	Percentage difference
MALE	12.35		
FEMALE	14.80	0.834	16.55

TABLE:1 Total Lipid content in (mg%) in male and female worms of Ascaridia galli



Total lipid content in (mg%) in male and female worms of Ascaridia galli



International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 9 Issue II Feb 2021- Available at www.ijraset.com

IV. DISSCUSSION

The result obtained on the total lipids of the *Ascaridia galli* has necessary prerequisites conductive to its parasites to its parasite mode of life. The value of total lipid (wet.weight) are comparable in trend with that of *A.lumbricodies*,male1.3% and female1.6% as reported by Cavier et.al.,1958 and in dry tissue they agree in trend with those of *A.lumbricoides*,male6.7% and female11.9%,Roberts and Fairbairn,1965.In *Ascaridia galli* and *Ascaridia lumbricoides* female has higher lipid content which is justifiable as per the structural needs for their large number of eggs. There is a variation in the individual valve of the parasites mentioned above which is due to species difference, different host, their metabolism and habit of the parasites. The results of the lipid contents are significantly more on female worm. The difference between the two sexes. It can be inferred that the female worms are metabolically more active.

V. ACKNOWLEDGEMENT

The author are thankful to the Principal Nutan Mahavidyalaya sailu, for providing laboratory facilities.

REFERENCES

- [1] Cavier R., SavelJ. and Monteoliva M.1958. Nature at repartition selonle sexe des substances lipidques chez. A. lumbricoides Linne, 1758. Bull. Soc. Chim. Biol. pp 177-187.
- [2] Ginger C.D. and Fairbairn D.1966.Lipid Metabolism in helminth parasites I the lipid of H.diminuta.J.Parasitol(52)pp 1086-1096.
- [3] Roberts L.S. and Fairbaim D,1965. Metabolic studies on adut Nippostronggybus brasiliensis. J.Parasitol(51).pp 129-138.
- [4] ShorbD.A. and ShorbM.S.1966. Phospholipids of some worm parasites proc.Inter. Congr. Parasitol Vol.(2)pp 59-61.
- [5] Yamaguti S,1961.Systema Helminthum Vol.III.The nematode of Vertebrate Part I &II .Interscience Publishers ltd.London.











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24*7 Support on Whatsapp)