



# **iJRASET**

International Journal For Research in  
Applied Science and Engineering Technology



---

# **INTERNATIONAL JOURNAL FOR RESEARCH**

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume: 3**

**Issue: XII**

**Month of publication: December 2015**

**DOI:**

**[www.ijraset.com](http://www.ijraset.com)**

**Call:  08813907089**

**E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)**

# **Assessment of Groundwater potential zone using remote sensing and GIS, A case study in Nandhiyar sub basin, Tamilnadu, India**

Ramamoorthy.P<sup>1</sup>, Raju.S<sup>2</sup>, Senthilraj.G<sup>2</sup>

<sup>1</sup>University of Madras, Chennai, Tamilnadu, India.

<sup>2</sup>Mailam Engineering College, Mailam, Tamilnadu, India.

**Abstract:** Groundwater is one of the greatest hidden natural resources of the earth's crust. It has ubiquitous occurrence. The deformative forces and different earth processes depending upon the nature and composition of the rocks produces characteristic structural, morphological features, which themselves are responsible for the groundwater accumulation, storage and movement, a detailed understanding of all these pathfinders and appreciation of their relative significance in different hydro geological environment is essential in proper planning and management of the groundwater resources. Assessing the groundwater potential zone is particularly important for the safety of water quality and the management of groundwater systems. A case study was conducted in Nandhiyar sub basin to find out the groundwater potential zone. The thematic maps such as geomorphology, geology, soil, drainage, lineament map were prepared in Arc GIS. All the thematic layers were integrated in to Arc GIS 9.3 software to find a groundwater potential zone map of the study area. Thus, three different groundwater potential zones were identified, namely 'Poor', 'Moderate' and 'Good'.

**Key words:** Groundwater, Nandhiyar, Geomorphology, Soil, Lineament etc

## **I. INTRODUCTION**

Water is one of the most essential commodities for mankind and the largest available source of fresh water lays underground (Todd,1980). It is one of the most significant natural resources which support both human needs and economic development. Tremendous increase in the agricultural, industrial and domestic activities in recent years has increased the demand for good quality water to meet the growing needs. Groundwater is mostly preferred to meet this growing demand because of its lower level of contamination and wider distribution (Biswas Arkoprovo et.al.,2012).

Groundwater resources are an important natural resource for its use in domestic, agriculture, and industries purposes. There has been a tremendous increase in the demand for groundwater due to increase in population, advanced irrigation practices and industrial usages (Ramamoorthy and Rammohan,2015).

The occurrence of groundwater at any place on the earth is not a matter of chance but a consequence of the interaction of the climatic, geological, hydrological, physiographical and ecological factors. Groundwater exploration operation is essentially a hydrogeological and geophysical inference operation and is dependent on the correct interpretation of the hydrological indicators and evidences (Antony Ravindran,2012).

Several researches have utilized the GIS technology and the remotely sensed derived data for water resources management, groundwater assessment and modeling. This study addresses the strategies for an integrated approach of remote sensing and GIS techniques to delineate groundwater prospective zones. The technique of integrated remote sensing and GIS has proved to be an efficient tool in groundwater studies (Edet et al. 1998; Krishnamurthy et.al 1996 and Murthy 2000). The Remote Sensing and GIS tools have opened new paths in water resources studies. One of the greatest advantages of using remote sensing and GIS techniques for hydrological investigations and monitoring is its ability to generate information in spatial and temporal domain, which is very crucial for successful analysis, prediction and validation (Sarma and Saraf, 2002).

## **II. STUDY AREA**

The study area Nandhiyar Sub basin is located on the western part of Chennai basin group and it spread over parts of Vellore and



## International Journal for Research in Applied Science & Engineering Technology (IJRASET)




Fig:2 Geology Map




Fig:3 Geomorphology Map

2) *Geomorphology*: Geomorphic features play a vital role in the evaluation of surface and ground water resources. The hydro geomorphic units such as flood plain, valley fill, buried pediment is good sources of groundwater where as structural hills, pediment zone and gullied land are poor recharge zones (Subagunasekar et al.2012).

The following geomorphological features are identified from the study area, Pediment, Buried Pediment (Shallow), Buried Pediment( Medium), Buried Pediment(deep),Structural hill, dyke etc.

The north west portion of the sub basin covered by vast pediment zone.Some portion in the northeast and northwest part of study area covered by Buried pediment (shallow), Buried Pediment (Moderate) is found in the upper central portion of the sub basin(Fig:3).

3) *Soil*: Soil map (Fig:4) of Nandhiyar sub basin was prepared based on soil classification National Bureau of Soil Survey and Land use Planning, Bangalore (NBSS) in co-operation with the Department of Agriculture, Tamil Nadu. The predominant soil types found in this sub basin is Alfisols, and Inceptisols. Due to different stages of weathering of parent material, the above soil types are met with in mixture.

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)




Fig: 4 Soil Map




Fig:5 Lineament Map

- 4) **Lineament:** Lineaments are surface manifestation of the linear fractures like faults, joints and fractures, groundwater potential of high order is indicated where lineament run along and across the weathering rock deposit are intersect on another (Radhakrishnan and Ramamoorthy, 2014). Lineament features indirectly reveals the groundwater potential zones, areas with high lineament density are good for groundwater potential zones (Haridas et al., 1998). The lineament intersection areas are considered to be good groundwater potential zones (Pothiraj and Rajagopalan, 2012). The lineament trending North east and South west are identified in the study area indicating regions of good groundwater source (Fig:5).
- 5) **Drainage:** Nandhiyar is the tributary of Kosasthalaiyar river which runs towards east in the central part of the sub basin. The river Nandhiyar originates from Puttur hill at an altitude of 582 m near Singasamudram in Andhra Pradesh. Drainage map (Fig-6) of the study area shows in predominantly dendritic pattern.

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)




Fig. 6. Drainage Map




Fig. 7. Groundwater Potential Map

6) *Groundwater potential zones*: By integrating the various thematic maps such as geology, geomorphology, soil, lineament, and drainage, with Arc GIS 9.3 software using weightage index overlay analysis and the groundwater potential zone was classified as poor, moderate and good (Fig.7) and assigning the weighted values and there rank was shown in table:1.

TABLE :1 - RANKS AND WEIGHTAGES FOR VARIOUS PARAMETERS FOR GROUNDWATER POTENTIAL ZONE.

Criteria	Features	Weightage	Rank
Geology	Sand stone /shale	3	High
	Quartz gravel	3	
	Talc schist	2	Medium
	Amphibolite/Pyroxenite	1	Low
	Epidote,,Hornblende Gneiss	1	
	Granite/Charnockite, Dolerite/ Gabbro	1	
Geomorphology	Buried pediment(Deep)	3	High
	Buried pediment(Moderate)	2	Medium
	Buried Pediment (shallow)	2	
	Pediment	1	Low
	Structural hill	1	
Lineament	Present	5	High
	Absent	1	Low

### III. CONCLUSION

GIS plays a more important role in the evaluation of groundwater potential. It is a prime component for the integration and analysis of the spatial and non-spatial information. By integrating all the thematic layers, the study area classified in to (i) Poor (ii) Moderate and (iii) Good groundwater potential zone. The study reveals that combination of Buried pediment (deep), Lineament, lineament intersection and drainage shows good source of groundwater. From this study observed that remote sensing and GIS tool can utilize successfully to categorize the groundwater potential zones. This groundwater potential zone map will be useful for effective

## International Journal for Research in Applied Science & Engineering Technology (IJRASET)

identification of apt locations for withdrawal of groundwater. It can be used for future development and management the groundwater source in the study area.

### REFERENCES

- [1] Antony Ravindran A., (2012). Azimuthal Square Array Resistivity Method and Groundwater Exploration in Sanganoor, Coimbatore District, Tamilnadu, India, Res. J. recent sci, 1(4), 41–45.
- [2] Biswas Arkoprovo, Jana Adarsa and Sharma Shashi Prakash(2012) Delineation of Groundwater Potential Zones using Satellite Remote Sensing and Geographic Information System Techniques: A Case study from Ganjam district, Orissa, India, Research Journal of Recent Sciences, Vol. 1(9), 59-66.
- [3] Edet, A.E., Okereke, C.S., Teme, S.C. and Esu, E.O.,(1998) "Application of Remote Sensing Data Groundwater Exploration: A Case Study of the Cross River State, Southeastern Nigeria," Hydrogeology, vol. 6, pp. 394–404,.
- [4] Haridas, V.R., Aravindan,S., Girish,G.,(1998).Remote sensing and its applications for groundwater favourable area identification. Quarterly journal of GARC 6, 18-22.
- [5] Krishnamurthy, J., Kumar, N.V., Jayraman, V. and Manivel, M.,(1996) "An Approach to Demarcate Groundwater Potential Zones through Remote Sensing and GIS," International Journal of Remote Sensing, vol. 17, no. 10, pp. 1867–1884,.
- [6] Murthy K.S.R.,(2000) "Groundwater Potential in a Semi-Arid Region of Andhra Pradesh: A Geographical Information System Approach," International Journal of Remote Sensing, vol. 21, no. 9, pp. 1867-1884,.
- [7] Prabu Pothiraj & Baskaran Rajagopalan ,(2012),A GIS and remote sensing based evaluation of groundwater potential zones in a hard rock terrain of Vaigai sub-basin, India. Arab J Geosci. DOI 10.1007/s12517-011-0512-3
- [8] Radhakrishnan.D., and Ramamoorthy.P,(2014) Delineation of Groundwater Recharge Potential zones in Mailam Block, Villupuram district, Using GIS,"International Journal of Water Research 2014; 2(2): pp.71-75.
- [9] Ramamoorthy.P and Rammohan.V(2015) Assessment of Groundwater potential zone using remote sensing and GIS in Varahanadhi watershed, Tamilnadu, India, International Journal for Research in Applied Science & Engineering Technology (IJRASET), Volume 3 Issue V,p-695- 702.
- [10] Sarma, B. and Saraf, A.K.,(2002) Study of Land Use-Groundwater Relationship Using an Integrated Remote Sensing and GIS Approach, <http://www.gisdevelopment.net>,.
- [11] Subagunasekar,M.,Sashikkumar,M.C.,(2012).INT J CURR SCI ,159-162.
- [12] Todd D.K.,(1980) Groundwater Hydrology, Second Edition, John Wiley and Sons Inc., New York, 556



10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)