



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 9 Issue: VII Month of publication: July 2021

DOI: https://doi.org/10.22214/ijraset.2021.36978

www.ijraset.com

Call: 🕥 08813907089 🔰 E-mail ID: ijraset@gmail.com



Sunanda Ravindranath<sup>1</sup>, Nirdosh Patil<sup>2</sup>, Shweta Patil<sup>3</sup>, Ambika Bhusange<sup>4</sup>

<sup>1, 2, 3</sup>Department of Chemistry, Appa Institute of Engineering and Technology, Vidya Nagar, Kalabuaragi-585103, Karnataka, India <sup>4</sup>Department of Chemistry, Godutai Engineering College for women, Vidya Nagar, Kalabuaragi-585103, Karnataka, India

Abstract: A few complexes 2-{[4-(5-bromo thiophen-2-yl)-6-phenyl pyrimidin-2-yl] carbonoimidoyl} phenol [HL<sup>5</sup>] and 2-{[4-(5-bromo thiophen-2-yl) -6- phenyl pyrimidin-2-yl]carbon imidoyl}-5-methoxyphenol [HL<sup>6</sup>] with first row transition metal ions have been reported in this paper. The stichiometry of type (1:1) [M(L)]2H<sub>2</sub>O is analysed by these complexes. Complexes with ligands (HL<sup>5</sup> and HL<sup>6</sup>) of metal ions Co(II), Ni(II), Cu(II) and Zn(II) reveal co- ordination number six, by means of magnetic susceptibility, electronic, IR, 1H NMR, X-ray diffraction and ESR spectrum data. Copper(II) complexes with (HL<sup>5</sup>) and (HL<sup>6</sup>) were studied spectrophotometrically at different temperatures having an absorption maxima of 535 nm. The findings demonstrate that copper(II) and (HL<sup>5</sup>) and (HL<sup>6</sup>) are combined at pH 5.0 at a molar ratio of 1:1. The stability constants of the complexes were calculated to be  $13.06 \times 10^4$ ,  $14.51 \times 10^4$  and  $10.40 \times 10^4$  and  $13.18 \times 10^4$ ,  $15.06 \times 10^4$ ,  $0.55 \times 10^4$ ,  $0.79 \times 10^4$  for (HL<sup>5</sup>) and (HL<sup>6</sup>) respectively by Jobs continuous variation method at 30, 35 and  $40^{\circ}$ C, respectively. Both ligands and their metal complexes were screened for their antibacterial and antifungal activity by cup plate method and antioxidant activity by using standard DPPH radical scavenging ability method.

Keywords: Antifungal, complexes, antioxidant, antibacterial, magnetic,

#### I. INTRODUCTION

In many physiologically active natural compounds nitrogen heterocycles, in general, and pyrimidine in particular, show substantial therapeutic promise. Pyrimidine is one of the most famous nucleic acid chemistry structures. Many chemotherapeutic drugs in the pyrimidine nucleus are now in clinical use, including antifungal (flucytocine), antifungal (pyrimethamine), antifungal (iodoxuridine, trifluridine and zidovudine), antiviral (sulfadiazine, sulphamarazine and sulfamethazine), antimalarial (pyrimidamine) antifungal agents (flucytocine). The large spectrum of biological and pharmacological activities of pyrimidines, including anticarcinogenicity[1-3], antibacterial[4], anti-inflammatory[5], antiviral[6], anti HIV [7], anti-hypertensive[8-9], anticonvulsant and analgesic[10], is of particular concern. In recent years, an increase in drug-resistant bacterial strain isolates has created a therapeutic hurdle to physicians when choosing antimicrobials. Consequently, it is critically necessary to produce novel antibacterial medicines with strong antibacterial action and reduced adverse effects. 2-acetyl-5-bromothiophene produced equivalent chalcones when condensed with different aromatic aldehydes in the alcoholic KOH solution. The chalcones were then reacted in the presence of a base with ethanol guanidine hydrochloride that yielded pyrimidine derivatives. The freshly synthesized heterocycles were characterized by the NMR, IR, and mass spectroscopic data. The antibacterial characteristics of all freshly synthesized compounds were examined[11]. We have reported in the current research that certain novel amino pyrimidines, such as  $2-\{[4-(5-bromothiophen-2-yl)-6-phenylpyrimidin-2-yl]carbonoimidoyl\} phenol [HL<sup>1</sup>] and <math>2-\{[4-(5-bromothiophen-2-yl)-6-phenylpyrimidin-2-yl]carbonoimidoyl} phenol [HL<sup>2</sup>] and their metal complexes.$ 



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429

Volume 9 Issue VII July 2021- Available at www.ijraset.com

### II. MATERIALS AND METHODS

All the chemicals and solvents used were of AR grade and were obtained from Sisco-Chem Industries. The metal salts were purchased from commercial sources. Metal contents were estimated using standard methods[12]. The electronic spectra were recorded on an Elico SL 159 spectrophotometer in the 200-1000nm ranges in DMF solution (10-3). Elemental analyses were obtained from HERAEUS C, H, N-O rapid analyzer. ESR measurements were carried out on a VARAN E-109 GHz. The experiment was carried by taking DPPH as the reference with the field set at 3200 Gauss. Magnetic susceptibilities were determined by the Faraday method using a model 300 Lewis Coil Force Magnetometer of one Tesla field strength at room temperature. The instrument was calibrated with Hg [Co (SCN)<sub>4</sub>][13]. All melting points were uncorrected and measured using an Electro-thermal IA 9100 apparatus (Shimadzu, Japan). IR spectra were recorded as potassium bromide pellets on a Perkin-Elmer 1650 spectrophotometer (Shimadzu). A Horiba F.8 pH meter, calibrated with standard buffer solutions, was used for pH measurements. Copper(II) sulphate pentahydrate was purchased from Merck, Germany. All other chemicals used were of analytical grade purchased from Merck, Germany. Double-distilled water was used throughout this study[14]. Both ligands and their metal complexes were screened for their antibacterial and antifungal activity by cup plate method and antioxidant activity by using standard DPPH radical scavenging ability method.



Fig. 1. Structure of ligands,  $HL^1 = R = H$  and  $HL^2 = R = OCH_3$ 

### A. Preparation of Ligands

- Preparation of 1-(5-bromothiophen-2-yl)-3-phenylprop-2-en-1-one: The substituted chalcones were prepared by reacting equimolar quantity of aromatic aldehydes (0.001mol) and 2-acetyl-5-bromothiophene iniethanol. 40% KOH was added to the reaction mixture at room temperature, drop by drop with regular stirring. After the completion of reaction, the precipitate thus obtained was separated by filtration, washed with cold ethanol, dried and purified by recrystallization from ethanol.
- 2) Preparation of 4-(5-bromothiophene-2yl)-6-pheneylpyrimidine-2- Amine: Refluxing a combination of alternative chalcones (0.001mol), guanidine hydrochloride (0.001mol) and alkali (KOH), into 20ml ethanol, replaced pyrimidine. The reaction mixture was dumped into ice cold water when the reaction was finished. The resulting precipitation was filtered, washed with ethanol, dried and purified using ethanol recrystallation.
- 3) Preparation of 2-{[4-(5-bromo thiophene-2-yl)-6-phenyl pyrimidin-2- yl] carbonimidoyl} phenol [HL<sup>1</sup>] and 2-{[4-(5-bromothiophene-2-yl)-6- phenylpyrimidin-2- yl] carbon imido yl}-5-methoxyphenol [HL<sup>2</sup>]: Equimolar 4-(5-bromothiophene-2yl)-6-pheneylpyrimidine-2-amine (0.001 mol) solutions and replaced salicylaldehyde (0.001mol) with 4 hour reaction mixture in 40 ml of ethanol reflux, with a 40% addition of KOH to the reaction mixture. Afterwards, for the reaction mixture, the precipitate was collected, washed with ethanol, dried and purified by ethanol recrystallization.

#### B. Microwave Irradiated Synthesis of Co(II), Ni(II) and Cu(II) metal complexes

Equimolar solution of Schiff base ligands of  $2-\{[4-(5-bromo thiophene-2-yl)-6-phenyl pyrimidin-2-yl] carbon imidoyl\}$  phenol  $[HL^1] / 2-\{[4-(5-bromo thiophene-2-yl)-6-phenylpyrimidin-2-yl]carbonoimidoyl\}-5-methoxyphenol <math>[HL^2]$  (0.001 mol) and metal chloride (0.001mol) in ethanol, the reaction mixture was refluxed in microwave irradiation for 3 to 4 minutes, then the colour of the mixture was turned, it was treated with sodium acetate (0.5 g) the resultant reaction mixture was further refluxed in microwave irradiation for 2 more minutes, and then it was decomposed by pouring to distilled water (80 to 100 ml) with stirring. The separated solid (complex) was allowed to settle and collected by filtration, washed several times with distilled water and then with hot ethanol. The solid complex obtained was dried in desiccators over anhydrous calcium chloride.



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 9 Issue VII July 2021- Available at www.ijraset.com

### III. RESULTS AND DISCUSSION

The analytical data shown in Table-1: indicate that all the metal ions from 1:1 (M:L) complexes with both the ligands. All the complexes are coloured and are soluble in DMF, DMSO and acetonitrile and are sparingly soluble in other organic solvents. Molar conductance measurements of these complexes in DMF correspond to electrolytes. The cobalt(II) complexes magnetic moment is 5.22-5.38 B.M. The magnetic moment of the cobalt(II) octahedral complex is shown to be between 4.46 and 5.33 B.M.[15]. Thus, the reported magnetic moment for the investigated Co(II) complexes demonstrates that they are octahedral geometry. Nickel (II) is a d<sup>8</sup> ion with two electrons unpaired which may form six coordinated octahedral complexes[16]. The magnetic moments reported in the octahedral Ni(II) complexes vary from 2.82 to 3.4, depending on the magnitude of the orbital contribution B.M.[17-18]. In this study 3.17-3.34 B.M. were discovered to be magnetic moments at room temperature for the Ni(II) complexes. This investigation implies the octahedral structure of Ni(II) complexes. The copper(II) complexes magnetic moment is 1.85 to 1.90 B.M., which is the same as the spin only value. It is reported that the distorted octahedral geometry of copper (II) complexes devoid of spin interaction exhibit magnetic moment in the range of 1.80-1.93 B.M[19]. Hence it may be concluded that copper (II) complexes have major spin interaction. The electronic spectral band maxima with their assignments for the complexes were discussed. In the present Co(II) complexes, the transitions may be ascribed to three bands: 14725.28 cm<sup>-1</sup>, 18192.32 cm<sup>-1</sup>, and 32233.05 cm<sup>-1</sup>. These transitions indicate the octahedral structure of the present Co(II) complexes. The investigated Ni(II) complexes were having transition of three bands: 10210.05 cm<sup>-1</sup>, 12825.55 cm<sup>-1</sup>, and 22730.32 cm<sup>-1</sup>, respectively, to  $v_1$ ,  $v_2$  and  $v_3$ , indicating the octahedral geometry of the Ni(II) ion.

	Table 1. Elemental analysis and conductance data for figures and metal complexes.							
<b>C1</b>		Analysis % Found (Calculated)						Molar
SI.	Ligands / Complexes							Conductance
No.		С	Н	N	S	Br	М	$\Box^{-1} \text{cm}^2 \text{mol}^{-1}$
		-			~			
	$HL^{1}$	57.82	3.24	9.67	7.28	18.25		-
	$C_{21}H_{14}BrN_3OS$	(57.78)	(3.20)	(9.62)	(7.33)	(18.31)		_
01	$Co(L^1)2H_2O$	47.45	3.15	7.95	6.05	15.11	11.13	73
	$Co(C_{21}H_{13}BrN_3OS)2H_2O$	(47.43)	(3.20)	(7.90)	(6.02)	(15.04)	(11.09)	00
02	$Ni(L^1)2H_2O$	47.48	3.25	7.97	5.98	15.13	11.10	9.4
	Ni(C <sub>21</sub> H <sub>13</sub> BrN <sub>3</sub> OS)2H <sub>2</sub> O	(47.45)	(3.21)	(7.91)	(6.03)	(15.05)	(11.05)	84. 29
03	$Cu(L^1)2H_2O$	47.10	3.13	7.90	6.03	14.96	11.80	0.4
	Cu(C <sub>21</sub> H <sub>13</sub> BrN <sub>3</sub> OS)2H <sub>2</sub> O	(47.03)	(3.17)	(7.84)	(5.97)	(14.91)	(11.85)	84. 96
0.4	$Zn(L^1)2H_2O$	46.92	3.21	7.86	6.00	14.90	12.10	05
04	Zn(C <sub>21</sub> H <sub>13</sub> BrN <sub>3</sub> OS)2H <sub>2</sub> O	(46.87)	(3.16)	(7.81)	(5.95)	(14.85)	(12.15)	85. 40
	$HL^2$	56.55	3.38	9.05	6.92	17.18		02
	$C_{22}H_{16}BrN_3O_2S$	(56.61)	(3.43)	(9.00)	(6.86)	(17.13)		63
05	$Co(L^2)2H_2O$	47.10	3.42	7.43	5.65	14.26	10.57	02
05	$Co(C_{22}H_{15}BrN_3O_2S)2H_2O$	(47.03)	(3.38)	(7.48)	(5.70)	(14.23)	(10.50)	93. 00
	$Ni(L^2)2H_2O$	47.09	3.45	7.52	5.76	14.31	10.40	
06	Ni(C <sub>22</sub> H <sub>15</sub> BrN <sub>3</sub> O <sub>2</sub> S)2H <sub>2</sub> O	(47.05)	(3.39)	(7.49)	(5.70)	(14.24)	(10.46)	85. 02
-	$Cu(L^2)2H_2O$	46.58	3.42	7.36	5.63	14.07	11.17	02
07	$Cu(C_{22}H_{15}BrN_{3}O_{2}S)2H_{2}O$	(46.65)	(3.35)	(7.42)	(5.65)	(14.12)	(11.22)	93.
L		(10.05)	(3.55)	(7.12)	(0.00)	(1.1.2)	(11.22)	40
08	$Zn(L^2)2H_2O$	46.55	3.37	7.46	5.69	14.12	11.25	92.
2.0	$Zn(C_{22}H_{15}BrN_3O_2S)2H_2O$	(46.50)	(3.34)	(7.39)	(5.63)	(14.06)	(11.18)	32

Table 1: Elemental analysis and Conductance data for ligands and metal complexes

The values shown in the parenthesis are calculated ones.



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 9 Issue VII July 2021- Available at www.ijraset.com

Three bands are seen at 14735.81 cm<sup>-1</sup>, 18155.21 cm<sup>-1</sup>, and 32260.06 cm<sup>-1</sup> in this study, which can be assigned to the transitions  ${}^{2}B_{1g} \rightarrow {}^{2}A_{1g}$ ,  ${}^{2}B_{1g} \rightarrow {}^{2}B_{2g}$  and  ${}^{2}B_{1g} \rightarrow {}^{2}E_{g}$  respectively. According to these findings, the Cu(II) complexes have six-coordinated octahedral structure. In the IR spectra of ligands 2-{[4-(5-bromo thiophen-2- yl)-6-phenyl pyrimidin-2-yl] carbonoimidoyl} phenol  $[HL^{1}]$  and 2-{[4-(5-bromo thiophen-2-yl) -6- phenyl pyrimidin-2-yl]carbon imidoyl}-5- methoxyphenol  $[HL^{2}]$  and their complexes, The azomethine group found in the Schiff bases is attributed to the significant absorption in the 1620 cm<sup>-1</sup> region. The azomethine group[20] in metal complexes is moved to a lower frequency in the range 1600 – 1580 cm<sup>-1</sup>. The appearance of a new band around this region 1618 - 1610 cm<sup>-1</sup> due to  $v_{C=N}[21]$  adds to this. The C=N changes to a higher frequency area in the present complexes, showing that azomethine nitrogen is involved in coordination. The spectrum of the base ligands HL<sup>1</sup> and HL<sup>2</sup> displayed a medium intensity band (O-H), which is absent from the complexes spectrum, of 3364 cm<sup>-1</sup> and shows that the base ligands HL<sup>1</sup> and HL<sup>2</sup> were deprotonated before coordination by the oxygen atom. In ligands, HL<sup>1</sup> and HL<sup>2</sup> a medium intensity band is observed in the range of 1090-950 cm<sup>-1</sup> are assigned to the  $v_{(C-S)}$  vibrations[22], In complexes these bands are shifted to lower frequency due coordination through S atom to central metal. Metals have respectively heavy mass and M-S and M-O bonds have low bond order and hence the group vibration due to these occurs in the far infrared region. Based on the literature data in the present study the complexes of  $v_{M-S}$  vibration occur in the region 540–490 cm<sup>-1</sup> and  $v_{M-O}$  occur in the region 450–400 cm<sup>-1</sup> have been assigned[23]. <sup>1</sup>H NMR spectra of ligands HL<sup>1</sup> and HL<sup>2</sup> shows two singlets 2.20 δ(ppm) at 3.83 δ(ppm) methoxy (OCH3) protons. But the ligands also shows a multiplet at 6.90-7.56  $\delta$ (ppm) due to aromatic protons. In the complexes the multiplet observed between  $6.66 - 7.25 \delta$  (ppm) due to aromatic protons and the signals at 3.62 - 3.77  $\delta$  (ppm) are due to methoxy protons, were shifted to down field due to the compelsation through oxygen and S of thiophene group and nitrogen of C=N group. A signal at 11.40 - 11.51 $\delta$ (ppm) due to resonance acting of phenolic OH protons, in complexes these signals were absent due to complexation through oxygen of phenolic OH group were shifted to down field due to the complexation through nitrogen of C=N and S of thiophene group to the metal ion respectively. ESR studies of the complex have been made. The magnetic parameters have been evaluated from the spectra and tend  $g_{II} > g_{\perp}$  observed in the g' values suggests the presence of an unpaired electron in the  $d_{x^2-v^2}$ orbital[24]. Based on these observations copper(II) complex may have octahedral geometry. The covalent character of the metalligand relation has a g value less than 2.3 and larger than 2.3 for ionic character has been reported by Kivelson and Neiman[25].  $Cu(L^2)2H_2O$ , complex has a g value of 2.47 in this study, indicating that the metal-ligand interactions are more ionic in nature. The powder x-ray diffraction for Cu(C<sub>21</sub>H<sub>13</sub>BrN<sub>3</sub>OS)2H<sub>2</sub>O have been studied. It is seen that the complexes consists of some reflections between  $8 - 69^{\circ}$  (2 $\Theta$ ). In Bragg's relation, interplanar spacing(d), is computed using the locations of the intense peak ( $n\lambda$ =2d Sin 1,54056 A<sup>0</sup>), where  $\lambda$  is the wavelength of the x-ray utilised (Cu k $\lambda$ = 1,54056 A<sup>0</sup>). The calculated spacing together with relative intensities with respect to most intense peaks are recorded in table. All main peaks have index [26] and their  $\sin^2 \Theta$ values are compared with the calculated values. The comparision of these values reveals that there is good agreement with the calculated and observed values. The complex shown broad peaks indicates that amorphous nature of the complexes.

On the basis of elemental analysis, magnetic susceptibility, conductivity, electronic, IR, <sup>1</sup>H NMR, ESR spectral data and powder Xray powder diffraction observations indicated the tentative structures of these complexes for the coordination number six having octahedral geometry have been proposed.



Fig.2. Proposed structure of complexes, where M= Co(II), NI(II), Cu(II) and Zn(II)



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 9 Issue VII July 2021- Available at www.ijraset.com

- A. Determination of stability constant of Cu(II) complexes by spectrophotometric method:
- 1) Determination of  $\lambda_{max}$ : A solution of  $(10^{-2}M)$  Cu (II) is mixed a solution of  $(10^{-2}M)$  ligands 2-{[4-(5-bromo thiophen-2-yl)-6-phenyl pyrimidin- 2-yl] carbonoimidoyl} phenol [HL<sup>1</sup>] and 2-{[4-(5-bromo thiophen-2-yl) -6-phenyl pyrimidin-2-yl]carbon imidoyl}-5-methoxyphenol [HL<sup>2</sup>] solution a distinct greenish colored complexes are formed. The absorbance at 5nm wavelength interval of the complex solutions were measured in the visible range at [350-750nm], it is found that both the complexes having maximum absorbance ( $\lambda_{max}$ ) at 535nm.
- 2) *Effect of pH:* The following method was used to investigate the effect of pH on the colour intensity of the reaction mixture. In a 25 ml standard flask, 1 ml of copper(II) solution  $(1 \times 10^{-3} \text{M})$  and 10 of buffer solution of known pH (Range 1 to 10) and 1 ml of HL<sup>1</sup> and HL<sup>2</sup>  $(1 \times 10^{-2} \text{ M})$  solutions were taken and diluted up to the mark with distilled water. The absorbance of the reaction mixture was measured at 535nm at different p<sup>H</sup> and plot the graph absorbance verses p<sup>H</sup>, It is found that maximum absorbance at p<sup>H</sup> = 5. In view of the above reasons the author has selected a pH of 5 for further studies[27].
- 3) Job's method: Prepare a series of solutions, copper(II) penthydrate sulphide standard solution 0 to  $40x10^{-4}$  M was pippeted into seven volumetric flasks and the mole ratio was maintained by adding an aliquot of  $40x10^{-4}$  to 0 M HL<sup>1</sup> and HL<sup>2</sup>, respectively. All measurements were made at 3 different temperatures at 535 nm, i.e. 30, 35 and 40 °C. The curve revealed a maximum of  $\chi$ metal = 0.5 in a mole fraction indicating the development of a complex with a ratio of 1:1 metal to ligand[28]. The following is the Job's approach equation used in this study:



Fig 3: Effect of pH on absorbance

$$K_f = \frac{[ML]}{[m][L]}$$
$$A_2 / A_1$$

$$Kf = \frac{2}{[1 - A_2 / A_1] \times [C_L - C_M \times A_2 / A_1]}$$

Where,  $A_1$  = absorbance at break point,  $A_2$  = actual absorbance,  $C_M$  = concentration of metal, and  $C_L$  = concentration of ligand.



Fig. 4: Representative Job's curves of equimolar solutions of Cu(II) complexes with (HL<sup>1</sup>) at  $30^{0}$ C,  $35^{0}$ C and  $40^{0}$ C



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 9 Issue VII July 2021- Available at www.ijraset.com

4) *Mole Ratio Method:* By using the mole ratio method[29], at constant Cu (II) concentration  $(20x10^{-4})$  and varying HL<sup>1</sup> and HL<sup>2</sup> concentrations (0 to  $40x10^{-4}$ ) at 535nm and at these three temperatures, a sharp bands were observed at 1:1mole ratio of HL<sup>1</sup> and HL<sup>2</sup> with Cu(II). A graph is plotted between HL<sup>5</sup> and HL<sup>6</sup> ligands volume and the absorbance (Fig.5.). Mole ratio method also confirms that the ratio between the metal ion and the reagent is 1:1(M:L). The corresponding equation for mole ratio method is as follows:

$$Kf = \frac{\frac{A}{\epsilon b}}{\left[C_M - \frac{A}{\epsilon b}\right] \times \left[C_L - \frac{A}{\epsilon b}\right]}$$

Where  $\varepsilon b$  = molar absorptivity constant, and A = absorbance at peak point. The stability constant has been determined using Job's and mole ratio approaches, based on the data provided[30-31], and the mean value of K<sub>f</sub> obtained using two separate approaches is in fair agreement (Table-2).



Fig. 5.: Representative Mole ratio curves for Copper(II) complexes with HL<sup>1</sup> at  $30^{\circ}$ C , $35^{\circ}$ C and  $40^{\circ}$ C

Table 2: Metal-ligand formation constants calculated from Job's continuous variation method and mole ratio method a	at
$30^{0}$ C, $35^{0}$ C and $40^{0}$ C and P <sup>H</sup> of $5 \pm 0.05$	

S.No	Methods	Metal: Ligand	Formation Co	Formation Constants at temperatures			
			25 <sup>0</sup> C	35 <sup>0</sup> C	$45^{0}C$		
1	Job's method	1:1	13.06x10 <sup>4</sup>	14.51x10 <sup>4</sup>	10.40x10 <sup>4</sup>		
a.	Copper(II) complexes with (HL <sup>1</sup> )						
b.	Copper(II) complexes with (HL <sup>2</sup> )	1:1	13.18x10 <sup>4</sup>	15.06x10 <sup>4</sup>	$11.23 \times 10^4$		
2	Mole ratio method	1:1	$0.24 \text{x} 10^4$	$0.57 \mathrm{x} 10^4$	$0.77 \mathrm{x} 10^4$		
А	Copper(II) complexes with (HL <sup>1</sup> )						
В	Copper(II) complexes with $(H_2L^2)$	1:1	$0.28 \times 10^4$	$0.55 x 10^4$	$0.79 \mathrm{x} 10^4$		



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 9 Issue VII July 2021- Available at www.ijraset.com

## B. Antimicrobial Activities

Both the investigated ligands, of 2-{[4-(5-bromo thiophene-2-yl)-6-phenyl pyrimidin-2-yl] carbon imidoyl} phenol [HL<sup>1</sup>] and 2-{[4-(5-bromo thiophene-2-yl)-6-phenylpyrimidin-2-yl]carbonoimidoyl}-5-methoxyphenol [HL<sup>2</sup>] and their metal complexes were screened for their antibacterial and antifungal activity by cup plate method, at different concentrations in DMSO solvent by minimum inhibitory concentration (MIC) method. Antibacterial activity in opposition to *Escherichia coli* and *Staphylococcus aureus* and Antifungal activity in opposition to *Aspergillus niger* and *Cladosporium* were carried out. A normal drugs gentamycine and fluconazole were also tested for their antibacterial and antifungal activity at the same concentration under the circumstances analogous to that of the test compound. The results (given in Table-3) suggest that the among all the complexes Cu(II) complexes have exhibited good bacterial and fungal activities which is due to the faster diffusion of the Cu(II) complexes[32-33], whereas, ligands (HL<sup>1</sup>) and (HL<sup>2</sup>) have shown moderate activity.

Ligands /	Zone of inhibition in mm						
Complexes	Antibac	cterial activity	Antifungal activity				
	Escherichia Coli	Staphylococcus Aureus	Aspergillus Niger	Cladosporium			
$HL^1$	13	12	14	13			
Co(II) with HL <sup>1</sup>	18	17	15	17			
Ni(II) with HL <sup>1</sup>	16	18	16	18			
Cu(II) with HL <sup>1</sup>	24	22	23	24			
$HL^2$	15	17	17	18			
$Co(II)$ with $HL^2$	15	14	13	14			
Ni(II) with HL <sup>2</sup>	16	17	19	16			
Cu(II) with HL <sup>2</sup>	15	18	17	17			
Gentamycine	22	23	24	22			
Fluconazole	18	16	16	17			

Table 3: The antimicrobial activity of ligands and their Co (II), Ni(II), and Cu(II) complexes evaluated by (mm).

## C. Antioxidant Activity

The utilization the stable free radical diphenylpicrylhydrazyl is one such approach that is (DPPH)[34]. This currently popular approach is based on the creation of the non-radical form DPPH-H when DPPH is reduced in methanol solution in the presence of a hydrogen donating antioxidant[35]. Therefore, the present study has been carried out to assess the antioxidant the newly synthesized Co(II), Ni(II), Cu(II) and Zn(II) metal complexes of the Schiff's base ligands  $2-\{[4-(5$ activity of bromothiophen-2-yl)-6-phenylpyrimidin-2-yl]carbon imidoyl} phenol [HL<sup>5</sup>] and 2-{[4-(5-bromothiophen-2-yl)-6-phenylpyrimidin-2-yl] carbon imidoyl-5 metho xyphenol [HL<sup>6</sup>]. The ligands and their complexes were diluted in methanol to a final concentration of various stock solutions (mg/ml), 2.5 ml of test solution of various concentrations was added to 1 ml of DPPH solution (0.1 mmol) and allowed to react at room temperature. The absorbance was measured at 517 nm after 30 minutes, using butylated hydroxyl anisole (BHA) as standards. On the y-axis, percentage scavenging effects were plotted, and on the x-axis, concentration (g/ml)[36]. The results of DPPH scavenging activity both the ligands and their metal complexes were presented in figure-6. The analysis revealed that metal complexes of Co(II), Ni(II), Cu(II), and Zn(II) scavenge DPPH more efficiently than ligands, implying that they have better free radical scavenging and antioxidant characteristics.



Fig. 6. Antioxidant results of ligands 2-{[4-(5-bromo thiophene-2-yl)-6-phenyl pyrimidin-2-yl] carbon imidoyl} phenol [HL<sup>1</sup>] and their metal (II) complexes



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 9 Issue VII July 2021- Available at www.ijraset.com

### IV. CONCLUSION

A few complexes of 2-{[4-(5-bromo thiophen-2-yl)-6-phenyl pyrimidin-2-yl] carbonoimidoyl} phenol [HL<sup>1</sup>] and 2-{[4-(5-bromo thiophen-2-yl) -6- phenyl pyrimidin-2-yl]carbon imidoyl $\}$ -5-methoxyphenol [HL<sup>2</sup>] with first row transition metal ions have been reported in this chapter. These complexes analyze for the stoichiometry of the type  $(1:1) [M(L)]2H_2O$ . The conductivity data shows the electrolytic nature of these complexes. The magnetic susceptibility measurements obtained at room temperature for Co(II), Ni(II) Cu(II) complexes agrees with the spin only values. The electronic, IR, <sup>1</sup>H NMR, ESR spectral data and X-ray diffraction data suggest that Co(II), Ni(II), Cu(II) and Zn(II) complexes with ligands (HL<sup>1</sup> and HL<sup>2</sup>), exhibit coordination number six. The formation of copper(II) complex with ligands has been studied spectrophotometrically at an absorption maximum of 535 nm at different temperatures. The data shows that copper(II) and (HL<sup>1</sup>) and (HL<sup>2</sup>) combine in the molar ratio of 1: 1 at  $p^{H}$  5.0. The stability constants of the complexes were calculated to be  $13.06 \times 10^4$ ,  $14.51 \times 10^4$  and  $10.40 \times 10^4$  and  $13.18 \times 10^4$ ,  $15.06 \times 10^4$ ,  $11.23 \times 10^4$  for  $(H_2L^1)$  and  $(H_2L^2)$  respectively by Jobs continuous variation method and  $0.24 \times 10^4$ ,  $0.57 \times 10^4$ , and  $0.77 \times 10^4$  and  $0.28 \times 10^4$ ,  $0.55 \times 10^4$ ,  $0.79 \times 10^4$  for (HL<sup>1</sup>) and (HL<sup>2</sup>) respectively by mole ratio method at 30, 35 and 40°C, respectively. The results of antimicrobial activity data suggest that the among all the complexes Cu(II) complexes have exhibited good bacterial and fungal activities which is due to the faster diffusion of the Cu(II) complexes, whereas, ligands (HL<sup>1</sup>) and (HL<sup>2</sup>) have shown moderate activity. Antioxidant analysis results of DPPH scavenging activity both the ligands and their metal complexes were studied. The analysis revealed that metal complexes of Co(II), Ni(II), Cu(II), and Zn(II) scavenge DPPH more efficiently than ligands, implying that they have better free radical scavenging and antioxidant characteristics.

#### V. ACKNOWLEDGEMENTS

The authors are thankful to Poojya Dr.Sharanbasvappa Appa, President Sharanbasveshwar Vidya vardhak Sangha, Kalaburagi, Dr. Anilkumar Bidve, Dean, Principal Dr. Basavaraj Mathapathi, Appa Institute of Engineering and Technology, Kalaburagi, Karnataka, for their constant encouragement and support during the process of performing this work.

#### REFERENCES

- [1] Cordeu L., Cubedo E., Bandres E., Rebollo A., Saenz X., Chozas H., Dominguez M.V., Echeverria M., Mendivil B., Sanmartin C., Palop J.A., Font M. and Garcia-Foncillas J. Biological profile of new apoptotic agents based on 2,4-pyrido[2,3-d]pyrimidine derivatives, Bioorg. Med. Chem., 15; 1659-1669 ,2007
- [2] Mohamed Y.A., ABD El- Galil E. Amr, Mohamed S.F., Abdalla M.M., Al-omr M.A. and Shfik S.H., Cytotoxicity and anti-HIV evaluations of some new synthesized quinazoline and thioxopyrimidine derivatives using 4-(thiophen-2-yl)-3,4,5,6-tetrahydrobenzo[h]quinazoline-2(1H)-thione as synthon., J. Chem. Sci.,124(3); 693-702.,2012
- [3] Gurdeve M.B., Kamo K., Yaglioglu A.S., Budak Y. and Ceylen M., Synthesis and in vitro anticancer evaluation of 1,4-phenylene-bis-pyrimidine-2-amine derivatives., Turk. J. Chem., 41;263-271., 2017.
- [4] Deshmukh M.B., Salunkhe S.M., Patil D.R. and Anbhule P.V.A novel and efficient one step synthesis of 2-amino-5-cyano-6-hydroxy-4-aryl pyrimidines and their anti-bacterial activity ,Eur. J. Med. Chem., 44, 2651-2654, 2009.
- [5] Stephen G. Pyne, Andrew S. Davis, Nicole J. Gates, Joseph P. Hartley, Karl B. Lindsay, Theeraphan Machan, Minyan Tang, Asymmetric Synthesis of Polyfunctionalized Pyrrolidines and Related Alkaloids., Synlett (15): 2670-2680, 2004.
- [6] Hockova D., Holy A., Masojidkova M., Andrei G., Snoeck R., De Clercq E. and Balzarini J., Synthesis and antiviral activity of 2,4-diamino-5-cyano-6-[2-(phosphonomethoxy)ethoxy]pyrimidine and related compounds., Bioorg. Med. Chem., 12;3197-3202, 2004.
- [7] Fujiwara N., Nakajima T., Ueda Y., Fujita H. and Kawakami H.Novel piperidinylpyrimidine derivatives as inhibitors of HIV-1 LTR activation, Bioorg. Med. Chem., 16; 9804-9816, 2008..
- [8] Amin K.M., Awadalla F.M., Eissa A.A.M., Abou-Seri S.M. and Hassan G.S., Design, synthesis and vasorelaxant evaluation of novel coumarin-pyrimidine hybrids., Bioorg. Med. Chem., 19; 6087-6097,2011.
- [9] Alam O., Khan S.A., Siddiqui N., Ashan W., Warma S.P. and Gilani S.J., Antihypertensive activity of newer 1, 4-dihydro-5-pyrimidine carboxamides: Synthesis and pharmacological evaluation., Eur. J. Med. Chem.,; 45,;5113-511.2010.
- [10] Amin K.M., Hanna M.M., Ebo-Youssef H.E. and George R.F., Synthesis, analgesic and anti-inflammatory activities evaluation of some bi-, tri- and tetracyclic condensed pyrimidines., Eur. J. Med. Chem.; 44:4572-4584.2009.
- [11] Dhandayutham Saravanan,, R. Kannadhasan, Murugesan Sankaranarayanan., Synthesis and antimicrobial evaluation of novel sulphur bridged 2-oxo-1,2,3,4-tetrahydropyrimidine derivatives., Research journal of chemistry and environment 21(9):21-30,2017
- [12] Vogel A.I., A Text book of Quantitative Inorganic Chemistry, ELBS and Longmanns Green and Co. Ltd., London, 3rd ed.1962.
- [13] Setwood P.W., Magneto Chemistry, 2nd ed., Interscience, New York, 1958.
- [14] Zumdal S.S., Chemistry, 4th ed., Houghton Mifflin Company, 1997.
- [15] Kumar Shalin, Dhar Durga Nath Saxena, P N., Applications of metal complexes of Schiff bases-A review., Journal of scientific and industrial research., 68;181-187, 2009.
- [16] F.A.Cotton and G.Wilkinson, "Advanced Inorganic Chemistry", Weley Easterns Pvt. Ltd., 1969.
- [17] Suman Malik1, Supriya Das1, Bharti Jain2, Transition Metal Complexes of Omeprazole An Anti-Ulcerative Drug., Ecl6 ca Química, 36 (3), 31-36, 2011.
- [18] N.B. Kolthup, L.H.Dally, S.E.Wiberley, Introduction to Infrared and Raman spectroscopy, Acad. Press, New York, 1964
- [19] M Kato, H B Jonassen and J C Fanning. Copper(II) Complexes with Subnormal Magnetic Moments , Chem. Rev.; 64; 99, 1964.



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429

Volume 9 Issue VII July 2021- Available at www.ijraset.com

- [20] C N R Rao "Chemical application Infrared spectroscopy" Academic press New York,; 260, 262., 1963.
- [21] K.K. Mohammed Yusuff., S Mayadevi Synthesis and Characterization of Some New Transition Metal Complexes of the Schiff Base Derived from Quinoxaline-2-Carboxaldehyde and 2- Aminophenol., Synthesis and Reactivity in Inorganic and Metal-Organic Chemistry., ; (2);27,2006.
- [22] M F Iskandar, L E I Sayed, A F M Hafry, and S E Zayan,, Coordination compounds of hydrazine derivatives with transition metals—XII Nickel(II) and copper(II) chelates of bis(n-salicylidene)dicarboxylic acid dihydrazides J. Inorg. Nucl. Chem.1976; 38; 2209.
- [23] K Nakamoto "IR Spectra of Inorganic and Coordination compounds" Wiley Inter-sciences New York. 1976.
- [24] B.A. Goodmm and J.B. Rayner, "Advanced in Inorganic Chemistry and Radio Chemistry" Ed.By H.T. Emelous and A.G. Sharpe,; 13;135,1970..
- [25] D.Kivelson and R.Neiman., ESR Studies on the Bonding in Copper Complexes, J.Chem. Physc.,; 35; 149,1961.
- [26] B D Cullity "Elements of X-ray diffraction", Wesley publishing company, Inc., England, 1959.
- [27] Tong Li, Yuheng Bao, Huiqian gQiu, Weijun Tong., Boosted peroxidase-like activity of metal-organic framework nanoparticles with single atom Fe(III) sites at low substrate concentration., Analytica Chimica Acta, ; 1152,; 338299.,2021.
- [28] E. J. Scott, L. H. Tung and H. G. Drickamer, Diffusion through an Interface J. Chem. Phys.;19;1075,1951.
- [29] L.Sacconi and M.Ciampolini, Pseudo-tetrahedral structure of some α-branched copper(II) chelates with Schiff bases., J. Chem.Soc ,; 276,1964.
- [30] Billig E, Williams R W, Bernal I and Gray H.B., The Electronic Structures of Square-Planar Metal Complexes. II. The Complexes of Maleonitriledithiolate with Copper(II), Nickel(II), Palladium(II), and Platinum(II)., Inorg. Chem.,; 3; 663.1964,
- [31] K. P. Rakesh, C. S. Shantharam, D. M. Suyoga Vardhan, N. M. Made Gowda, M. B. Sridhara and D. Channe Gowda., Synthesis and antiglycation activity of benzene-1,3,5- tricarboxylic acid mediated new series of schiff base derivatives, ejpmr, ;3(3);362-367, 2016
- [32] L. Mishra and V.K. Singh, Indian J. Chem., Synthesis, structural and antifungal studies of Co(II), Ni(II), Cu(II) and Zn(II) complexes with new Schiff bases bearing benzimidazoles.;32A;446. 1993.
- [33] R. Malhotra, S. Kumar and K.S. Dhindsa, Synthesis, characterization and microbiocidal activity of cobalt(II), nickel(II), copper(II) and zinc(II) complexes of aryloxyacetic acid hydrazides, Indian J. Chem., 32A; 457,1993.
- [34] Philip molyneux The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity, SJST, 26; 211. 2004.
- [35] P. Jegadeeswari, E. D. Daffodil and V. R. Mohan, Quantification of total phenolics, flavonoid and in vitro antioxidant activity of Aristolochia bracteata retz. Int J Pharm Pharm Sci, 6;747.2014
- [36] P. Ikechukwu, Ejidike and A. A. Peter,; Synthesis, Characterization and Biological Studies of Metal(II) Complexes of (3E)-3-[(2-{(E)-[1-(2,4-Dihydroxyphenyl) ethylidene]amino}ethyl)imino]-1-phenylbutan-1-one Schiff Base, Molecules, 20; 9788. 2015.











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24\*7 Support on Whatsapp)