



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 9 Issue: IX Month of publication: September 2021 DOI: https://doi.org/10.22214/ijraset.2021.37912

www.ijraset.com

Call: 🕥 08813907089 🔰 E-mail ID: ijraset@gmail.com



Diversity and Seasonal Variations of Zooplankton in Kampli Water Tanks in and Around Hosapete City, Karnataka (India)

G. Mallanagoud¹, Vijaya S Gulalakayi², L. R. Dindalkoppa³, S. Manjappa⁴, Suresh B⁵

^{1, 2}Department of Zoology, Veerashiva College Ballari- 583104, Karnataka, India
³Department of Zoology, KRCES GGD Arts BMP Com & SVS Science College Bailhongal, Belgavi- 591102, Karnataka, India
⁴Department of Chemistry, University BDT College of Engineering, Davangere – 577005, Karnataka, India
⁵Department of Civil Engineering, Bapuji Institute of Engineering & Technology, Davangere – 577004, Karnataka, India

Abstract: This study focuses on assessment of zooplankton with special reference to seasonal variation and selected physicchemical variables of Kampli water tanks in Hosapete city, Karnataka state. A large quantity of manmade activities near the water tank. During the study period, 48 species of zooplankton - 14 species of Rotifera, 12 species of Cladocera, 6 species of Ostracoda and 4 species of Copepoda were observed. Among zooplankton, Rotifera was (645 no./Org) noticed as the dominant group in the entire the study period and the maximum count was identified in the summer period while less numbers were noticed during winter season. Zooplankton community is also significantly correlated with some physico-chemical variables. The analytical results during the study period indicate that the scattering and density of zooplankton species were encouraged by prevailing physical and chemical features of the aquatic ecological condition.

Keywords: Kampli, water tank, zooplankton, diversity, seasonal and domestic activity.

I. INTRODUCTION

Water is one of the abundantly available substance in nature. It is an essential constituent of all the animals and vegetable matter and forms about 75% of the matter of earth's crust. Water is the mother liquid of all forms of life. It is the vital essence, miracle of nature and the great sustainer of life. The essentiality of water for living systems is quite evident as without water, there is no life (Omar WMW, 2010). Water pollution due to organic materials is one of the most significant issues in present days. Most of the freshwater bodies are under remarkable pressure from human communities and developmental actions in an around the water tanks. Increasing in the addition of nutrient into the water tanks from the surrounding has been deteriorating water quality of surface water ecosystems (Kamble, et al., 2005). Physico-chemical constraints of any surface water tanks though, provide a good indication about the chemistry and quality of water. These variables will not give the clear picture of the ecological condition of the surface water body due to lack of proper assimilation with ecological factors (Karr, et al., 2000). Plankton are free moving, unicellular, microscopic and colonial autotrophic organisms that grow in aquatic environments whose movement is more or less dependent upon water movements (Cecilia Medupin, 2011 and Suresh, 2015). Plankton are also called as biotic communities, microscopic organism and minute species since these are living inside the water. The biotic community is the outcome of the integration and interaction of different physical, chemical and geo-morphological characteristics of any water body, biological appraisal is a useful alternative in assessing those systems (Stevenson and Pan, 1999). Plankton are considered as significant component of aquatic fauna and flora, play a key role in maintaining equilibrium between abiotic and biotic components of aquatic ecosystem (Arfi, et al., 2003). The surface water bodies are most significant for all livening organism including other human activities like drinking and agricultural practices. Research studies on the zooplankton counting aspects are of great significance in developing resources of a water tanks including all types of water body. A number of researchers have studied the zooplankton diversity of lentic and lotic water bodies (Kar and Kar (2013); Pullie and Khan (2003); Manjare, (2015); Miah, et al., (2013); Suresh, 2015). In India 80 % of the surface water is vulnerable to pollution as more than 95 % of the sewage in the country is not treated. Lotic water bodies like rivers and streams play a very significant role in maintaining the bio-diversity and over all ecological balance in nature. However, the water quality of fluvial systems is deteriorating due increase in the amount of raw sewage entering the rivers. The increase of pollution is caused by population growth and increasing urbanization. Related to this is the industrialization that also causing huge environmental problems (Zargar and Ghosh, 2006). Karnataka state is endowed with 6.31 lakhs hectare of freshwater resources consisting of 4.15 lakhs hectare which includes ponds and tanks and 2.16 lakh hectare reservoirs. In addition, the state has 6000 kms of river stretch and 3000 kms length of canal. Water pollution is the introduction into fresh/ground/ocean waters of chemical, physical or biological material that degrades the quality of the water and affects the organisms living in it (Pandey, et al., 2009).



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 9 Issue IX Sep 2021- Available at www.ijraset.com

The main objectives of the study are: collection of water samples using plankton net at three different water tank in the Hosapete city of Karnataka state. Since, no work have so far neither been done in enumerating zooplankton nor in analyzing water quality in Kamapli water tank in Hosapete city. The present attempt made an endeavor to appraise the water quality variables and to assess pollution status of the Kampli water tanks using Palmer's scale.

II. MATERIALS AND METHODS

A. Topography Of The Study Area

Hospet is a town head quarter situated 66 Kms away from Bellary district in the Central part of Karnataka state, India. Hospet-Shimoga Highway (SH-25) passes through the study area. Almost all the villages of the area are connected by unmetalled and metalled roads and regular bus facility exits from Hosapete to different villages. The study area falls in the survey of India topo map numbers 57 B/6 on 1:50000 scale. The area is bounded by 14.74° to 14.88° N latitude and 75.88° to 76° E longitude. The location map of the study area is represented in Fig. 1. Topography of the study area is generally undulating to rolling topography with frequent mound like structures. Soils of the area are affected by erosion. Isolated hills and hill ranges are also seen. The geology (rock) of the study area consists of metamorphic rocks like gray wacke, argillite and granodiorite and tonalitic gneisss. The study area received a maximum rainfall of 742 mm in the year 2015 and a minimum of 361.9 mm in the year 2003. The normal rainfall of the study area is 656.70 mm.

1) Kampli Tank: It is a natural, perennial fresh water body situated in the Hosapete city and located 35 km away from Hosapete. The water body lies at 150.301 N latitude and 760.61 E longitude. The area of the water body is 20 acres and depth is about 8 feet. The colour of the water body is pale greenish. The water body is rectangular in shape. The Kampli tank received water from rainfall, city sewage and agricultural run-off. The water is used to grow the crops like paddy, sugarcane, banana and some vegetable crops. Besides this, water is also used for washing of vehicles, cattle washing and other domestic activities.

B. Methods

The water samples for physico- chemical as well as zooplankton analysis were collected at monthly interval for a year from March 2019 to February 2020 from three collection points taking randomly at Kampli water tank. The data thus generated were summed up as average data on the basis of seasons viz. summer (April to July), monsoon (August to October) and winter (November to March). Grab surface water samples were collected in all the selected water tanks and were analyzed for the physico-chemical variables (temperature, pH, total dissolved solids (TDS), and dissolved oxygen (DO)) in situ using and pH meter and conductivity meter. DO was estimated as per the standard method means Wrinkles methods (APHA 2012). The zooplankton sampling was carried out by filtering 50 L of water through a planktonic net and was placed in 20 ml plastic vials to which 4% formalin was added for preservation. The preserved zooplankton samples were scanned under compound microscope in the laboratory at magnifications of 75x to 300x and were further identified using the taxonomic keys Tenenbaum, *et al.* (2004).

III. RESULT AND DISCUSSION

Water quality data of the selected Kampli water tank were showed in the Table 1. The surface water temperatures recorded during the study period was between the range from $25.20(\pm 1.21)^{\circ}$ C at Kampli tank during Winter Season to $28.90 (\pm 1.38)^{\circ}$ C during monsoon. The maximum temperature was observed during monsoon the study period while the minimum was found during winter season. Yadav et al. (2013), Niroula et al. (2010) observations at urban ponds are agreement with the current results. The pH value of the selected water tank in and around the Hosapete city during the study period was in between 7.24 (\pm 0.29) at Somalapura water tank during the summer season to 8.02 (± 0.52) during monsoon at Kampli water tank. In the present study Kampli water tank showing the maximum value of pH and were slightly higher towards alkaline range, it is extremely acidic during monsoon season. In this present investigation dissolved oxygen concentration of the selected water tanks in and around the Hosapete city was varied from 8.60 (\pm 2.89) mg/l during winter to 6.10 (\pm 1.68) mg/l during monsoon at Kampli water tank. From the present study the results area revealed that and was noticed that the monsoon rain play key role in seasonal dynamics of studied physicochemical properties of the water samples. The runoff water during the rainy season carried large amount of organic matter in the form of community and home waste to the selected water tanks in and around the Hosapete city. As the runoff water were rich in clay, silt and colloidal organic matter which also attributed for excessive plankton growth and thus increase turbidity during monsoon season (Radhika et al. 2004, Pathak and Limaye, 2012 and Dhanalakshmi et al. 2013). Total dissolved solids (TDS) was varied 198.6 (± 3.84) at Kampli water tank during monsoon season to 139.2 (\pm 2.36) mg/l during summer season. The highest value was recorded during monsoon and the lowest was observed during summer season period.



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 9 Issue IX Sep 2021- Available at www.ijraset.com

Raised ionic concentration due to nutrient deposition and organic pollution attributed highest electrical conductivity (Fokmare and Musaddique, 2001). In the present study TDS was higher during High during monsoon since TDS and EC are most correlated. Water temperature could raise the rate of microbial decomposition of the rain water carried organic load resulting reduction of dissolved oxygen content in water sample (Hulyal and Kaliwal, 2011; Ramulu and Benarjee, 2013) and on the other hand. (Dhanalakshmi, et al., 2013) particularly during monsoon.

A. Statistical Analysis

The statistical analysis of Pearson's correlation coefficient is presented in the Table 2. The surface water temperature was significantly positively correlated with total dissolved solids and temperature. On the other hand, surface water temperature showed strict negative relation pH in water tanks in all the selected water tanks also. The pH showed significant negative correlation with temperature and dissolved oxygen during the entire study period and all the selected water tanks in and around the Hosapete city. Dissolved oxygen has positive correlation with temperature and showed negative correlation with pH. Due to the accelerated microbial decomposing activity the requirement of oxygen was increased (Mwebaza-Nadwula, *et al.*, 2005) resulting lower value of DO during monsoon season. Runoff from the surrounding human settlement consisting domestic sewage rich in organic matters was the main cause of nutrient enrichment of the selected water tanks in and around the Hosapete city (Verma *et al.*, 2012).

B. Zooplankton Study

A total of 14 genera of zooplankton individuals were collected in the present study. Of these, the most abundant taxon was Rotifera (5 or 35.71%), which was distantly followed by Cladosera (3 or 24.420%), Copepoda (3 or 21.42%) and finally Ostracoda (2 or 14.28%). The relative abundance of the major taxa of phytoplankton are presented in Table 4 alongside that of species (Table 3). Kampli – 1 (K1) showed a maximum species compared to other two (Kampli-II and III) location suring the study period. At Kampli-II and III also the Rotifera (72% and 57%) were dominant followed by Cladosera (30% and 25%), Copepoda (20% and 25%) and Ostrocoda (10% and 12.5%).

In the current study, at Kampli water tank, the occurrence of *Brachionus sp., Conochilus sp., Plationus sp., Daphnia sp., Alonella sp., Moina sp., Cyclops sp., Heliodiapt omus sp., Cyprinotus sp.,* was indicating pollutants of organic and biological origin. The current study agreed with the observation of Gadag et al. (2005). The pollution level was observed to increase in the Kampli water tank in Hosapete city onwards as it was confirmed by using Shannon-Weaver index (Table 3). The abundance of *Daphnia sp., Alonella sp., Moina sp., Cyclops sp., Heliodiapt omus sp.,* were maximum at Kampli water tank this indicating the highest degree of organic pollution. Unpolluted Kampli (K3) is slightly characterized by abundance of Rotifera species followed by cladocera and Copepoda, as it was supported by earlier workers (More and Nandan, 2000; Nandan and Aher, 2005; Tas and Gonulal, 2007).

Distribution of zooplankton depends partly upon the aquatic environment, their requirements and their range of tolerance. The organisms with many requirements and a limited range of tolerance are very narrowly distributed and usually rare (Amphorn Sakset and Wanninee Chankaew, 2013).

The distribution of zooplankton may explain disparities in Frequency in the present study. Among the zooplanktons, 14 (100%) occurred in Tank-I (K1) (See Table 4). The various zooplankton taxa presented different presence performances: most species in the Rotifera class showed presence in all the selected locations of Kampli water tank the trends with the Tank 1 - K1 (14), Tank II - K2 (10) andf Tank III - K2 (8).

A study of dominance among the zooplankton taxa shows that the Rotifera, followed by the Cladocera, Copepoda and Ostrocoda were dominant during the study period in the Kampli water tank of the Hosapete city. However we believe that the habitat suffers frequent variability, and according to Tiwari et al., (2006), conditions are not ideal for any one competing species and the competitive advantage swings from one species to another before the latter has had the opportunity to replace the former.

The seasonal variation of species diversity index in 4 studied sectors is given in Table 3. The index is based on the principle that in clean water, the species diversity is high while, in polluted water the diversity becomes low. The Shannon-Weaver diversity index proposed as diversity index greater than (>4) is clean water; between 3-4 is mildly polluted water; between 2-3 is moderately polluted water and less than 2(<2) is heavily polluted water. The index computed in the present analysis showed that zooplankton species diversity ranged from 2.188 in Tank I (K1) location water representing moderately polluted water (medium the Shannon-Weaver index medium level of pollution), 2.924 in Tank III (K3) water indicating mildly polluted water and 4.285 in Tank III (K3) water indication slightly polluted water in Kampli lake.



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 9 Issue IX Sep 2021- Available at www.ijraset.com

IV. CONCLUSION

In the present investigation, based on physico-chemical parameters and zooplankton load was higher at Tank – I (K1) and Tank – II (K2). Hence, Shannon-Weaver index values observed to be lower (Table 3) in Tank – I (K1) and Tank – II (K2) water tank. The index was maximum at Tank – III (K3) location, since the water is slightly due to absence of human anthropogenic activities near this location. Comparatively, Tank-II (K2) location showing is mildly polluted when compared to Tank-I (K1) and Tank –II (K2) location because of there is no human activity near and around these locations.

It is summarized from the results that Kampli water tank in the Hosapete city, which are the most productive water tanks. The results show that the improvement of diversity index from Kampli water tank, was due to the decline in pollution level. The findings of this investigation clearly revealed that in respect to domestic waste and human activity the pollution, zooplankton perhaps were more tolerant to pollution. The study emphasizes the necessity of using zooplankton as effective and appropriate method of biomonitoring for evaluation of lentic water quality.

V. ACKNOWLEDGEMENT

No conflict for interest

Parameters	IS Standards	Summer Season				
		Tank – I (K1)	Tank – II (K2)	Tank – III (K3)		
Temperature (^o C)		26.81 ± 0.05	26.90 ± 0.05	26.20 ± 0.11		
рН	6.5 - 8.5	7.89 ± 0.02	7.92 ± 0.02	7.24 ± 0.03		
TDS (ppm)	1000	145.6 ± 5.84	148.5±0.69	139.2±1.36		
DO (ppm) minimum	5.0	7.1±0.09	7.3±0.02	7.8±0.03		
		Winter Season				
Temperature (^o C)		25.01 ± 0.05	26.90 ± 0.05	25.20 ± 0.11		
рН	6.5 - 8.5	7.56 ± 0.02	7.32 ± 0.02	7.96 ± 0.03		
TDS (ppm)	1000	165.6 ±5.84	171.5±0.69	151.2±1.36		
DO (ppm) minimum	5.0	8.1±0.09 8.3±0.02		8.6±0.03		
			Monsoon Season	n		
Temperature (^O C)		27.08 ± 0.05	28.90 ± 0.05	27.20 ± 0.11		
pН	6.5 - 8.5	8.01 ± 0.02	8.02 ± 0.02	7.96 ± 0.03		
TDS (ppm)	1000	198.6 ±5.84	156.5±0.69	161.2±1.36		
DO (ppm) minimum	5.0	6.1±0.09	6.3±0.02	6.8±0.03		

Table 1. Mean Value of seasonal variation in Physico-chemical Parameters of Kampli water

Table 2 Correlation	n sigificance betwe	een selectd physico	-chemical variables
---------------------	---------------------	---------------------	---------------------

	Tank – I (K1)			Tank – II (K2)			Tank – III (K3)					
	pН	DO	Temp.,	TDS	pН	DO	Temp.,	TDS	pН	DO	Temp.,	TDS
pН	1.00				1.00				1.00			
DO	-0.08	1.00			-0.09	1.00			-0.06	1.00		
Temp.,	-0.48	0.69	1.00		-0.56	0.70	1.00		-0.51	0.71	1.00	
TDS	0.86	-0.72	0.61	1.00	0.79	-0.68	0.70	1.00	0.82	-0.58	0.59	1.00

Note: in bold correlation is significant at the level (2 tailed)



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429

Volume 9 Issue IX Sep 2021- Available at www.ijraset.com

		Locations in and	around the Hospet	t City		
				Kampali	Kampali	Kampali
				Tank - I	Tank – II	Tank - III
Zooplankton	Rotifera	Rotifera	Rotifera			
diversity	Brachionus	Brachionus	Conochilus sp.			
	sp.	sp.	Polyarthra sp			
	Conochilus	Conochilus	Plationus sp			
	sp.	sp.	Cladosera			
	Polyarthra sp	Filinia sp.	Alonella sp			
	Filinia sp.	Plationus sp.	Diaphanos			
	Plationus sp	Cladosera	oma sp.			
	Cladosera	Daphnia sp	Copepoda			
	Daphnia sp	Alonella sp	Nauplius sp.			
	Alonella sp	Moina sp.	Heliodiapt			
	Diaphanos	Copepoda	omus sp			
	oma sp.	Cyclops sp.	Ostracoda			
	Moina sp.	Heliodiapt	Cypris sp			
	Copepoda	omus sp				
	Cyclops sp.	Ostracoda				
	Nauplius sp.	Cyprinotus sp.				
	Heliodiapt					
	omus sp					
	Ostracoda					
	Cypris sp					
	Cyprinotus sp					







International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429

Volume 9 Issue IX Sep 2021- Available at www.ijraset.com

Parameters	IS Standards	Summer Season				
		Tank – I (K1)	Tank – II (K2)	Tank – III (K3)		
Temperature (^O C)		26.81 ± 0.05	26.90 ± 0.05	26.20 ± 0.11		
pН	6.5 - 8.5	7.89 ± 0.02	7.92 ± 0.02	7.24 ± 0.03		
TDS (ppm)	1000	145.6 ±5.84	148.5±0.69	139.2±1.36		
DO (ppm) minimum	5.0	7.1±0.09	7.3±0.02	7.8±0.03		
		Winter Season				
Temperature (⁰ C)		25.01 ± 0.05	26.90 ± 0.05	25.20 ± 0.11		
pН	6.5 - 8.5	7.56 ± 0.02	7.32 ± 0.02	7.96 ± 0.03		
TDS (ppm)	1000	165.6 ±5.84	171.5±0.69	151.2±1.36		
DO (ppm) minimum	5.0	8.1±0.09 8.3±0.02		8.6±0.03		
			Monsoon Season	1		
Temperature (⁰ C)		27.08 ± 0.05	28.90 ± 0.05	27.20 ± 0.11		
pН	6.5 - 8.5	8.01 ± 0.02	8.02 ± 0.02	7.96 ± 0.03		
TDS (ppm)	1000	198.6 ±5.84	156.5±0.69	161.2±1.36		
DO (ppm) minimum	5.0	6.1±0.09	6.3±0.02	6.8±0.03		

Table 1. Mean Value of seasonal variation in Physico-chemical Parameters of Kampli water

Table 2 Correlation sigificance between selectd physico-chemical variables

		Tank	– I (K1)		Tank – II (K2)			Tank – III (K3)				
	pН	DO	Temp.,	TDS	pН	DO	Temp.,	TDS	pН	DO	Temp.,	TDS
pН	1.00				1.00				1.00			
DO	-0.08	1.00			-0.09	1.00			-0.06	1.00		
Temp.,	-0.48	0.69	1.00		-0.56	0.70	1.00		-0.51	0.71	1.00	
TDS	0.86	-0.72	0.61	1.00	0.79	-0.68	0.70	1.00	0.82	-0.58	0.59	1.00

Note: in bold correlation is significant at the level (2 tailed)

Table 3 Phytoplankton Diversity of selected water tanks in and around the Hosapete city

Locations in and around the Hospet City						
	Kampali Tank - I	Kampali Tank – II	Kampali Tank - III			
Zooplankton diversity	Rotifera	Rotifera	Rotifera			
	Brachionus sp.	Brachionus sp.	Conochilus sp.			
	Conochilus sp.	Conochilus sp.	Polyarthra sp			
	Polyarthra sp	Filinia sp.	Plationus sp			
	Filinia sp.	Plationus sp.	Cladosera			
	Plationus sp	Cladosera	Alonella sp			
	Cladosera	Daphnia sp	Diaphanos oma sp.			
	Daphnia sp	Alonella sp	Copepoda			
	Alonella sp	Moina sp.	Nauplius sp.			
	Diaphanos oma sp.	Copepoda	Heliodiapt omus sp			
	Moina sp.	Cyclops sp.	Ostracoda			
	Copepoda	Heliodiapt omus sp	Cypris sp			
	Cyclops sp.	Ostracoda				
	Nauplius sp.	Cyprinotus sp.				
	Heliodiapt omus sp					
	Ostracoda					
	Cypris sp					
	Cyprinotus sp					



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429

Volume 9 Issue IX Sep 2021- Available at www.ijraset.com

Frequency %	Total count	Rotifera	Cladosera	Copepoda	Ostracoda				
Tank – I (K1)	14 (100.0%)	5 (35.71%)	4(28.057%)	3 (21.43%)	2(14.28%)				
Tank – II (K2)	10 (72.0%)	4 (40.0%)	3 (30.0%)	2 (20.0 %)	1 (10.0 %)				
Tank – III (K3)	08 (57.0%)	3 (37.5%)	2 (25.0%)	2(25.0%)	1 (12.5%)				
	14	5 (35.71%)	3(21.42%)	3(21.42%)	2(14.28%)				

Table 4 Percentage presence of Phytoplankton Diversity in and around the Hosapete city

REFERENCE

- Amphorn Sakset and Wanninee Chankaew. (2013). Phytoplankton as a Bio-indicator of Water Quality in the Freshwater Fishing Area of Pak Phanang River Basin (Southern Thailand), Chiang Mai J. Sci. 2013; 40(3): 344-355
- [2] APHA 2012 Standard Methods for examination of water and wastewater (22nd ed.), 1175 pp. American Public Health Association, Washington DC.
- [3] Arfi R, Bouvy M, Cecchi P, Corbin D, Pagano M. (2003). Environmental conditions and phytoplankton assemblages in two shallow reservoirs of Ivory Coast (West Africa). Archive Fur Hydrobiologie 156: 511-534
- [4] Cecilia Medupin. (2011). Phytoplankton community and their impact on water quality: An analysis of Hollingsworth Lake, UK, J. Appl. Sci. Environ. Manage. Vol. 15 (2) 347 – 350.
- [5] Dhanalakshmi, V., K.Shanthi and K.M.Remia. (2013). Physicochemical study of Eutrophic pond in Pollachi town, Tamilnadu, India. Int.J.Curr.Microbiol.App.Sci. 2(12): 219-227.
- [6] Fokmare, A. K. and M. Musaddiq. (2001) Comparative Studies of Physico-Chemical and Bacteriological Quality of Surface and Ground Water at Akole (MS). Pollution Research. 4(1): 56-61.
- [7] Gadag, S.S., M.S. Kodashettar, N.R. Birasal and M.I. Sambrani. (2005). A checklist of the microphytes and macrophytes i n and around Heggeri lake (Haveri district). Proc. St ate level UGC sponsored semi nar on biodiversity and its conservation held at KLE societ y's Gudl eppa Hallikeri College, Haveri, 28-29. p. 91
- [8] Hulyal S.B. and B.B. Kaliwal. (2011). Seasonal Variations in Physico-Chemical Characteristics of Almatti Reservoir of Bijapur district, Karnataka State. I.J.E.P. 1(1):58-67.
- [9] Kamble BB, Meshram CB (2005) A preliminary study on Zooplankton diversity at Khatijapur tank, near Achlapur, District Amravati, Maharastra. J Aqua Biol. 20(2):45-47.
- [10] Kar S and Kar D (2013) Studies on zooplankton diversity of an oxbow lake of South Assam, India. International Journal of Current Research, 5(12):3652-3655.
- [11] Karr, J.R, J D. Allen, and A. C. Benke. (2000). River conservation in the United States and Canada. In P. J. Boon, Davies and B.R. Petts, G E (Ed.), Global perspectives on River conservation, pp 3–39 Science, Policy, and Practice. Wiley, New York.
- [12] Lacdan, Natividad F., Louise Mae V. Javier, John Vincent A. Pagaddu and Glenn L. Sia Su. (2014). Assessing water quality of Dao River, Batangas using phytoplankton biomonitoring, International Journal of Current Scinece, 12: E 98-102
- [13] Manjare SA (2015) Qualitative and quantitative study of zooplankton from fresh water tanks of kolhapur district, (maharashtra). Research Journal of Life sciences, Bioinformatics, Pharmaceuticals and Chemical Sciences 1(1) 54-58.
- [14] Miah MF, Roy S, Jinnat E and Khan Z K (2013) Assessment of Daphnia, Moina and Cylops in Freshwater Ecosystems and the Evaluation of Mixed Culture in Laboratory. American International Journal of Research in Formal, Applied & Natural Sciences, 4(1): 1-7.
- [15] More, Y.S. and S.N. Nandan. (2000). Hydrobiological study of algae of Panzara river (Maharashtra). Ecol. Environ. Cons., 6, 99-103.
- [16] Mwebaza-Nadwula, M, Sekiranda L, and Kiggundu V. (2005) Variability in zooplankton community along a section of the Upper Victoria Nile, Uganda. Afr. J. Ecol., 43: 251-257.
- [17] Nanda S N and N H Aher. (2005). Algal community used for assessment of water quality of Haranbaree dam and Mosam river of Maharashtra. J. Environ.Biol., 26, 223-227.
- [18] Natividad F. Lacdan, Louise Mae V. Javier, John Vincent A. Pagaddu and Glenn L. Sia Su, (2014). Assessing water quality of Dao River, Batangas using phytoplankton biomonitoring, INT J CURR SCI 2014, 12: E 98-102.
- [19] Niroula B., K.L.B. Singh, G.B. Thapa and J. Pal. (2010). Seasonal Variations in Physico-Chemical Properties and Biodiversity in Betana Pond, Eastern Nepal. Our Nature. 8: 212-218.
- [20] Omar WMW. (2010). Perspectives on the use of algae as biological indicators for monitoring and protecting aquatic environments, with special reference to Malaysian freshwater ecosystems. Tropical Life Sciences Research 21(2): 51-67.
- [21] Pandey BN, Ambasta OP, Thakur AK, Kumar S, Kumar R (2009) Zooplankton diversity in relation to certain physicochemical parameters of swamp of Kishanganj, Bihar. Environ Conserv J. 10(1, 2):9-14.
- [22] Pathak, H., D. Pathak and S. N. Limaye. (2012). Studies on the physico-chemical status of two water bodies at Sagar city under anthropogenic Influences. Advances in Applied Science Research. 3 (1): 31-44.
- [23] Pullie JS, Khan AM (2003). Studies on zooplankton community of Isapur dam water. Ind Poll Res. 22:451-455.
- [24] Radhika, C. G., I. Mini and T. Gangadevi. (2004). Studies on abiotic parameters of a tropical fresh water lake Vellayani Lake, Trivandrum, Kerala. Poll. Res 23(1):49-63.
- [25] Ramulu N. K. and G. Benarjee. (2013). Physico-chemical factors influenced plankton biodiversity and fish abundance- A case study of Andhra Pradesh. Int. J. Lifesc. Bt. & Pharm. Res.1 (2):248-260.
- [26] Shannon CE, Wiener. (1963). The mathematical theory of communications: 117.
- [27] Stevenson, R. J. and Y. Pan. (1999). Assessing environmental conditions in Rivers and streams using diatoms. In E. F. Stoermer and J. P. Smol, (Ed.), The diatoms, Applications for the environmental and earth sciences, pp. 11–40 Cambridge University Press, Cambridge.
- [28] Suresh B. (2015). Multiplicity of phytoplankton diversity in Tungabhadra River near Harihar, Karnataka (India), International Journal of Current Microbiology and Applied Sciences, 4(2): 1077-1085.



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.429 Volume 9 Issue IX Sep 2021- Available at www.ijraset.com

- [29] Tas, Beyhan and Arif Gonulol. (2007). An ecologic and taxonomic study on phytoplankton of a shallow lake, Turkey. J. Envi ron. Bi ol., 28, 439-445.
- [30] Tenenbaum D R, Villac M C, Viana S C, Matos M, Hatherly M, Lima IV, Menezes M. (2004). Phytoplankton atlas of Sepetiba Bay, Rio de Janeiro, Brazil. Globallast Monograph Series No. 16. IMO, London: UK.
- [31] Tiwari, Ashesh and S.V.S. Chauhan. (2006). Seasonal phytoplanktonic diversity of Kitham lake, Agra. J. Environ Biol., 27, 35-38.
- [32] Verma, P.U., A. R. Purohit and N. J. Patel. (2012). Pollution Status of Chandlodia Lake Located in Ahmedabad Gujarat, IJERA. 2:1600-1606.
- [33] Yadav, P., V. K. Yadav, A.K. Yadav and P.K. Khare. (2013). Physico-Chemical Characteristics of a Fresh Water Pond of Orai, U. P., Central India. Octa. J. Biosci. Vol. 1(2): 177-184.
- [34] Zargar, S and T.K. Ghosh. (2006). Influence of cooling water discharges from Kaiga nuclear power plant on selected indices applied to plankton population of Kadra reservoir. J. Environ. Biol., 27, 191-198.











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24*7 Support on Whatsapp)