iJRASET
International Journal For Research in Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 2 **Issue:** VI **Month of publication:** June 2014

DOI:

www.ijraset.com

Call: 08813907089 | E-mail ID: ijraset@gmail.com

ISSN: 2321-9653

INTERNATIONAL JOURNAL FOR RESEARCH IN APPLIED SCIENCE AND ENGINEERING TECHNOLOGY (IJRASET)

Existence of solutions for neutral functional mixed-volterra integrodifferential evolution equations with nonlocal conditions

S. Chandrasekaran[#1] and S. Karunanithi[#2]

#1 *Department of Mathematics, SNS College of Technology, Coimbatore -641 035, Tamil Nadu, India.*
#2 *Department of Mathematics, Kongunadu Arts and Science College, Coimbatore - 641 029, TamilNadu, India.*

Abstract*— *In this work, we study the existence of mild solutions for nonlinear neutral functional mixed-volterra integrodifferential evolution equations with nonlocal conditions in Banach spaces. The results are obtained by using the fractional powers of operators and Sadovskii's fixed point theorem.

Keywords*— *Semigroup, mild and strong solution, Neutral equations, nonlocal condition.
2000 subject Classi fication: 34G20, 47D03, 47H10, 47H20.

I. INTRODUCTION

Abstract neutral differential equations arise in many areas of applied mathematics. As such, they have been largely studied during the last few decades. The literature related to ordinary neutral differential equations is very extensive. The work in partial neutral functional differential equations with infinite delay was initiated by Hernandez and Henriquez. First-order partial neutral functional differential equations have been studied by different authors. The reader can consult Adimy [1], Hale [13, 14] and Wu [25] for systems with finite delay and Hernandez Henriquez [17, 18] and Hernandez [16] for the unbounded delay case. Hernandez [15] established the existence results for partial neutral functional differential equations with nonlocal conditions modeled as

$$\frac{d}{dt}\big(u(t)+F(t,u_t)\big)=Au(t)+G(t,u_t),\quad 0\le t\le T$$
$$u_0=\varphi+q(u_{t_1},u_{t_2},u_{t_3},\ \dots\ u_{t_n})\in\Omega$$

Bahuguna and Agarwal [2] studied the approximation of solution to a partial neutral functional differential equation with nonlocal history condition

$$\frac{d}{dt}(u(t)+g(t,u(t-\tau_1))+Au(t)=f(t,u(t),u(t-\tau_2)),\quad t>0,$$
$$h(u)=\varphi,on\ [-\tau,0]$$

in a separable Hilbert space, where $\tau = max\{\tau_1, \tau_2\}$, τ_1, $\tau_2>0$. An extensive theory for ordinary neutral functional differential equations which includes qualitative behavior of classes of such equations and applications to biological and engineering processes. Several authors have studied the existence of solutions of neutral functional differential equations in Banach space [2, 3, 4, 6, 11, 12, 13, 15, 17, 18, 23]. The nonlocal Cauchy problem for semi linear evolution equations in Banach space was studied first by Byszekswi [7, 8, 9] where he established the existence and uniqueness of mild and classical solutions. The nonlocal conditions were motivated by physical problems and their importance is discussed in [?, ?, ?]. Balachandran et al [2, 4, 5, 21] studied the nonlocal Cauchy problem for various type of nonlinear integrodifferntial equations. In addition, our result can also be

IJRASET: ISSN: 2321-9653

ISSN: 2321-9653

regarded as an extension of the corresponding results on classical problem in [10, 21].

In this paper, we study the following neutral functional integrodifferential equation with nonlocal condition

$$\frac{d}{dt}\Big(x(t)+F\big(t,x(t),x(b_1(t),\dots x(b_n(t))\big)\Big)+A(t)x(t) = G\left(t,x(t),x\big(a_1(t)\big),\dots\dots,x\big(a_m(t)\big)\right) + K(t,x(t),\int_0^t k\big(t,s,x(s)\big)ds\,,\int_0^T \boldsymbol{h}\big(t,s,x(s)\big)ds\,), \quad t \in [0,T]$$

$$x(0)+\ g(x)=x_0 \qquad (1.1)$$

II. PRELIMINARIES

Let $-A$ be the infinitesimal generator of a compact analytic semi group of uniformly bounded linear operators $U(t, s)$ defined in the Banach space X. Let $0 \in \rho(A)$, then define the fractional power A^α, for $0 \le \alpha \le 1$, as a closed linear operator on its domain $D(A^\alpha(t))$ which is dense in X. Further $D(A(t))$ is a Banach space under the norm

$$\|x\|_\alpha = \|A^\alpha x\|\,, x \in D(A^\alpha(t))$$

which we denote by X_α. Then for each $0 \le \alpha \le 1$, $X_\alpha \to X_\beta$ for $0 < \beta < \alpha \le 1$ and the imbedding is compact whenever the resolvent operator of A is compact. We assume that

(a) there is a $M \ge 1$ such that $\|U(t,s)\| \le M$, for all $0 \le t \le a$.

(b) for any $a > 0$, there exists a positive constant C_α such that

$$\|A_\alpha U(t,s)\| \le \frac{C_\alpha}{t^\alpha}, 0 < t \le T.$$

Now we represent the basic assumptions on equation (1.1).

(H_1) $F : [0, T]\times X^{(n+1)} \to X$ is a continuous function, $F(0, T]\times X^{(n+1)} \subset D(A(t))$ with n a positive integer, and there exists constants L, $L_1 > 0$ such that the function $A(t)F$ satisfies the Lipschitz condition:

$$\|A(t)F(s_1,x_0,x_1,\dots,x_n) - A(t)F(s_2,\overline{x_0},\overline{x_1},..,\overline{x_n})\| \le L(|s_1-s_2|+\max_{i=0,1,\dots n}\|x_i-\overline{x_i}\|) \qquad (2.2)$$

for every $0 \le s_1, s_2 \le T$; $x_i, \overline{x_i} \in X$, $i = 0, 1,\dots n$, and the inequality

$$\|A(t)F(t,x_0,x_1,\dots x_n)\| \le L_1(max(\|x_i\| : i = 0, 1, \dots, n) + 1) \qquad (2.3)$$

holds for any $(t, x_0, x_1,\dots x_n) \in [0, T] \times X^{n+1}$.

(H_2) The function $G : [0, T] \times X^{m+1} \to X$ satisfies the following condition:

(**i**) for each $t \in [0, T]$, the function $G(t, .) : X^{m+1} \to X$ is continuous, and for each $(x_0, x_{1,\dots,} x_n) \in X^{m+1}$, the function $G(., x_0, x_{1,\dots,} x_n) : [0; T] \in X$ is strongly measurable.

(**ii**) for each positive constant $k \in N$, there is a positive function $g_k \in L'([0, T])$ such that

$$\sup(\|x_0\|,\dots\dots\|x_m\|) \le k\|G(.,x_0,x_1,\dots\dots\dots,x_m)\| \le g_k\,(t,)$$

and

$$\liminf_{k\to+\infty}\frac{1}{k}\int_o^T g_k\,(s)ds = \gamma < \infty$$

(H_3) The function $K : [0, T] \times X \times X \times X \to X$ satisfies the following condition:

(**i**) For each $t \in [0, T]$, the function $K(.,.,.,.) : X \times X \times X \to X$ and for each $x, y \in X$, $K(.,., x, y) : [0, T] \to X$ is strongly measurable.

(**ii**) For each positive number $r \in N$, there is a positive function $\mu_r \in L'([0, T])$ such that

$$\sup_{\|x\|\le r}\left\|K(s,x(s),\int_0^s k\big(s,\tau,x(\tau)\big)d\tau,\ \int_0^s h\big(s,\tau,x(\tau)\big)d\tau\right\| \le \mu_r(s)$$

and

IJRASET: ISSN: 2321-9653

ISSN: 2321-9653

$$\lim in\ f_{r\to+\infty}\frac{1}{r}\int_o^T \mu_r(s)ds = \gamma_1 < \infty$$

(H_4) $ai, aj \in C([0, T]; [0, T])$, $i = 1, 2,\ldots, m$, $j = 1, 2,\ldots, n$, $g \in C(H;X)$ is completely continuous, where $H = C([0, T];X)$, and there exists a constant $L_2 > 0$ such that $\|g(x)\| \leq L_2\|x\|$ for each $x \in H$.

Theorem 2.1 (Sadovskii's fixed point theorem, [24]).
Let P be a condensing operator on a Banach space X, i,e., P is continuous and takes bounded sets into bounded sets, and α(P(B)) ≤ α (B) for every bounded set B of x with α(B) > 0. If P(H) ⊂ H for convex, closed and bounded set H of X, then P has a bounded point in H (where α(.) denotes the Kuratowski's measures of non-compactness).

III. EXISTENCE OF MILD SOLUTIONS

Definition 3.1: A continuous function $x(.) : [0, T] \to X$ is said to be a mild solution of the nonlocal Cauchy problem(1.1), is the function

$$U(t,s)F\left(s, x(b_1(s)), \ldots\ldots\ldots, x(b_n(s))\right),\quad s \in (0,t)$$

in integrable on [0, t) and the following integral equation is verified:

$$x(t) = U(t,0)\left[x_0 + F\left(0, x(0), x(b_1(0)), \ldots., x(b_n(0))\right) - g(x)\right]$$

$$-F\left(t, x(b_1(t)), \ldots\ldots., x(b_n(t))\right)$$
$$+\int_0^t U(t,s)A(s)F(s, x(s), x(b_1(s)), \ldots\ldots\ldots, x(b_n(s)))ds$$

$$+\int_0^t U(t,s)G(s, x(s), x(a_1(s)), \ldots\ldots\ldots, x(a_m(s)))ds$$
$$+\int_0^t U(t,s)[K(s, x(s), \int_0^t k(s,\tau,x(\tau))d\tau, \int_0^t h(s,\tau,x(\tau))d\tau]ds$$

Theorem 3.1: If the assumption $(H_1) - (H_4)$ are satisfied and $x_0 \in X$, then the nonlocal Cauchy problem (1.1) has a mild solution provided that

$$L0 : L[(M + 1)M_0, MT] < 1$$
and $M_0L_1 + (L_2 + \gamma + \gamma_1 + M_0L_1 + L_1T)M < 1$,

where M is from property (f), $M_0 = \sup \|A^{-1}(t)\|$.

Proof: For the sake of brevity, we rewrite (t, $x(t)$, $x(b_1(t)),\ldots,x(b_n(t))$) = ($t$, $v(t)$) and (t, $x(t)$, $x(a_1(t)),\ldots$, $x(a_m(t))$) = (t, $u(t)$). Define the operator P on $C([0, T];X)$ by the formula

$$(Px)(t) = U(t,0)\left[x_0 + F(0, v(0)) - g(x)\right] - F(t, v(t)) + \int_0^t U(t,s)A(s)F(s, v(s))ds$$
$$+\int_0^t U(t,s)G(s, u(s))ds + \int_0^t U(t,s)K(s, x(s), \int_0^s k(s,\tau,x(\tau))d\tau, \int_0^s h(s,\tau,x(\tau))d\tau)ds,$$
$$0 \leq t \leq T.$$

for each positive number k, let $B_k = \{x \in C([0, T];X) : \|x(t)\| \leq k,\ 0 \leq t \leq T\}$, then for each k, B_k is clearly a nonempty bounded closed convex set in $X([0, T];X)$, since the following relation holds

$$\|U(t,s)A(s)F(s, v(s))\| \leq \|U(t,s)\|\|A(s)F(s, v(s))\| \leq ML_1(k + 1)$$

then from Bouchenr's theorem [20] it follows that $U(t, s)$ $A(s)F(s, v(s))$ is integrable on [0, t] since it is obviously strongly measurable, so P is well defined on B_k. We claim that there exists a positive number k such kthat $P(B_k) \subseteq B_k$. It is not true, then for each positive number k, there is a function $x_k(.) \in B_k$, but $Px_k \notin B_k$, that is

$$\|Px_k(t)\| > k$$

For some $t(k) \in [0, T]$. However, on the other hand, we have

$$k < \|Px_k(t)\| = \|U(t,0)\left[x_0 + F(0, v(0)) - g(x_k)\right] -$$

IJRASET: ISSN: 2321-9653

ISSN: 2321-9653

$$F(t,v_k(t)) + \int_0^t U(t,s)A(s)F(s,v_k(s))ds\| + \left\|\int_0^t U(t,s)\right.$$
$$G(s,u_k(s))ds + \int_0^t U(t,s)K(s,x_k(s),\int_0^s k(s,\tau\, x_k(\tau))d\tau,$$
$$\left.\int_0^T h(s,\tau,x_k(\tau))d\tau\right\|ds$$
$$\leq \|U(t,0)\,[x_0 - g(x_k) + F(0,v_k(0))]\|$$
$$+\|A(t)A^{-1}(t)F(t,v_k(t))\|$$
$$+\int_0^t \|U(t,s)\|\,\|A(s)F(s,v_k(s))\|ds$$
$$\int_0^t \|U(t,s)G(s,u_k(s))\|$$
$$+\int_0^t \left\|U(t,s)\right\|\left\|K(s,x_k(s),\int_0^s k(s,\tau,x_t(\tau))d\tau,\right.$$
$$\left.\int_0^T h(s,\tau,x_k(\tau))d\tau\right\|ds$$
$$\leq M\,[\|x_0\| + L_2k + M_0L_1(k+1)] + M_0L_1(k+1)$$
$$+ ML_1(k+1)T$$
$$+M\int_0^T g_k(s)ds + M\int_0^T \mu_r(s)ds.$$

Dividing on both sides by k and taking the lower limit $k\to +\infty$, we get

$M_0L_1 + (L_2 + M_0L_1 + L_1T + \gamma + \gamma_1)\mathrm{M} \geq 1.$

This is contradicts (7). Hence some positive k, $P(B_k) \leq B_k$.

We will show that the operator P has a fixed point on B_k, which implies that equation (1.1) has a mild solution. To this end, we decompose P into $P = P_1 + P_2$, where the operator P_1, P_2 are dfined on B_k respectively by

$$(P_1x)(t) = U(t,0)F(0,v(0)) - F(t,v(t))$$
$$+\int_0^t U(t,s)A(s)F(s,v(s))ds$$

and

$$(P_2x)(t) = U(t,0)[x_0 - g(x)] + \int_0^t U(t,s)$$
$$G(s,u(s))ds + \int_0^t U(t,s)K[(s,x(s),\int_0^s k(s,\tau,x(\tau))d\tau,$$
$$\int_0^T h(s,\tau,x_k(\tau))d\tau\,]ds$$

$0 \leq t \leq T$, and will verify that P_1 is contraction which P_2 is compact operator.

To prove P_1 is a contraction, we take x_1, $x_2 \in B_k$, then for each $t \in [0, T]$ and by condition (H_1) and (6), we have

$$\|(P_1x_1)(t)(P_2x_2)(t)\| \leq \|U(t,0)[F(0,v_1(0)) -$$
$$F(0,v_2(0))]\| + \|F(t,v_1(t)) - F(t,v_2(t))\|$$
$$+\left\|\int_0^t U(t,s)A(s)[F(s,v_1(s)) - F(s,v_2(s))]ds\right\|$$
$$\leq (M+1)M_0\,L\sup_{0\leq t\leq T}\|x_1(s) - x_2(s)\|$$
$$= L_0\sup_{0\leq t\leq T}\|x_1(s) - x_2(s)\|$$
$$\|(P_1x_1)(t) - (P_2x_2)(t)\| \leq \mathrm{L}_0\|x_1(s) - x_2(s)\|,$$

which shows that P_1 is contraction.

To prove that P_2 is compact, firstly we prove that P_2 is continuous on B_k, Let $\{x_n\} \subseteq B_k$ with $x_n \to x$ is B_k, then by (H_2), we have

$$G(s,u_n(s)) \to G(s,u(s)), n \to \infty$$
$$K\left(t,x_n(t),\int_0^t k(t,s,x_n(s))ds,\int_0^T h(t,s,x_n(s))ds\right)$$
$$\to K\left(t,x(t),\int_0^t k(t,s,x(s))ds,\int_0^T h(t,s,x(s))ds\right)$$

as $n \to \infty$.Since

$$\|G(s,u_n(s)) - G(s,u(s))\| \leq 2g_k(s),$$
$$\left\|K\left(t,x_n(t),\int_0^t k(t,s,x_n(s))ds,\int_0^T h(t,s,x_n(s))ds\right)\right.$$
$$\left.- K\left(t,x(t),\int_0^t k(t,s,x(s))ds,\int_0^T h(t,s,x(s))ds\right)\right\|$$
$$\leq 2\mu_r(s),$$

then by nominated convergence theorem we have,

$$\|P_2\,x_n - P_2\;x\| = \sup_{0\leq t\leq T}\|U(t,0)[x_n(0) - x(0)]$$
$$+\int_0^t U(t,s)[\,G(s,u_n(s)) - G(s,u(s))]ds\|$$

IJRASET: ISSN: 2321-9653

ISSN: 2321-9653

$+ \sup_{0\le t\le T} \left\| \int_0^t U(t,\ s)\,[K\left(s, x_n(s), \int_0^s k(s,\tau,x_n(\tau)\right) d\tau,$
$\int_0^T h\big(s,\tau,x_n(\tau)\big)d\tau)ds -$
$K\left(s, x(s), \int_0^T h\left(s,\tau,x(\tau)\right)d\tau\right) ds]\Big\| \to 0$, as n$\to \infty$.
That is P_2 is continuous.

We prove that the family $\{P_2x : x \in B_{k}\}$ is family of equicontinuous functions. To do this, let $0 \le t_1 \le t2 \le T$; $0 < \epsilon < t_1$, then

$$\|(P_2x)(t_2) - (P_2x)(t_1)\| \le \|U(t_2,0) - U(t_1,0)\|\|x(0)\| + \int_0^{t-\epsilon} \|U(t_2,s) - U(t_1,s)\|\,\|G(s,u(s))\|ds + \int_{t_1-\epsilon}^{t_1} \|U(t_2,s) - U(t_1,s)\|\,\|G(s,u(s))\|ds + \int_{t_1}^{t_2} U(t_2,s)\|\,\|G(s,u(s))\|ds + \int_0^{t_1-\epsilon} \|U(t_2,s) - U(t_1,s)\| \left\|K(s,x(s), \int_0^s k\big(s,\tau,x(\tau)\big)d\tau,\ \int_0^T h\big(s,\tau,x(\tau)\big)d\tau\right\| ds + \int_{t_1-\epsilon}^{t_1} U(t_2,s)\| \left\|K(s,x(s), \int_0^s k\big(s,\tau,x(\tau)\big)d\tau, \int_0^T h\big(s,\tau,x(\tau)\big)d\tau\right\| ds + \int_{t_1}^{t_2} U(t_2,s)\| \left\|K(s,x(s), \int_0^s k\big(s,\tau,x(\tau)\big)d\tau,\ \int_0^T h\big(s,\tau,x(\tau)\big)d\tau\right\| ds$$

Noting that $\|G\big(s,u(s)\big)\| \le g_k(s)$ and $g_k(s) \in L'$, we see that $\|(P_2x)(t_2)-(P_2x)(t_1)\|$ tends to zero independently of $x \in B_k$ as $t_2 - t_1 \to 0$ since the compactness of $\{U(t, s), t > s\}$ implies the continuity of $\{U(t, s), t > s\}$ in t in the uniform operator topology uniformly for s. Hence P_2 maps B_k into a family of equicontinuous functions.

It remains to prove that $V(t) = \{(P_2x)(t) : x \in B_{y}\}$ is relatively compact in X, $V(0)$ is relatively compact in X. Let $0 < t \le T$ be fixed, $0 < \epsilon < t$, for $x \in B_k$, we define

$$(P_2,\epsilon^x)(t) = U(t,0)x(0) + \int_0^{t-\epsilon} U(t,s)G(s,u(s))\,ds + \int_0^{t_{1-\epsilon}} U(t,s)\,K(\,s,x(s),\ \int_0^s k\big(s,\tau,x(\tau)\big)d\tau,\ \int_0^T h\big(s,\tau,x(\tau)\big)d\tau\,)ds$$

$$= U(t,0)x(0) + U(t,t-\epsilon)\int_0^{t-\epsilon} U(t-\epsilon,t)G(s,u(s))\,ds + U(t-\epsilon,s)\int_0^{t_{1-\epsilon}} U(t-\epsilon,s)\,K(\,s,x(s),\ \int_0^s k\big(s,\tau,x(\tau)\big)d\tau,\ \int_0^T h\big(s,\tau,x(\tau)\big)d\tau\,)ds$$

Then from the compactness of $U(t, s)$ $(t - s>0)$, we obtain that

$$V_{\epsilon}(t) = \{(P_{2,}\epsilon^x)(t): x \in B_y\}$$

is relatively compact in X for every, $0<\epsilon< t$. Mortover, $x \in B_y$, we have

$$\|(P_2x)(t) - (P_2,\epsilon^x)(t)\| \le \int_{t-\epsilon}^t \|U(t,s)G(s,u(s))\|ds + \int_{t-\epsilon}^t \|U(t,s)\|\left\|K(s,x(s), \int_0^s k\big(s,\tau,x(\tau)\big)d\tau,\ \int_0^T h(s,\tau,x(\tau)d\tau\right\| \le M\int_{t-\epsilon}^t g_k(s)ds + M\int_{t-\epsilon}^t \mu_r(s)ds$$

IJRASET: ISSN: 2321-9653

ISSN: 2321-9653

Therefore, there are relatively compact sets arbitrarily close to the set $V(t)$. Hence the set $V(t)$ is also relatively compact in X.

Thus by Arzela-Ascoli theorem P_2 is compact operator. These arguments above enable us to conclude that $P = P_1 + P_2$ is condense mapping on B_k, and by Theorem 2.1 there exists a fixed point $z(.)$ for P on B_k, therefore the nonlocal Cauchy problem (1.1) has mild solution. Then the proof is completed.

REFERENCES

[1] M. Adimy and K. Ezzinbi, A class of linear partial neutral functional differential equations with nondense domain, J. Differential Equations, 147(1998) 285 - 332.

[2] D. Bahuguna and S. Agarwal, Approximations of solutions to neutral functional differential equations with nonlocal history conditions, J. Math. Anal. Appl., 317 (2006) 583 - 602.

[3] K. Balachandran and M. Chandrasekaran, Existence of solutions of a de-lay differential equation with nonlocal condition, Indian J.Pure appl. Math., 27(1996,443-449.

[4] K. Balachandran and F. Paul Samuel, Existence of solutions for quasilinear delay integrodifferential equations with nonlocal conditions, Electronic J.Di_erential Eqns., 6 (2009), 1 - 7.

[5] K. Balachandran, R. Sakthivel: Existence of solutions of neutral functional integrodi_erential equation in Banach space, Proc. Indian Acad. Sci.Math.Sci., 109 (1999) 325 - 332.

[6] M. Benchohra and S.K. Ntouyas: Nonlocal Cauchy problems for neutral functional Differential and integrodifferential inclusions, J. Math. Anal. Appl.,258 (2001) 573 - 590.

[7] L. Byszewski and H. Akca, Existence of solutions of semilinear functional differential evolution nonlocal problem, Nonl.Anal.,34(1998),65-72.

[8] L. Byszewski, Theorem about existence and uniqueness of a solution of a semilinear evolution nonlocal Cauchy problem, J.Math.Anal.Appl.,162(1991),496-505.

[9] L. Byszewski, Uniqueness criterion for solution to abstract nonlocal Cauchy problem, J. Appl. Math. Stoch. Anal., 6 (1993), 49-54.

[10] A. Fridman, Partial Differential Equations, Holt, Rienhat and Winston, New York, 169.

[11] X. Fu, On solutions of neutral nonlocal evolution equations with nondense domain, J. Math. Anal. Appl., 299(2004), 392-410.

[12] K. Fu and K. Ezzinbi, Existence of solution for neutral functional evolution equations with nonlocal conditions, Nonl. anal., 54(2003), 215-227.

[13] J.K. Hale, Partial neutral functional-differential equations, Rev. Roumaine Math. Pures Appl., 39 (1994) 339 - 344.

[14] J.K. Hale and M. Verduyn Lunel Sjoerd, Introduction to Functional Differential Equations, Applied Math. Sci., 99. Springer-Verlag, New York, 1993.

[15] E.Hern_andez: Existence results for partial neutral functional differential equations with nonlocal conditions, Cadenos De Mathematica., 02 (2001) 239 - 250.

[16] E. Hern_andez, Existence results for partial neutral integro-differential equations with unbounded delay, J. Math.Anal. Appl., 292 (2004) 194 - 210.

[17] E. Hern_andez and H.R. Henr__quez, Existence of periodic solutions of partial neutral functional-differential equations with unbounded delay, J. Math. Anal.Appl. 221 (2) (1998) 499 - 522.

[18] E. Hern_andez and H.R. Henr__quez, Existence results for partial neutral functional-differential equations with unbounded delay, J. Math. Anal. Appl.,221 (1998) 452 - 475.

[19] Y. Lin and H. Liu, Semilinear integrodifferential equations with nonlocal Cauchy problem, Nonl. anal., 26(1996), 1023-1033.

[20] C.M. Marle, Measure et Probabilities, Herman, Paris, 1974.

[21] F. Paul Samuel and K. Balachandran, Existence results for impulsive quasilinear integrodifferential equations in Banach spaces, Vietnam Journal of Math., 38 (2010), 305 - 321.


[22] A. Pazy, Semigroup of Linear Operators and Applications to Partial Differential Equations, Springer-Verlag, New York, 1983.10

[23] S.M. Rankin III, Existence and asymptotic behavior of a functional differential equations in Banach space, J.Math.Anal.Appl., 88(1982), 531-542.

[24] B.N. Sadovskii, On a fixed point principle, Funct. Anal. Appl., 1(1967), 7476.

[25] J.Wu and H. Xia, Self-sustained oscillations in a ring array of coupled lossless transmission lines, J. Differential Equations, 124 (1996) 247 - 278.

IJRASET: ISSN: 2321-9653



10.22214/IJRASET

45.98

IMPACT FACTOR: 7.129

IMPACT FACTOR: 7.429

INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 (24*7 Support on Whatsapp)