



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 5 Issue: III Month of publication: March 2017 DOI: http://doi.org/10.22214/ijraset.2017.3129

www.ijraset.com

Call: 🛇 08813907089 🕴 E-mail ID: ijraset@gmail.com

www.ijraset.com IC Value: 45.98

International Journal for Research in Applied Science & Engineering Technology (IJRASET) Food Adulteration: A Review

Ameeta Sharma¹, Neha Batra², Anjali Garg³, Ankita Saxena⁴ ^{1,2,3,4}Department of Biotechnology, The IIS University, Jaipur, Rajasthan, India.

Abstract: Food is one of the basic needs for every living being and is very important aspect for life. But now a day's foods are affected by different adulterants. Adulteration is a substance which reduces the vital importance of food. Adulterants like metanil yellow (an artificial color) is used to intensify the color of the turmeric powder also adulterants like chalk powder, brick powder and toxic substances are added to gain profit and to lower the cost so as to compete with the market. Such adulterants can lead to anemia, paralysis, brain damage, stomach disorders and also cancer. Spices may also be contaminated because of conditions under which they are cultivated and harvested. Contaminated spices have been reported to cause certain food-borne illnesses and spoilage.

Keywords: food, adulteration, microbial contamination.

I. INTRODUCTION

Adulteration in food is often present in its most crude form as prohibited substances are either added partially or wholly substituted. Contamination or adulteration in food is added for various reasons which includes financial gain, carelessness and lack in proper hygienic condition of processing, storing, transportation and selling. Therefore, the consumer is either fooled or usually become victim of diseases (Abraham *et al.*, 1997). In backward or developing countries, such kind of adulteration is usually seen. It is essential for the consumer to be aware of common adulterants and their effect on health (Acharya and Shah, 1999). Market has high number of food producers also food is imported in large amount which permits the producer to mislead and cheat consumer. There are producers who take advantage of legal rules and others who commit food adulteration and to differentiate between the two is extremely troublesome. Adulteration in foods not only decreases our social value but morality too (Ankleshwaria and Shah, 1999). The awareness of consumers plays important role in preventing food adulteration. Unawareness and unfair market behaviour might endanger consumer health and misleading can lead to poisoning. Therefore basic screening tests should be known to common people (Vasanthakalaam, 1996).

II. NATURE OF ADULTERATION

Within the past few decades, adulteration of food has become one amongst the intense issues. Many agencies came upon by the Government of India to get rid of adulterants from food stuff. Screening of adulterated and non-adulterated food is important for daily life to make sure that such foods don't cause any health problems (Babu and Shenolikar, 1995). Toxic contaminants are present in ppm level therefore, it's impossible to make sure wholesome food solely on visual examination (Rao, 1994). However, visual examination of the food before purchase bounds to guarantee absence of insects, visual flora, foreign particles etc. In addition, label declaration on packed food is very important in order to know the ingredients and its nutritional value (Babu and Shenolikar, 1995). It also helps in checking the freshness of the food and the period of best before use. The buyer must avoid taking food from a place which avoids hygienic conditions. Food purchased from unhygienic places may cause various diseases (Anyawu and Jukes 1990). Fruits that are being peeled off or cut before selling should not be purchased from unhygienic conditions. It is perpetually better to buy certified food from acknowledged shop.

During different festivals also people use varied composite food colours in excess that cause toxicity (Rao *et al.*, 2005). Before iusing any type of intentional or unintentional adulterant an initial risk analysis should be done (Mathur, 2000) Based on the type of contamination, intention of the producer and processing methodology, adulteration can be divided into 3 types. :

A. Intended Adulterants

It may include adulterants like sand, marble chips, stone, water, coal tar dyes, mud, mineral oil and chalk powder. Such adulterants harm the human body.

B. Metallic Contamination

Metals like arsenic, lead and tin that are present in pesticide, water and cans respectively are responsible for metallic contamination

International Journal for Research in Applied Science & Engineering Technology (IJRASET)

of food products. Such adulterants unintentionally become a part of food during processing (Pappi and Murugesan, 2004).

C. Accidental Adulterants

Accidental adulterants are pesticides residues, dropping of rodents, larvae in foods, etc. Metallic contamination with arsenic lead, mercury can also occur accidentally. Accidental adulterants also involve pests such as rodents and insects that trespasses the food at high degree and produces impurity in the form of excreta, bodily secretions and spoilage through micro-organisms. Most common accidental adulterants are pesticides, D.D.T. and residues present on the plant product (Pandit *et al.*, 2002). Maximum permissible limit for D.D.T. is 3 ppm which sometimes is over crossed (Kannan *et al.*, 1997).

III. IMAPACT OF ADULTERANTS

Now-a-days it is very common to hear or read news about the food items being adulterated and such products are being openly sold out and are consumed by people, which cause various health hazards. News has shown how milk and milk products are being adulterated with urea, soap and other hazardous chemicals (Kalra *et al.*, 1999; Venkateswar, 2001). Also vegetables are being injected to make them grow overnight. In chickens, steroids were being injected to make them into a hen in a very short period of time. Bacteriological profile of street foods in Managlore was also done whose results were in accord (Bhaskar *et al.*, 2004). Adulteration of food leads to several heath issues in humans (Nageswararao *et al.*, 1989; Bhatia *et al.*, 1999). Few health hazards include stomach ache, body ache, anaemia, paralysis, and increase within the incidence of tumors, pathological lesions in very important organs, abnormalities of skin and eyes (Shah *et al.*, 1999). Thus food adulteration ought to be very important because of its impact within the health significance of the public (Beniwal and Khetarpaul, 1999). The individuals are laid low with heart disease, kidney failure, skin diseases, asthma attack and alternative chronic diseases. The individuals are unfortunate victims of this adulteration trade running in full swing and uncurbed.

IV. RELATED CASES

A. Case 1

In Chennai, a commercial shop was inspected for the standard products and the result shows that the products sold were adulterated and it was sold in the name of branded products. About 70% of adulteration was seen in oils and ghee and 10% in masala products (Abhirami and Radha, 2015). Adulerants like metanil yellow (an artificial colour) is used to intensify the colour of the turmeric powder and saw dust are added to increase the quantity of turmeric powder for gaining more profit.

B. Case 2

A study was carried out on the microbiological quality of the spice mix used in the production of Kilishi. Kilishi is a sun dried spiced and grilled meat snack that can be kept for significantly longer duration without getting spoiled because of its dry nature. Application of spice is one among the foremost vital stages throughout the production of kilishi since it has a critical control point (Shamsuddeen and Ameh, 2008). According to International Standard Organization, spices could be defined as the natural vegetable product or mixtures without any extraneous matter that is used for flavoring, seasoning and imparting aroma to foods (ISO, 1972). Spices like other food substances, may carry some bacteria, yeast, mould spores and even some insects. The predominant flora is usually composed of aerobic spores and non spore forming bacteria, indicator organisms and a few pathogens may also be found (ICMSF, 1986). Coliforms from mixed spices were isolated and characterised to be *E. coli, Klebsiella* spp, *Pectobacterium* spp and *Enterobacter* (Patel *et al.*, 1976). According to Frazier and Westhoff (2006), spices don't have a marked bacteriostatic effect in the concentrations used in meat products and they may even serve as source of contamination of processed products.

C. Case 3

This study evaluated the presence of adulterant from daily used food materials like pulses, wheat flour, black pepper, chilli powder and coffee powder which were collected from different departmental and local grocery stores and were checked for the presence of different adulterants through biochemical qualitative analysis. According to the reagents added the colour of the sample changed which indicated the presence of various adulterants (Shruti *et al.*, 2014).

D. Case 4

Another study was done in which 15 packed and 27 unpacked spice samples of three kinds- red pepper, turmeric and coriander were

www.ijraset.com IC Value: 45.98 Volume 5 Issue III, March 2017 ISSN: 2321-9653

International Journal for Research in Applied Science & Engineering Technology (IJRASET)

collected from different markets and bazaars in Bangladesh. Standard microbiological analysis was carried out for the detection and enumeration of microorganisms using standard media. Results were compared with International Microbiological Standard which showed that few microbiological parameters of unpacked sample were higher than the recommended limits (Parveen *et al.*, 2014). Therefore, it's suggested to keep up good sanitary practice and hygienic quality throughout production stages of spices in our country.

V. CONCLUSIONS

From the above review and discussions lastly we can summarize that the adulteration which is been added in the food items can cause tremendous affect on health without our knowledge. Adulteration can be prevented by few alerting steps of our society. Hike of price of food items should be checked by government. While purchasing food items, selection of wholesome and non-adulterated food is necessary to make sure that such food do not cause and health problems. Though presence of adulterants cannot be ensured by visual examination as toxic contaminants are present in very low level but visual examination before purchase can ensure absence of insects, fungus and other foreign materials. The consumer should avoid buying food from places which do not maintain proper hygiene conditions. Both local and branded food stores should be inspected by government bodies. The above general consciousness is simple and easy to initiate for our healthy life. If we tend to actively participate in these changes then we can bring about a healthy and non venturous future for the upcoming generations.

VI. ACKNOWLEDGMENT

We thank IIS University for providing financial grant and necessary facilities to carry out this work.

REFERENCES

- [1] Abhirami, S. and Radha (2015). Detection of food adulteration in selected food items procured by homemaker. International Journal of Recent Scientific Research. 6 (8): pp 5938-5843.
- [2] Abraham, M., Pai, M., Kang, G., Asokan, G.V., Magesh, S.R., Bhattacharji, S. and Rama
- [3] Acharya, M.R. and Shah, R.K. (1999). Some microbiological and chemical attributes of mango pulp samples. Journal of Food Science Technology. 36(4): pp 339-341.
- [4] Ankleshwaria, N. and Shah, S.R. (1999). A study on the awareness pertaining to the use of safe edible food colours amongst Indian Housewives. Applied Nutrition. 24 (2): 21-27.
- [5] Anyawu R. C. and Jukes D. J. (1990), "Food Safety Control Systems for Developing Countries", Food Control, 1(1), 17-26.
- [6] Babu, S. and Shenolikar, I.S. (1995). Health and nutritional implications of food colours. Indian J. of Medical Research. 102: 245-249.
- [7] Beniwal, A. and Khetarpaul, N. (1999) Knowledge of consumers during regarding the nature and extent of adulteration of Indian foods. Nutr Health. 13(3):153-160.
- [8] Bhaskar, J., Usman, M., Smitha, S. and Bhat, G.K. (2004). Bacteriological profile of street foods in Managlore. Indian J. of Medical Microbiology. 22(3): 197.
- [9] Bhatia V, Swami, H.M., Bhatia M. and Bhatia S.P.S. (1999). Attitude and Practice regarding diarrhea in rural community in Chandigarh. Indian Journal Pediatrics. 66 : 499-503.
- [10] Dipak, K. and Dash (2011). National worldwide survey to check food adulteration; Department of food adulteration.
- [11] Frazier, W.C. and Westthroff, W.C. (2006). Food microbiology 3rd Edition, McGraw Hill Publishing Company Limited New York. 163-165, 223-236, 419-543.
- [12] International Commission on microbiological Specification for Food (ICMSF) (1986). Microorganisms in food sampling for microbiological analysis: Principle and specific appilication 2nd edition. Blackwell Scientific Publications 139-140; 213-215.
- [13] International Standards Organisation, (IOS) (1972). Spices and condiments. Women culture Finchizes Draft Proposal. Tc-34/Sc-7, 150 Budapest.
- [14] Kalra, R.L., Kaur, H.Sharma, S., Kapoor, S.K. Chakraborty, S.S. Kshirsagar, R.B. Vaidya, R.C. Sagade, R.B., Shirolkar, S.B., Dikshith, T.S., Raizada, R.B., Srivastava, M.K., Singh, V. Nagaraj, K.V., Appaiah, K.M., Srinivasa, M.A., Rani, M.U., Rao, S.N., Toteja, G.S., Dasgupta, J., Ghosh, P.K. and Saxena, B.N. (1999). DDT and HCH residues in dairy milk samples collected from different geographical regions of India: a multicentric study. Food Add and Contaminants. 16(10): 411-417.
- [15] Kannan, K., Tanabe, S., Gisey, J.P. and Tasukawa, R. (1997). Organochlorine pesticides and polychlorinated biphenyls in foodstuffs from Asian and Oceanic countries. Rev. Environ. Contamin. Toxicol. 152: 1-55
- [16] Krishna B.S. (1997). An outbreak of food poisoning in Tamil Nadu associated with Yersenia enterocolytic. 106: 465-468.
- [17] Nageswara Rao, R., Sudhakar, P. Ramesh V.Bhat and Gupta C.P. (1989). A study of recorded cases of Foodborne diseases at Hyderabad during 1984-1985. J.Trop Med Hyg.1989; 92:320- 324.
- [18] Pandit, G.G., Sharma, S., Srivastava, P.K. and Sahu, S.K. (2002). Persistent organochlorine pesticide residues in milk and dairy products in India. Food Add. and Contaminants. 19(2): 153-157
- [19] Patel, J.D., Krishnaswamy, M.A. and Nair, K.K.S. (1976). Biochemical characteristics of some coliforms isolated from spices. Journal of Food Science Technology, India 13; 37-40.
- [20] Pratima, R., Ramesh, V.B., Sudershan, R.V., T, Prasanna, Krishna. (2005). "Consumption of Synthetic food colors during festivals in Hyderabad, India". British Food Journal. 105 (4&5):276-284
- [21] Praveen, S., Das, S., Begum, A., Sultana, N., Hoque, M.M. and Ahmad, I. (2014). Microbiological quality assessment of three selected spices in Bangladesh.

International Journal for Research in Applied Science & Engineering Technology (IJRASET)

International Food Research Journal 21(4): 1327-1330.

- [22] Pulkith Mathur (2000). Risk Analysis of Food Adulterants. Doctoral Thesis. Delhi University, New Delhi, India.146-183.
- [23] Shah, R.C., Wadher, B.J. and Reddy, B.G.L. (1996). Incidence and characteristics of bacillus cereus isolated from Indian foods. Journal Institute of Food Science Technology. 33(3): 249- 250.
- [24] Shamusuddeen, U. and Ameh, J.B. (2008). Survey on the possible critical control points in kilishi (a traditional dried and grilled meat snack) produced in kano. International Journal of Bioscience, 3(2): 34-38.
- [25] Shruti, A., Kirti, J., Anwesha, D., Raza, A., Ganesh, S. and Kishan, N. (2014). Analysis of food quality and food adulterants from different departmental & local grocery stores by qualitative analysis for food safety. IOSR Journal Of Environmental Science, Toxicology And Food Technology (IOSR-JESTFT) e-ISSN: 2319-2402,p- ISSN: 2319-2399.Volume 8, Issue 2 Ver. III (Mar-Apr. 2014), PP 22-26.
- [26] Sudershan Rao, V. (1994). Analytical and Toxicological studies on selected newer food adulterants. Doctoral thesis, Mangalore University, Karnataka, India. 126-130.
- [27] Sumar, S. and Ismail, H. (1995). Adulteration of food- past and present. Nutr. Fd. Sci. 4 : 11-15.
- [28] Vasanthakalaam, H. (1996). Studies on food handling and microbiological quality of street foods in Madurai city. Ph.D. Thesis. Dept. of Food Science and Nutrition, Agricultural and Research Institute. TNAU.
- [29] Venkateswar Rao, L. (2001). Quality of milk and milk product marketed in Hyderabad city with special reference to public health. Ph.D. Thesis. A.N.G.R. Agricultural University, Hyderabad, India.











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24*7 Support on Whatsapp)