



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 5

Issue: IV

Month of publication: April 2017

DOI:

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

Operations on Hesitant Fuzzy Hypergraph

Dr. A Thamilsai¹, R Rajeswari²

¹Assistant Professor, Kamaraj College of Engineering, Virudhunagar

²Assistant Professor, Department of Mathematics, Fatima College, Madurai.

Abstract : The concept of hypergraphs was extended to fuzzy hypergraph. In this paper, we extend the concepts of fuzzy hypergraphs into that of hesitant fuzzy hypergraphs. Based on the definition of hesitant fuzzy graph, order of the hfhg, size of the hfhg, operations like complement, join, union, intersection, ringsum, cartesian product, composition are defined for hesitant fuzzy graphs. The authors further proposed to apply these operations in clustering techniques.

I. INTRODUCTION

Hypergraph theory, originally developed by C.Berge in 1960, is a generalization of graph theory. The concept of hypergraphs can model more general types of relations than binary relations. The notion of hypergraphs has been extended in fuzzy theory and the concept of fuzzy hypergraphs was proposed by Lee-Kwang and S.M.Chen. The authors have already introduced the concept of Hesitant fuzzy hypergraph. Operations on HFHGs have also been analyzed to a real-life problem with a numerical example.

II. PRELIMINARIES

A. Definition 1.1

Let X be a fixed set, a Hesitant fuzzy set (HFS) on X is in terms of a function that when applied to X returns a subset of $[0, 1]$. The HFS is defined by a mathematical symbol:

$$A = \{ \langle x, h_A(x) \rangle / x \in X \}$$

where $h_A(x)$ is a set of some values in $[0, 1]$, denoting the possible membership degrees of the element $x \in X$ to the set A . Xu and Xia (2011b) called $h = h_A(x)$ a hesitant fuzzy element (HFE) and Θ the set of all HFEs.

B. Definition 1.2

A Hesitancy Fuzzy Graph is of the form $G = (V, E)$, where

$V = \{v_1, v_2, \dots, v_n\}$ such that $\mu_1: V \rightarrow [0, 1]$, $\gamma_1: V \rightarrow [0, 1]$ and $\beta_1: V \rightarrow [0, 1]$ denote the degree of membership, non-membership and hesitancy of the element $v_i \in V$ respectively and $\mu_1(v_i) + \gamma_1(v_i) + \beta_1(v_i) = 1$ for every $v_i \in V$, where $\beta_1(v_i) = 1 - [\mu_1(v_i) + \gamma_1(v_i)]$ and $E \subseteq V \times V$ where $\mu_2: V \times V \rightarrow [0, 1]$, $\gamma_2: V \times V \rightarrow [0, 1]$ and $\beta_2: V \times V \rightarrow [0, 1]$ are such that,

$$\mu_2(v_i, v_j) \leq \min [\mu_1(v_i), \mu_1(v_j)]$$

$$\gamma_2(v_i, v_j) \leq \max [\gamma_1(v_i), \gamma_1(v_j)]$$

$$\beta_2(v_i, v_j) \leq \min [\beta_1(v_i), \beta_1(v_j)]$$

$$\text{and } 0 \leq \mu_2(v_i, v_j) + \gamma_2(v_i, v_j) + \beta_2(v_i, v_j) \leq 1 \text{ for every } (v_i, v_j) \in E.$$

C. Definition 1.3

1) Hypergraph H is an ordered pair $H = (V, E)$ where

a) $V = \{v_1, v_2, v_3 \dots v_n\}$, a finite set of vertices.

b) $E = \{E_1, E_2, E_3 \dots E_m\}$, a family of subsets of V .

c) $E_j \neq \emptyset, j = 1, 2, \dots, m$ and $\cup E_j = V$.

The set V is called the set of vertices and E is the set of edges (or hyper edges).

D. Definition 1.4

In a hypergraph, two or more vertices $x_1, x_2, \dots, x_n$ are said to be adjacent if there exist an edge E_j which contains those vertices. In a hypergraph two edges E_i & $E_j, i \neq j$ is said to be adjacent if their intersection is not empty.

E. Definition 1.5

A regular hyper graph is one in which every vertex is contained in k edges for some constant k .

International Journal for Research in Applied Science & Engineering Technology (IJRASET)

F. Definition 1.6

1) Hesitant Fuzzy HyperGraph (HFHG) is an ordered pair $H = \langle V, E \rangle$ where

- a) $V = \{v_1, v_2, v_3 \dots v_n\}$, a finite set of vertices
- b) $E = \{E_1, E_2, E_3 \dots E_m\}$, a family of Hesitant fuzzy subsets of V .
- c) $E_j = \{ \langle v_i, \mu_j(v_i), \gamma_j(v_i), \beta_j(v_i) \rangle / \mu_j(v_i), \gamma_j(v_i), \beta_j(v_i) \geq 0 \text{ and } 0 \leq \mu_j(v_i, v_j) + \gamma_j(v_i, v_j) + \beta_j(v_i, v_j) \leq 1, j = 1, 2, \dots, m \}$
- d) $E_j \neq \emptyset, j = 1, 2, \dots, m$

Here the edges E_j are HFHGs. $\mu_j(v_i), \gamma_j(v_i), \beta_j(v_i)$ denote the degree of membership, non-membership and hesitancy of the element $v_i \in V$ respectively of the vertex v_i to edge E_j .

G. Definition 1.7

Let $H = (X, E)$ be a Regular Hesitant Fuzzy Hypergraph. The **order** of a Regular Hesitant Fuzzy Hypergraph, H in

$$O(H) = \left(\sum_{x \in X} \mu_{E_i}(x), \sum_{x \in X} \gamma_{E_i}(x), \sum_{x \in X} \beta_{E_i}(x) \right) \text{ for every } x \in X.$$

The **size** of Regular Hesitant Fuzzy Hypergraph is

$$S(H) = \sum_{i=1}^n S(E_i) \text{ where } S(E_i) = \left(\sum_{x \in E_i} \mu_{E_i}(x), \sum_{x \in E_i} \gamma_{E_i}(x), \sum_{x \in E_i} \beta_{E_i}(x) \right)$$

1) *Example 1:* Consider the Hesitant Fuzzy Hypergraph $H = (X, E)$, Define $X = \{a, b, c, d\}$ and

$E = \{E_1, E_2, E_3, E_4\}$, where

$$E_1 = \{(a, 0.1, 0.7, 0.2) (b, 0.1, 0.7, 0.2)\}$$

$$E_2 = \{(b, 0.1, 0.7, 0.2) (c, 0.1, 0.7, 0.2)\}$$

$$E_3 = \{(c, 0.1, 0.7, 0.2) (d, 0.1, 0.7, 0.2)\}$$

$$E_4 = \{(d, 0.1, 0.7, 0.2) (a, 0.1, 0.7, 0.2)\}$$

Here, Order of the HFHG, $O(H) = (0.4, 2.8, 0.8)$,

Size of the HFHG, $S(H) = (0.8, 5.6, 1.6)$.

III. OPERATIONS ON HESITANT FUZZY HYPERGRAPHS

A. Definition 2.1

Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be two HFHGs. Then the **Union** of G_1 and G_2 is an HFHG $G = G_1 \cup G_2 = (V_1 \cup V_2, E_1 \cup E_2)$ defined by

$$(\mu_1 \cup \mu_1')(v) = \begin{cases} \mu_1(v) & \text{if } v \in V_1 - V_2 \\ \mu_1'(v) & \text{if } v \in V_2 - V_1 \end{cases}$$

$$(\gamma_1 \cup \gamma_1')(v) = \begin{cases} \gamma_1(v) & \text{if } v \in V_1 - V_2 \\ \gamma_1'(v) & \text{if } v \in V_2 - V_1 \end{cases}$$

$$(\beta_1 \cup \beta_1')(v) = \begin{cases} \beta_1(v) & \text{if } v \in V_1 - V_2 \\ \beta_1'(v) & \text{if } v \in V_2 - V_1 \end{cases}$$

$$\text{And } (\mu_2 \cup \mu_2')(v_i v_j) = \begin{cases} \mu_{2ij} & \text{if } e_{ij} \in E_1 - E_2 \\ \mu_{2ij}' & \text{if } e_{ij} \in E_2 - E_1 \end{cases}$$


International Journal for Research in Applied Science & Engineering Technology (IJRASET)

$$(\gamma_2 \cup \gamma'_2)(v_i v_j) = \begin{cases} \gamma_{2ij} & \text{if } e_{ij} \in E_1 - E_2 \\ \gamma'_{2ij} & \text{if } e_{ij} \in E_2 - E_1 \end{cases}$$

$$(\beta_2 \cup \beta'_2)(v_i v_j) = \begin{cases} \beta_{2ij} & \text{if } e_{ij} \in E_1 - E_2 \\ \beta'_{2ij} & \text{if } e_{ij} \in E_2 - E_1 \end{cases}$$

where $(\mu_1, \gamma_1, \beta_1)$ and $(\mu'_1, \gamma'_1, \beta'_1)$ refer the vertex membership, non-membership and hesitant element of G_1 and G_2 respectively; $(\mu_2, \gamma_2, \beta_2)$ and $(\mu'_2, \gamma'_2, \beta'_2)$ refer the edge membership, non-membership and hesitant element of G_1 and G_2 respectively.

1) *Example 2:* Consider the Hesitant Fuzzy Hypergraphs $G_1 = (V_1, E_1)$ where $V_1 = \{a, b, c, d\}$ and $G_2 = (V_2, E_2)$ where $V_2 = \{u_1, u_2\}$.



B. Definition 2.2

Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be two HFHG. Then the **Join** of G_1 and G_2 is an HFHG $G = G_1 + G_2 = (V_1 \cup V_2, E_1 \cup E_2)$ defined by

$$(\mu_1 + \mu'_1)(v) = (\mu_1 \cup \mu'_1)(v) \quad \text{if } v \in V_1 \cup V_2$$

$$(\gamma_1 + \gamma'_1)(v) = (\gamma_1 \cup \gamma'_1)(v) \quad \text{if } v \in V_1 \cup V_2$$


$$(\beta_1 + \beta'_1)(v) = (\beta_1 \cup \beta'_1)(v) \quad \text{if } v \in V_1 \cup V_2$$

International Journal for Research in Applied Science & Engineering Technology (IJRASET)

$$(\mu_1 + \mu'_1)(v_i v_j) = \begin{cases} (\mu_1 + \mu'_1)(v_i v_j) & \text{if } v_i v_j \in E_1 \cup E_2 \\ \mu_1(v_i) \mu'_1(v_j) & \text{if } v_i v_j \in E' \end{cases}$$

$$(\gamma_1 + \gamma'_1)(v_i v_j) = \begin{cases} (\gamma_1 + \gamma'_1)(v_i v_j) & \text{if } v_i v_j \in E_1 \cup E_2 \\ \gamma_1(v_i) \gamma'_1(v_j) & \text{if } v_i v_j \in E' \end{cases} \quad (\beta_1 + \beta'_1)(v_i v_j) = \begin{cases} (\beta_1 + \beta'_1)(v_i v_j) & \text{if } v_i v_j \in E_1 \cup E_2 \\ \beta_1(v_i) \beta'_1(v_j) & \text{if } v_i v_j \in E' \end{cases}$$

1) *Example 3:* Consider the Hesitant Fuzzy Hypergraphs $G_1 = (V_1, E_1)$ where $V_1 = \{a, b, c, d\}$ and $G_2 = (V_2, E_2)$ where $V_2 = \{u_1, u_2\}$.



C. Definition 2.3

Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be two HFHG. Then the **Intersection** of HFHGs G_1 and G_2 , denoted by $G_1 \cap G_2$, is an HFHG defined by

$$(\mu_1 \cap \mu'_1)(v) = \begin{cases} \mu_1(v) & \text{if } v \in V_1 \text{ and } V_2 \\ \mu'_1(v) & \text{if } v \in V_2 \text{ and } V_1 \end{cases}$$

$$(\gamma_1 \cap \gamma'_1)(v) = \begin{cases} \gamma_1(v) & \text{if } v \in V_1 \text{ and } V_2 \\ \gamma'_1(v) & \text{if } v \in V_2 \text{ and } V_1 \end{cases}$$

$$(\beta_1 \cap \beta'_1)(v) = \begin{cases} \beta_1(v) & \text{if } v \in V_1 \text{ and } V_2 \\ \beta'_1(v) & \text{if } v \in V_2 \text{ and } V_1 \end{cases}$$


$$(\mu_2 \cap \mu'_2)(v_i v_j) = \begin{cases} \mu_{2ij} & \text{if } e_{ij} \in E_1 \text{ and } E_2 \\ \mu'_{2ij} & \text{if } e_{ij} \in E_2 \text{ and } E_1 \end{cases}$$

International Journal for Research in Applied Science & Engineering Technology (IJRASET)

$$(\gamma_2 \text{ I } \gamma'_2)(v_i v_j) = \begin{cases} \gamma_{2ij} & \text{if } e_{ij} \in E_1 \text{ and } E_2 \\ \gamma'_{2ij} & \text{if } e_{ij} \in E_2 \text{ and } E_1 \end{cases}$$

$$(\beta_2 \text{ I } \beta'_2)(v_i v_j) = \begin{cases} \beta_{2ij} & \text{if } e_{ij} \in E_1 \text{ and } E_2 \\ \beta'_{2ij} & \text{if } e_{ij} \in E_2 \text{ and } E_1 \end{cases}$$


1) Example 4:




D. Definition 2.4

Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be two HFHGs. Then the ringsum of HFHGs G_1 and G_2 is an HFHG defined by

$$G_1 \oplus G_2 = (G_1 \cup G_2) - (G_1 \text{ I } G_2).$$



International Journal for Research in Applied Science & Engineering Technology (IJRASET)



E. Definition 2.5

Let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be two HFHGs. Then the Cartesian product of HFHGs G_1 and G_2 is an HFHG, denoted by $G = G_1 \times G_2$, defined by $G = (V, E)$ $V = V_1 \times V_2$ and

$$E = \{(u, u_2)(u, v_2) : u \in V_1, u_2 v_2 \in E_2\} \cup \{(u_1, w)(v_1, w) : w \in V_2, u_1 v_1 \in E_1\}$$

where

(i)

$$(\mu_1 \times \mu_1')(u_1, u_2) = \mu_1(u_1) \cdot \mu_1'(u_2) \text{ for every } (u_1, u_2) \in V$$

$$(\gamma_1 \times \gamma_1')(u_1, u_2) = \gamma_1(u_1) \cdot \gamma_1'(u_2) \text{ for every } (u_1, u_2) \in V \text{ and}$$

$$(\beta_1 \times \beta_1')(u_1, u_2) = \beta_1(u_1) \cdot \beta_1'(u_2) \text{ for every } (u_1, u_2) \in V$$

(ii)

$$(\mu_2 \times \mu_2')(u, u_2)(u, v_2) = \mu_1(u) \cdot \mu_2(u_2 v_2) \text{ for every } u \in V_1, u_2 v_2 \in E_2$$

$$(\gamma_2 \times \gamma_2')(u, u_2)(u, v_2) = \gamma_1(u) \cdot \gamma_2(u_2 v_2) \text{ for every } u \in V_1, u_2 v_2 \in E_2$$


$$(\beta_2 \times \beta_2')(u, u_2)(u, v_2) = \beta_1(u) \cdot \beta_2(u_2 v_2) \text{ for every } u \in V_1, u_2 v_2 \in E_2 \text{ and}$$

$$(\mu_2 \times \mu_2')(u_1, w)(v_1, w) = \mu_1(w) \cdot \mu_2(u_1 v_1) \text{ for every } w \in V_2, u_1 v_1 \in E_1$$

$$(\gamma_2 \times \gamma_2')(u_1, w)(v_1, w) = \gamma_1(w) \cdot \gamma_2(u_1 v_1) \text{ for every } w \in V_2, u_1 v_1 \in E_1$$

$$(\beta_2 \times \beta_2')(u_1, w)(v_1, w) = \beta_1(w) \cdot \beta_2(u_1 v_1) \text{ for every } w \in V_2, u_1 v_1 \in E_1$$

1) Example 5:



F. Definition 2.6

Let $G = G_1 \circ G_2 = (V_1 \times V_2, E)$ be the Composition of two graphs G_1 and G_2 where $V = V_1 \times V_2$ and

$$E = \{(u, u_2)(u, v_2) : u \in V_1, u_2 v_2 \in E_2\} \cup \{(u_1, w)(v_1, w) : w \in V_2, u_1 v_1 \in E_1\}$$

$$\cup \{(u_1, u_2)(v_1, v_2) : u_1 v_1 \in E_1, u_2 \neq v_2\}$$

Then, the “Composition of HFHGs G_1 and G_2 ”, denoted by $G = G_1 \circ G_2$, is an HFHG defined by

(i)

$$(\mu_1 \circ \mu_1')(u_1, u_2) = \mu_1(u_1) \cdot \mu_1'(u_2) \text{ for every } (u_1, u_2) \in V_1 \times V_2$$

$$(\gamma_1 \circ \gamma_1')(u_1, u_2) = \gamma_1(u_1) \cdot \gamma_1'(u_2) \text{ for every } (u_1, u_2) \in V_1 \times V_2 \text{ and}$$

International Journal for Research in Applied Science & Engineering Technology (IJRASET)

$$(\beta_1 \circ \beta_1')(u_1, u_2) = \beta_1(u_1) \cdot \beta_1'(u_2) \text{ for every } (u_1, u_2) \in V_1 \times V_2$$

(ii)

$$(\mu_2 \circ \mu_2')(u, u_2)(u, v_2) = \mu_1(u) \cdot \mu_2(u_2 v_2) \text{ for every } u \in V_1, u_2 v_2 \in E_2$$

$$(\gamma_2 \circ \gamma_2')(u, u_2)(u, v_2) = \gamma_1(u) \cdot \gamma_2(u_2 v_2) \text{ for every } u \in V_1, u_2 v_2 \in E_2$$

$$(\beta_2 \circ \beta_2')(u, u_2)(u, v_2) = \beta_1(u) \cdot \beta_2(u_2 v_2) \text{ for every } u \in V_1, u_2 v_2 \in E_2$$

$$(\mu_2 \circ \mu_2')(u_1, w)(v_1, w) = \mu_1(w) \cdot \mu_2(u_1 v_1) \text{ for every } w \in V_2, u_1 v_1 \in E_1$$

$$(\gamma_2 \circ \gamma_2')(u_1, w)(v_1, w) = \gamma_1(w) \cdot \gamma_2(u_1 v_1) \text{ for every } w \in V_2, u_1 v_1 \in E_1$$

$$(\beta_2 \circ \beta_2')(u_1, w)(v_1, w) = \beta_1(w) \cdot \beta_2(u_1 v_1) \text{ for every } w \in V_2, u_1 v_1 \in E_1 \text{ and}$$

$$(\mu_2 \circ \mu_2')(u_1, u_2)(v_1, v_2) = \mu_1'(u_2) \cdot \mu_1'(v_2) \cdot \mu_2(u_1 v_1) \text{ for every } (u_1, u_2), (v_1, v_2) \in E - E''$$


$$(\gamma_2 \circ \gamma_2')(u_1, u_2)(v_1, v_2) = \gamma_1'(u_2) \cdot \gamma_1'(v_2) \cdot \gamma_2(u_1 v_1) \text{ for every } (u_1, u_2), (v_1, v_2) \in E - E''$$

$$(\beta_2 \circ \beta_2')(u_1, u_2)(v_1, v_2) = \beta_1'(u_2) \cdot \beta_1'(v_2) \cdot \beta_2(u_1 v_1) \text{ for every } (u_1, u_2), (v_1, v_2) \in E - E''$$

where

$$E'' = \{(u, u_2)(u, v_2) : u \in V_1, u_2 v_2 \in E_2\} \cup \{(u_1, w)(v_1, w) : w \in V_2, u_1 v_1 \in E_1\}$$

1) Example 6:



IV. CONCLUSION

In this paper, order of the HFHG, size of the HFHG, Operations like complement, join, union, intersection, ringsum, Cartesian product, composition on HFHGs are defined. Currently, the authors are working on existing clustering techniques. Further, it is proposed to apply the properties of HFHGs to develop a new clustering algorithm and the same may be checked with a numerical dataset.

REFERENCES

- [1] Z. Xu, Hesitant Fuzzy Sets Theory, Studies in Fuzziness and Soft Computing 314, 1 DOI: 10.1007/978-3-319-04711-9_1, © Springer International Publishing Switzerland 2014.
- [2] Mordeson N. John, Nair S. Premchand, Fuzzy Graphs and Fuzzy Hypergraphs, New York, Physica-Verlag, 2000, ISBN 3-7908-1286-2.
- [3] T. Pathinathan, J. Jon Arockiaraj and J. Jesintha Rosline, Hesitancy Fuzzy Graphs, Vol 8(35), DOI: 10.17485/ijst/2015/v8i35/86672, December 2015, Indian Journal of Science and Technology, ISSN (Print) : 0974-6846.
- [4] R.Parvathi, M.G.Karunambigai and K.T.Atanassov, Operations on Intuitionistic fuzzy graphs, Fuzzy Systems, 2009, Fuzz-IEEE 2009. IEEE international conference 1396-1401.
- [5] S.Thilagavathi, R.Parvathi and M.G.Karunambigai, Intuitionistic fuzzy hypergraphs, J.Cybernetics and Information Technologies 9(2), (2009), 46-53, Sofia.
- [6] I.Pradeepa and S.Vimala, Regular and Totally Regular Intuitionistic Fuzzy Hypergraph (IFH), Volume 4, Issue 1-C (2016), 137-142, Indian Journal of Mathematics And its Applications, ISSN (Print) : 2347-1557.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)