



# IJRASET

International Journal For Research in  
Applied Science and Engineering Technology



---

# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

---

**Volume: 5      Issue: VII      Month of publication: July 2017**

**DOI:**

**[www.ijraset.com](http://www.ijraset.com)**

**Call:  08813907089**

**E-mail ID: [ijraset@gmail.com](mailto:ijraset@gmail.com)**

# On Reducibility of Certain q-Double Hypergeometric Series and Clausen Type Identities

Rajesh Pandey

Department of Applied Science, Institute of Engineering & Technology, Sitapur Road, Lucknow 226021 India.

**Abstract:** In this paper, we have made use of certain known summations to establish transformations of q-double series in terms of single series. We have deduced Clausen type identities from these results.

**Keywords :** Hypergeometric functions, Summations, Transformation, Identities and Convergence.

## I. INTRODUCTION

For  $\alpha$ , real or complex and  $|q| < 1$ , we define the q-shifted factorials by

$$[\alpha; q]_n = \begin{cases} 1 & \text{if } n = 0 \\ (1 - \alpha)(1 - \alpha q) \dots (1 - \alpha q^{n-1}), & \text{if } n = 1, 2, 3, \dots \end{cases} \quad (1.1)$$

A basic hypergeometric function is defined as:

$${}_r\Phi_s \left[ \begin{matrix} a_1, a_2, \dots, a_r; q; z \\ b_1, b_2, \dots, b_s; q^\lambda \end{matrix} \right] = \sum_{n=0}^{\infty} \frac{[a_1, a_2, \dots, a_r; q]_n z^n q^{\lambda n(n-1)/2}}{[q, b_1, b_2, \dots, b_s; q]_n}, \quad (1.2)$$

Where  $[a_1, a_2, \dots, a_r; q]_n = [a_1; q]_n [a_2; q]_n \dots [a_r; q]_n$ .

The series  ${}_r\Phi_s$  converges absolutely for all z if  $\lambda > 0$  and for  $|z| < 1$  if  $\lambda = 0$ . we shall use the following series identity to establish our results.

$$\sum_{n=0}^{\infty} \sum_{k=0}^n B(n, k) = \sum_{n,k=0}^{\infty} B(n+k, k), \quad (1.3)$$

Provided the series on both sides of (1.3) exist.

## II. NOTATIONS AND DEFINITIONS

Notations and definitions appearing in this paper have their usual meaning. We shall use the following known summations of q-series in our analysis:

$${}_2\Phi_1 \left[ \begin{matrix} q^{-n}, a; q; zq^n/a \\ c \end{matrix} \right] = \frac{[c/a; q]_n}{[c; q]_n}. \quad (2.1)$$

$${}_3\Phi_2 \left[ \begin{matrix} a, b, q^{-n}; q; q \\ c, abq^{1-n}/c \end{matrix} \right] = \frac{[c/a, c/b; q]_n}{[c, c/ab; q]_n} \quad (2.2)$$

$${}_2\Phi_1 \left[ \begin{matrix} x, q^{-n}; q; -q/x \\ q^{-n}/x \end{matrix} \right] = \frac{[q; q]_n [x^2 q^2; q^2]_m}{[xq; q]_n [q^2; q^2]_m}, \quad (2.3)$$

Where m is the greatest integer  $\leq n/2$ .

$${}_4\Phi_3 \left[ \begin{matrix} q^{-n}, -q^{-n}/xy, x, y; q; q \\ -xyq, q^{-n}/x, q^{-n}/y \end{matrix} \right] = \frac{[q, xyq; q]_n [x^2 q^2, y^2 q^2; q^2]_m}{[xq, yq; q]_n [q^2, x^2 y^2 q^2; q^2]_m}, \quad (2.4)$$

Where m is the greatest integer  $\leq n/2$

$${}_2\Phi_1 \left[ \begin{matrix} x, q^{-n}; q; -1/x \\ q^{-n}/x \end{matrix} \right] = \frac{[q; q]_n [x^2 q^2; q^2]_m x^{n-2m}}{[xq; q]_n [q^2; q^2]_m}, \quad (2.5)$$

Where m is the greatest integer  $\leq n/2$ .

$$\begin{aligned}
 & {}_4\Phi_3 \left[ \begin{matrix} q^{-n}, -q^{-n}/xy, xq, yq \\ -xyq, q^{1-n}/x, q^{1-n}/y \end{matrix}; q; q \right] \\
 &= \frac{(-)^n [q; q]_n [xyq; q]_n [x^2q^2, y^2q^2; q^2]_m}{q^n [x, y; q]_n [q^2, x^2y^2q^2; q^2]_m}, \tag{2.6}
 \end{aligned}$$

Where m is the greatest integer  $\leq n/2$ .

$$\begin{aligned}
 & {}_4\Phi_3 \left[ \begin{matrix} q^{-n}, -q^{-n}/x^2, y, -y \\ q^{-n}/x, -q^{-n}/x, y^2q \end{matrix}; q; q \right] \\
 &= \frac{[q; q]_n [x^2y^2q^2; q^2]_n [x^2q^2, y^2q^2; q^2]_m}{[x^2q^2; q^2]_n [y^2q; q]_n [q^2, x^2y^2q^2; q^2]_m}, \tag{2.7}
 \end{aligned}$$

Where m is the greatest integer  $\leq n/2$ .

$${}_3\Phi_2 \left[ \begin{matrix} q^{-n}, q^{-n}/x^2, 0 \\ q^{-n}/x, -q^{-n}/x \end{matrix}; q; q \right] = \frac{[q; q]_n [x^2q^2; q^2]_m}{[x^2y^2; q^2]_n [q^2; q^2]_m}, \tag{2.8}$$

Where m is the greatest integer  $\leq n/2$ .

$$\begin{aligned}
 & {}_3\Phi_2 \left[ \begin{matrix} q^{-n}, q^{-n}/x^2, 0 \\ q^{-n}/x, -q^{-n}/x; q \end{matrix}; 1 \right] \\
 &= \frac{[q; q]_n [x^2q^2; q^2]_m q^{n(n+1)/2} x^{2n-2m}}{[x^2y^2; q^2]_n [q^2; q^2]_m}, \tag{2.9}
 \end{aligned}$$

Where m is the greatest integer  $\leq n/2$ .

$$\begin{aligned}
 & {}_4\Phi_3 \left[ \begin{matrix} q^{-n}, q^{-n}/x^2, yq, -yq \\ q^{1-n}/x, q^{1-n}/x, y^2q \end{matrix}; q; q \right] \\
 &= \frac{(-)^n [q; q]_n [x^2y^2q^2; q^2]_n [x^2q^2, y^2q^2; q^2]_m}{q^2 [x^2; q^2]_n [y^2q; q]_n [q^2, x^2y^2q^2; q^2]_m}, \tag{2.10}
 \end{aligned}$$

where m is the greatest integer  $\leq n/2$ .

$${}_3\Phi_2 \left[ \begin{matrix} q^{-n}, q^{-n}/x^2, 0 \\ q^{1-n}/x, q^{1-n}/x \end{matrix}; q; q \right] = \frac{(-)^n [q; q]_n [x^2q^2; q^2]_m}{q^n [x^2; q^2]_n [q^2; q^2]_m}, \tag{2.11}$$

Where m is the greatest integer  $\leq n/2$ .

$$\begin{aligned}
 & {}_3\Phi_2 \left[ \begin{matrix} q^{-n}, q^{-n}/x^2, 0 \\ q^{1-n}/x, q^{1-n}/x; q \end{matrix}; q^2 \right] \\
 &= \frac{(-)^n [q; q]_n [x^2q^2; q^2]_m q^{n(n-1)/2} x^{2n-2m}}{[x^2; q^2]_n [q^2; q^2]_m}, \tag{2.12}
 \end{aligned}$$

Where m is the greatest integer  $\leq n/2$ .

Putting  $yq^{-n}$  for y in [Verma and Jain 1; (2.20) P.1027] we get the following summation formula:

$$\begin{aligned}
 & {}_4\Phi_3 \left[ \begin{matrix} q^{-n}, q^{-n}/x^2, 1/xy, -1/xy \\ q^{1-n}/x, -q^{-n}/x, 1/x^2y^2 \end{matrix}; q; q \right] \\
 &= \frac{(-)^n (xq)^{-n} [q; q]_n [1/y^2; q^2]_n [x^2q^2; q^2]_m [y^2q^2; q^2]_{m-n}}{[x, xq; q]_n [1/x^2y^2; q]_n [q^2; q^2]_m [x^2y^2q^2; q^2]_{m-n}} \tag{2.13}
 \end{aligned}$$

Where m is the greatest integer  $\leq n/2$ .

$$\begin{aligned}
 & {}_3\Phi_2 \left[ \begin{matrix} q^{-n}, q^{-n}/x^2, 0 \\ q^{1-n}/x, -q^{-n}/x; q \end{matrix}; q \right] \\
 &= \frac{(-)^n [q; q]_n [x^2q^2; q^2]_m x^{n-2m}}{[x, -xq; q]_n [q^2; q^2]_m}, \tag{2.14}
 \end{aligned}$$

Where m is the greatest integer  $\leq n/2$ .

$$= \frac{(-)^n x^n q^{n(n-1)/2} [q; q]_n [x^2 q^2; q^2]_m}{[x, -xq; q]_n [q^2; q^2]_m} {}_3\Phi_2 \left[ \begin{matrix} q^{-n}, q^{-n}/x^2, 0; q; q \\ q^{1-n}/x, -q^{-n}/x; q \end{matrix} \right] \tag{2.15}$$

Where m is the greatest integer  $\leq n/2$ .

Putting  $wq^m$  for w in [Alsalam and Verma 1; (4.3) P.420] we get the summation formula:

$$= \frac{[w; q]_{2m} [w/a, -q; q]_m}{[w/a; q]_{2m} [w, -aq; q]_m} {}_4\Phi_3 \left[ \begin{matrix} a, aq, a^2 q^{2-2m}/w^2, q^{-2m}; q^2, q^2 \\ a^2 q^2, aq^{1-2m}/w, aq^{2-2m}/w \end{matrix} \right] \tag{2.16}$$

$$= \frac{[cd; q]_n [c, d, -q^{1/2}; q^{1/2}]_n}{[c, d; q]_n [cd; q^{1/2}]_n} {}_4\Phi_3 \left[ \begin{matrix} q^{-n}, c, d, \frac{1}{cd} q^{\frac{1}{2}-n}; q; q^2 \\ \frac{1}{c} q^{1-n}, \frac{1}{d} q^{1-n}, cdq^{1/2} \end{matrix} \right] \tag{2.17}$$

$$= \frac{[cdq^{-1/2}; q^{1/2}]_{2n} [c, d; q^{1/2}]_n [q; q]_n}{[cdq^{-1/2}; q^{1/2}]_n [cdq^{1/2}; q]_n [c, d; q]_n [q^{1/2}; q^{1/2}]_n} {}_4\Phi_3 \left[ \begin{matrix} q^{-n}, c, d, \frac{1}{cd} q^{\frac{3}{2}-n}; q; q \\ \frac{1}{c} q^{1-n}, \frac{1}{d} q^{1-n}, cdq^{1/2} \end{matrix} \right] \tag{2.18}$$

$$= \frac{[q, cd; q]_n [c, d; q^{1/2}]_n q^{-n/2}}{[c, d; q]_n [cdq^{-1/2}; q^{1/2}]_n [q^{1/2}; q^{1/2}]_n} {}_4\Phi_3 \left[ \begin{matrix} q^{-n}, c, d, \frac{1}{cd} q^{\frac{1}{2}-n}; q; q \\ \frac{1}{c} q^{1-n}, \frac{1}{d} q^{1-n}, cdq^{-1/2} \end{matrix} \right] \tag{2.19}$$

### III. MAIN RESULTS

In this section we shall establish certain transformations of double series in the term of single series.

(i) Multiplying both sides of (2.1) by an arbitrary sequence  $B_n$ , summing over n from 0 to  $\infty$ , applying the identity (1.3) and then replacing  $B_n$  by  $\frac{z^n}{[q; q]_n} A_n$ , where  $A_n$  is another arbitrary sequence, we get:

$$\sum_{n,k=0}^{\infty} A_{n+k} \frac{[a; q]_k (-cz/a)^k z^n q^{k(k-1)/2}}{[c; q]_k [q; q]_k [q; q]_n} = \sum_{n=0}^{\infty} A_n \frac{[c/a; q]_n z^n}{[q, c; q]_n} \tag{3.1}$$

This is a transformation which reduces a double series in terms of a single series.

Similarly, one can easily establish the following results:

$$(ii) \sum_{n,k=0}^{\infty} A_{n+k} \frac{[a, b; q]_k [c/ab; q]_n (cz/ab)^k z^n}{[q, c; q]_k [q; q]_n} = \sum_{n=0}^{\infty} A_n \frac{[c/a, c/b; q]_n z^n}{[q, c; q]_n} \tag{3.2}$$

(Using (2.2) with  $B_n = \frac{[c/ab; q]_n z^n}{[q; q]_n} A_n$ )

$$(iii) \sum_{n,k=0}^{\infty} A_{n+k} \frac{[x; q]_k [xq; q]_n (-zq)^k z^n}{[q; q]_k [q; q]_n} = \sum_{n=0}^{\infty} A_n \frac{[x^2 q^2; q^2]_m z^n}{[q^2; q^2]_m} \tag{3.3}$$

Where m is the greatest integer  $\leq n/2$ .

(Using (2.3) with  $B_n = \frac{[xq;q]_n z^n}{[q;q]_n} A_n$  )

$$(iv) \sum_{n,k=0}^{\infty} A_{n+k} \frac{[x,y;q]_k [xq,yq;q]_n (-zq)^k z^n}{[q,-xyq;q]_k [q,-xyq;q]_n} = \sum_{n=0}^{\infty} A_n \frac{[xyq;q]_n [x^2q^2,y^2q^2;q^2]_m z^n}{[-xyq;q]_n [q^2,x^2y^2q^2;q^2]_m} \quad (3.4)$$

Where m is the greatest integer  $\leq n/2$ .

(Using (2.4) with  $B_n = \frac{[xq,yq;q]_n z^n}{[q,-xyq;q]_n} A_n$  )

$$(v) \sum_{n,k=0}^{\infty} A_{n+k} \frac{[x;q]_k [xq;q]_n (-z)^k z^n}{[q;q]_k [q;q]_n} = \sum_{n=0}^{\infty} A_n z^n \frac{[x^2q^2;q^2]_m x^{n-2m}}{[q^2;q^2]_m} \quad (3.5)$$

Where m is the greatest integer  $\leq n/2$ .

(Using (2.5) with  $B_n = \frac{[xq;q]_n z^n}{[q;q]_n} A_n$  )

$$(vi) \sum_{n,k=0}^{\infty} A_{n+k} \frac{[x,-xq;q]_n (-z)^k z^n}{[q,x^2q;q]_n [q;q]_k} = \sum_{n=0}^{\infty} A_n (-z)^n \frac{[x^2q^2;q^2]_m x^{n-2m}}{[x^2q;q]_n [q^2;q^2]_m} \quad (3.6)$$

Where m is the greatest integer  $\leq n/2$ .

(Using (2.6) with  $B_n = \frac{[x-xq;q]_n z^n}{[q,x^2q;q]_n} A_n$  )

$$(vii) \sum_{n,k=0}^{\infty} A_{n+k} \frac{[x,-xq;q]_n (-z)^k z^n q^{k(k-1)/2}}{[q,x^2q;q]_n [q;q]_k} = \sum_{n=0}^{\infty} A_n \frac{(-zx)^n q^{n(n-1)/2} [x^2q^2;q^2]_m}{[x^2q;q]_n [q^2;q^2]_m} \quad (3.7)$$

Where m is the greatest integer  $\leq n/2$ .

(Using (2.7) with  $B_n = \frac{[x,-xq;q]_n z^n}{[q,x^2q;q]_n} A_n$  )

$$(viii) \sum_{n,k=0}^{\infty} A_{n+k} \frac{[a,aq;q^2]_k [wq/a,w/a;q^2]_n (zq)^k z^n}{[q^2,a^2q^2;q^2]_k [q^2,w^2/a^2;q^2]_n} = \sum_{n=0}^{\infty} A_n \frac{[w;q]_{2n} z^n}{[w;q]_n [q;q]_n [-aq,-w/a;q]_n} \quad (3.8)$$

(Using (2.8) with  $B_n = \frac{[w/a,wq/a;q^2]_n z^n}{[q^2,w^2/a^2;q^2]_n} A_n$  )

#### IV. CLAUSEN TYPE IDENTITIES

In this section, we deduce the Clausen type identities from the result established in section (3)

(i) Taking  $A_n = 1$  in (3.2) we get

$$= {}_2\Phi_1 \left[ \begin{matrix} c/a, c/b; q; z \\ c \end{matrix} \right], \quad {}_2\Phi_1 \left[ \begin{matrix} a, b; qcza/ab \\ c \end{matrix} \right] {}_1\Phi_0 \left[ \begin{matrix} c/ab; q; z \\ - \end{matrix} \right] \quad (4.1)$$

Which is the basic analogue of Euler’s transformation.

(ii) For  $A_n = 1$ , (3.4) yields the product formula:

$$\begin{aligned}
 & {}_2\Phi_1 \left[ \begin{matrix} x, y; q; -zq \\ -xyq \end{matrix} \right] {}_2\Phi_1 \left[ \begin{matrix} xq, yq; q; z \\ -xyq \end{matrix} \right] \\
 &= {}_4\Phi_3 \left[ \begin{matrix} xyq, xyq^2, x^2q^2, y^2q^2; q^2; z^2 \\ -xyq, -xyq^2, x^2y^2q^2 \end{matrix} \right] \\
 &+ \frac{z(1-xyq)}{(1+xyq)} {}_4\Phi_3 \left[ \begin{matrix} xyq^2, xyq^3, x^2q^2, y^2q^2; q^2; z^2 \\ -xyq^2, -xyq^3, x^2y^2q^2 \end{matrix} \right] \tag{4.2}
 \end{aligned}$$

Which is known result [Verma and Jain 1; (2.37)P.1031]

Similarly, taking  $A_n = 1$  in (3.3) - (3.8) we have the following results respectively.

$$\begin{aligned}
 \text{(iii)} \quad & {}_2\Phi_1 \left[ \begin{matrix} xq, yq; q; -z/q \\ -xyq \end{matrix} \right] {}_2\Phi_1 \left[ \begin{matrix} x, y; q; z \\ -xyq \end{matrix} \right] \\
 &= {}_4\Phi_3 \left[ \begin{matrix} xyq, xyq^2, x^2q^2, y^2q^2; q^2; z^2/q^2 \\ -xyq, -xyq^2, x^2y^2q^2 \end{matrix} \right]
 \end{aligned}$$

$$\begin{aligned}
 & - \frac{z(1-xyq)}{q(1+xyq)} {}_4\Phi_3 \left[ \begin{matrix} xyq^2xyq^3, x^2q^2, y^2q^2; q^2; z^2/q^2 \\ -xyq^2, -xyq^3, x^2y^2q^2 \end{matrix} \right] \tag{4.3}
 \end{aligned}$$

$$\begin{aligned}
 \text{(iv)} \quad & {}_2\Phi_1 \left[ \begin{matrix} y, -y; q; -zq \\ y^2q \end{matrix} \right] {}_2\Phi_1 \left[ \begin{matrix} xq, -xq; q; z \\ x^2q \end{matrix} \right] \\
 &= {}_4\Phi_3 \left[ \begin{matrix} xyq, -xyq, xyq^2, -xyq^2; q^2; z^2 \\ x^2q, y^2q, x^2y^2q^2 \end{matrix} \right]
 \end{aligned}$$

$$\begin{aligned}
 & + \frac{z(1-x^2y^2q^2)}{(1-x^2q)(1-y^2q)} {}_4\Phi_3 \left[ \begin{matrix} xyq^2, -xyq^2, xyq^3, -xyq^3; q^2; z^2 \\ x^2q^3, y^2q^3, x^2y^2q^2 \end{matrix} \right] \tag{4.4}
 \end{aligned}$$

$$\begin{aligned}
 \text{(v)} \quad & {}_2\Phi_1 \left[ \begin{matrix} xq, -yq; q; z \\ x^2q \end{matrix} \right] \\
 &= [-zq; q]_{\infty} {}_0\Phi_1 \left[ \begin{matrix} -; q^2; z^2 \\ x^2q \end{matrix} \right] + \frac{z[-zq; q]_{\infty}}{(1-x^2q)} {}_0\Phi_1 \left[ \begin{matrix} -; q^2; z^2 \\ x^2q^3 \end{matrix} \right] \tag{4.5}
 \end{aligned}$$

$$\begin{aligned}
 \text{(vi)} \quad & {}_2\Phi_1 \left[ \begin{matrix} c, d; q; zq^{1/2} \\ cdq^{1/2} \end{matrix} \right] {}_2\Phi_1 \left[ \begin{matrix} c, d; q; z \\ cdq^{1/2} \end{matrix} \right] \\
 &= {}_4\Phi_3 \left[ \begin{matrix} c, d, \sqrt{cd}, -\sqrt{cd}; q^{1/2}; z \\ cd, q^{1/4}\sqrt{cd}, -q^{1/4}\sqrt{cd} \end{matrix} \right] \tag{4.6}
 \end{aligned}$$

$$\begin{aligned}
 \text{(vii)} \quad & {}_2\Phi_1 \left[ \begin{matrix} c, d; q; zq^{1/2} \\ cdq^{1/2} \end{matrix} \right] {}_2\Phi_1 \left[ \begin{matrix} c, d; q; z \\ cdq^{-1/2} \end{matrix} \right] \\
 &= {}_4\Phi_3 \left[ \begin{matrix} c, d, \sqrt{cd}, -\sqrt{cd}; q^{1/2}; z \\ cdq^{-1/2}, q^{1/4}\sqrt{cd}, -q^{1/4}\sqrt{cd} \end{matrix} \right] \tag{4.7}
 \end{aligned}$$

$$\begin{aligned}
 \text{(viii)} \quad & {}_2\Phi_1 \left[ \begin{matrix} c, d; q; zq^{-1/2} \\ cdq^{-1/2} \end{matrix} \right] {}_2\Phi_1 \left[ \begin{matrix} c, d; q; z \\ cdq^{1/2} \end{matrix} \right] \\
 &= {}_4\Phi_3 \left[ \begin{matrix} c, d, \sqrt{cd}, -\sqrt{cd}; q^{1/2}; zq^{-1/2} \\ cdq^{-1/2}, q^{1/4}\sqrt{cd}, -q^{1/4}\sqrt{cd} \end{matrix} \right] \tag{4.8}
 \end{aligned}$$

## V. CONCLUSION

In this paper a new method has been developed to establish certain transformation of double q-series in terms of a single series. These results lead to certain Clausen type identities. With the help of these results it is also possible to establish certain continued fraction representation involving q- series.



## VI. ACKNOWLEDGEMENT

My thanks are due to Dr. G.C Chaubey Ex Associate Professor & Head department of Mathematics TDPG College Jaunpur, Professor B. Kunwar Department of Mathematics IET, Lucknow and Dr. S.K Mishra Assistant Professor department of Physics IET, Lucknow for their encouragement and for providing necessary support. I am extremely grateful for their constructive support.

## REFERENCES

- [1] Gasper, G. and Rahman, M. (1991): Basic hypergeometric series, Cambridge University Press
- [2] Agarwal, R.P., Manocha, H.L. and Rao, K.Srinivas (2001); Selected Topics in special functions, Allied Publisher Limited, New Delhi
- [3] Agarwal, R.P.: Generalized hypergeometric series and it's application to the theory of combinatorial analysis and partition theory (Unpublished monograph)
- [4] L.J. Slater : Generalized Hypergeometric Functions, Cambridge University Press,( 1966).
- [5] S. Ramanujan : Notebook, Vol. II, Tata Institute of Fundamental Research, Bombay, (1957).
- [6] Verma and Jain "Some summation formulae of basic hypergeometric series" , Indian J. of Pure and Applied Math. 11 (8),1021-1038
- [7] Verma and Jain "Some summation formulae for non terminating basic hypergeometric series" Siam.J. Math.Anal. Vol 16 No.3 1985.



10.22214/IJRASET



45.98



IMPACT FACTOR:  
7.129



IMPACT FACTOR:  
7.429



# INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24\*7 Support on Whatsapp)