



IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 5 Issue: VII Month of publication: July 2017 DOI:

www.ijraset.com

Call: 🛇 08813907089 🕴 E-mail ID: ijraset@gmail.com



Bolpur (Semi Urban), India

Suraj Ghosh¹, Rumi Rabha², Mallika Chowdhury³ and Pratap Kumar Padhy⁴

^{1,2,3,4} Department of Environmental Studies, Siksha Bhavana, Visva-Bharati Santiniketan – 731235, West Bengal, India

Abstract: Monitoring of particulate matter, gaseous pollutants and heavy metals has been done at Bolpur, a semi-urban area of West Bengal, India; during summer from April to June (2014) and winter from December to February (2015). In this study the elemental characterization of airborne particulate matter (PM_{10}) has been recognized as a significant air quality parameters with the study of criteria air pollutants like, PM, SO₂, NO₂, O₃, CO and Pb. In this study all pollutants were analyzed separatelyin both summer and winter seasons and were explained with the help of statistical techniques like ANOVA test, Pearson correlation test, box plot and cluster analysis. During this study period, variation of temperature, rainfall, wind speed and humidity were shown as one main factor for seasonal variation of air pollution. Seasonal variation of pollution levels in outdoor air of Bolpur was found to be highest at all the locations in the month of winter compared with summer. Three statistically significant cluster groups (Cluster-I, Cluster-II and Cluster-III) were displayed during Hierarchical Cluster Analysis depending on pollutants characteristics. Keywords: Particulate matter, Gaseous air pollutants, Heavy metals, Seasonal variation, Cluster analysis.

I. INTRODUCTION

In recent year's rapid growth of industrialization, urbanization, fossil fuel combustion and increasing economic growth are responsible for higher concentration of air borne particulate matter and gaseous air pollutants [1]. Particularly in developing countries like India, rapid industrialization and modernization activities led to the increased use of fossil fuels in huge amount for energy demand. In addition, India characterized by its rapid population growth, where 1.3 billon population also plays an important role in increasing environmental pollution [2]. It is of concern that atmospheric particulate pollution is one of the major pollutants that affect air quality in urban area and even rural areas is getting importance worldwide. Particulate matter is a complex mixture of solid and liquid particles that vary in size with aerodynamic diameters less than 2.5 mm and 10 mm (PM_{2.5} and PM_{10}) and contains heavy metals and ions and which remain suspended in ambient air. The major anthropogenic sources of particulate matter are the incomplete combustion of biofuel, fossil fuel burning, agricultural residual burning, municipal solid waste treatment, construction activities and emission from paved and unpaved roads [3]. Apart from the impact of particulate matter, gaseous air pollutants play significant roles on human health and environment. For example, the contribution and accumulation of many gaseous pollutants including SO₂, NO₂, O₃ and CO increase the pollution levels of atmosphere and susceptibility to respiratory diseases. Soluble components (e.g., metals and ionic compounds) and fine or ultrafine particulate matter may be absorbed through the respiratory tract surface into the systemic circulation. The aim of this paper is to examine the seasonal variation (summer and winter) of various air pollutants like total suspended particulate matter (TSPM), PM₁₀, PM_{2.5} and the harmful trace metals present in PM_{10} samples collected during 2014-2015 from Bolpur, West Bengal.

II. MATERIAL AND METHODS

A. Study Area

Bolpur is a non-industrial and semi-urban area with a population of 80,210 people and area of 18 km². The metropolis has the geographic coordinate of 23°40'N 87°43 E. People of some areas in Bolpur mostly depend on wood or other solid biomass fuels for cooking and other purposes. Total six sites (Table 1 & Figure 1) from Bolpur were selected i.e., Station Chowrasta (S₁), Chitra More (S₂), Bolpur Post Office (S₃), Bolpur Bus Stand (S₄), Siksha Bhavana (S₅) and Sriniketan (S₆).

International Journal for Research in Applied Science & Engineering Technology (IJRASET)



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor:6.887 Volume 5 Issue VIII, July 2017- Available at <u>www.ijraset.com</u>

TABLE 1									
Latitudinal and longitudinal locational details of sampling sites at Bolpur									
	Sampling sites	Latitude	Longitude						
	Bolpur								
\mathbf{S}_1	Station Chowrasta	23°39'41"	87°41'47"						
S ₂	Chitra More	23°40'04"	87°41'39"						
S ₃	Post Office	23°39'48"	87°41'27"						
S_4	Bolpur Bus Stand	23°39'55"	87°40'53"						
S ₅	Siksha Bhavana	23°40'52"	87°40'36''						
S ₆	Sriniketan	23°39'56"	87°39'43''						



Figure 1: Map of the study area showing location of the sampling site and its surroundings places in Bolpur, India.



ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor:6.887 Volume 5 Issue VIII, July 2017- Available at www.ijraset.com

B. Sampling Method and Processing

- 1) Particulate Matter (PM) Sampling: Particulate matter samples were collected on glass fibre filter paper using Portable MiniVol TAS (Tactical Air Sampler) instruments (Air Metrics, Eugene, OR, USA) for size selective particulate matter i.e. TSPM, PM₁₀ and PM_{2.5}. The low flow technology of the instrument was developed with the association of U. S. Environmental Protection Agency (EPA). The particulate mass concentrations were measured gravimetrically by weighing the particles collected and knowing the total volume of air sampled. Filter papers were kept in a desiccator for 24 h before and after the sample collection. Field and laboratory blank filter samples were routinely analysed for particulate matter to evaluate analytical bias and precision.
- 2) Sampling and Measurement of Gaseous Air Pollutants: Sulphur di-oxide (SO₂), nitrogen dioxide (NO₂), ozone (O₃) and carbon monoxide (CO) data were collected using the portable gas sensor and different sensor head with data logger (AeroQual, Series-500). Before proceeding with the experiment, the air quality sensor was switched on. The sensor head was placed perpendicular to the air flow with the monitor and prepared for warm up. When the monitor is first switched on, it will warm up for 3 minutes. The reading will then flash for the next 7 minutes to indicate that the sensor is still in the warm up phase. It is recommended that the monitor is kept in Stand By mode when not being used to keep the sensor heated and prevent the build-up of contaminants.

C. Analysis of Heavy Metals

After gravimetric analysis, collected PM_{10} sample filters were digested in mixed tri-acid i.e. HNO_3 , $HCIO_4$ and H_2SO_4 in the ratio of 5:1:1 using the method of Foley et al [4]. Reagent blank was also prepared by using unexposed filter paper following the same procedure. The digested solutions were then analyzed by using Anodic Stripping Voltameter (Metrohm 797 VA Computrace, Switzerland) for heavy metals like manganese (Mn), zinc (Zn), cadmium (Cd), lead (Pb), nickel (Ni), and cobalt (Co).

D. Statistical Analysis

One-way ANOVA was used to determine interrelationships between parameters and results are shown as the mean \pm standard deviation (SD). Multiple DA (Discriminant Analysis) was performed to reveal and explain essential air quality parameters. DA is a dimension reduction statistical technique to classify data. Where necessary, results were transformed to meet the requirements of normality and equal variance.

E. Meteorological Data

Monthly meteorological parameters were recorded from six different sampling sites of Bolpur during study period. Continuous maximum and minimum temperature (°C), relative humidity (%), wind speed (m/s) and total precipitation were obtained from local weathering station. In summer the average temperature, relative humidity (RH), wind speed and total rainfall of Bolpur were 33.58 °C, 47.81%, 2.9 m/s and 332.92 mm. respectively whereas in winter the average temperature, relative humidity (RH), wind speed and total rainfall of Bolpur were 20.79 °C, 46.50 %, 2.61 m/s and 7.99 mm respectively (Figure 2).







III. RESULTS AND DISCUSSIONS

A. Comparative Study

Six criteria pollutants (carbon monoxide, lead, nitrogen dioxide, ozone, particulate matter, and sulphur dioxide) and other pollutants (heavy metals) were collected from Bolpur area in summer and winter seasons. These seasonal pollution data were analysed statistically in this study and described briefly.



Figure 3: Seasonal variation of Particulate Matter in outdoor air of Bolpur.

In this study seasonal variation of TSPM, PM_{10} and $PM_{2.5}$ in outdoor air in six different sites of Bolpur are shown in Figure 3. Levels of TSPM in outdoor air sampling (24 hours) of Bolpur was varied from 242.23 µg/m³ to 519.3 µg/m³ in summer and 223.6 µg/m³ to 669.45 µg/m³ in winter, PM_{10} was varied from 85.75 µg/m³ to 280.1 µg/m³ in summer and 133.34 µg/m³ to 315.25 µg/m³ in winter and $PM_{2.5}$ was varied from 63.2 µg/m³ to 112.84 µg/m³ in summer and 61.67 µg/m³ to 139.04 µg/m³ in winter respectively. From the Figure 1, seasonal variation of particulate pollution levels shows that winter season had higher



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor:6.887 Volume 5 Issue VIII, July 2017- Available at <u>www.ijraset.com</u>

pollution level than summer. In this study period S_5 (Siksha Bhavana) site had very low pollution level compared to the other sites in both season whereas the pollution level was high at S_4 (Bus Stand) site in winter season.



Figure 4: Seasonal variation of Gaseous Air Pollutants levels in outdoor air of Bolpur

The seasonal variation of ambient gaseous air pollution levels in outdoor air in six different sites of Bolpur are shown in Figure 4. Levels of SO₂ in outdoor air sampling of Bolpur varied from 10 μ g/m³ to 130 μ g/m³ in summer and 10 μ g/m³ to 140 μ g/m³ in winter, NO₂ varied from 82 μ g/m³ to 237 μ g/m³ in summer and 105 μ g/m³ to 285 μ g/m³ in winter, O₃ varied from 0 μ g/m³ to 39 μ g/m³ in summer and 0 μ g/m³ to 42 μ g/m³ in winter and CO varied from 0.5 mg/m³ to 4.7 mg/m³ summer and 0.7 mg/m³ to 4.3 mg/m³ summer respectively. In this study, at S₅ (Siksha Bhavana) site ozone concentration were observed in zero level during



the summer and winter season. Sulphur di-oxide, NO_2 and O_3 levels in all sites except S_4 site were very high which exceeded the NAAQS. Higher levels of CO were found at S_2 and S_4 sites due to high vehicular activities.



Figure 5: Seasonal variation of heavy metal concentration levels in outdoor air of Bolpur

Figure 5 show that, concentration of manganese (Mn), zinc (Zn), cadmium (Cd), lead (Pb), nickel (Ni), and cobalt (Co) in PM₁₀ samples were found in the range of 0.013-0.032 μ g/m³, 0.092-0.263 μ g/m³, 1.148-3.302 ng/m³, 0.024-0.044 μ g/m³, 12.948-29.297 ng/m³ and 0.01-0.025 ng/m³ respectively during summer and the same for winter were 0.015-0.034 μ g/m³, 0.101-0.276 μ g/m³, 1.2063-3.833 ng/m³, 0.0278-0.071 μ g/m³, 12.083-34.688 ng/m³, 0.009-0.0367 ng/m³ respectively. Levels of heavy metals were found highest at all the locations in the month of winter compared to summer. Generally the winter season is characterized by low wind speed and serenity as compared to the summer season. High concentration of heavy metals may be because of poor dispersing condition due to low wind speed and surrounding conditions in this time. TABLE 2

Annual Mean ± 5D concentration of rariculate matter, gaseous ponutants and neavy metals in Bolpur									
Parameters S1		S2	S 3	S4	S4 S5				
TSPM ($\mu g/m^3$)	431.41 ± 50.50	549.8 ± 43.13	385.93 ± 10.79	564.15 ± 148.84	232.91 ± 13.17	520.24 ± 74.77			
$PM_{10}(\mu g/m^3)$	277.75 ± 22.74	297.67 ± 24.85	153.23 ± 46.33	206.01 ± 7.21	$\begin{array}{c} 109.54 \pm \\ 33.65 \end{array} 184.38 \pm 5. \end{array}$				
$PM_{2.5} (\mu g/m^3)$	$94.68 \hspace{0.1 in} \pm \hspace{0.1 in} 2.91$	110.74 ± 5.71	97.03 ± 12.79	125.97 ± 18.49	62.44 ± 1.08	79.83 ± 9.85			
$SO_2 (\mu g/m^3)$	75 ± 7.07	127.5 ± 10.6	115 ± 7.07	135 ± 7.07	10 ± 0	95 ± 7.07			
$NO_2 (\mu g/m^3)$	109 ± 11.31	214 ± 18.38	164.5 ± 12.02	261 ± 33.94	$93.5\ \pm 16.26$	217 ± 14.14			
$O_3(\mu g/m^3)$	30 ± 1.41	31 ± 0	25 ± 5.65	40.5 ± 2.12	0	20 ± 2.83			

International Journal for Research in Applied Science & Engineering Technology (IJRASET)



Pb

Ni

Co

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 6.887 Volume 5 Issue VIII, July 2017- Available at www.ijraset.com

0.49

0.749 1

1

0.739

1

$CO (\mu g/m^3)$	0.95 ± 0.07	2.25 ± 0.63	0.8 ± 0.28	4.55 ± 0.21	0.6 ± 0.14	1.55 ± 0.35
Mn ($\mu g/m^3$)	0.028 ± 0.007	$0.032\ \pm 0$	0.017 ± 0.003	0.029 ± 0.003	0.014 ± 0.001	0.021 ± 0.001
$Zn (\mu g/m^3)$	0.204 ± 0.028	0.259 ± 0.023	$0.161\ \pm 0$	0.263 ± 0.001	0.096 ± 0.006	$0.134\ \pm 0.03$
$Cd (ng/m^3)$	2.571 ± 1.151	3.169 ± 0.305	1.483 ± 0.104	$3.567 \ \pm 0.376$	1.177 ± 0.041	2.104 ± 0.398
Pb ($\mu g/m^3$)	0.054 ± 0.015	0.055 ± 0.021	0.035 ± 0.005	$0.052\ \pm 0.01$	0.025 ± 0.002	0.039 ± 0.006
Ni (ng/m ³)	19.029 ± 0.342	24.062 ± 0.883	18.086 ± 1.97	31.992 ± 3.812	12.516 ± 0.611	15.154 ± 0.917
$Co(ng/m^3)$	0.021 ± 0.009	0.015 ± 0.006	0.025 ± 0.001	$0.031 \ \pm 0.008$	$0.009\ \pm 0$	0.016 ± 0.005

	TABLE 3
Pearson	correlation of different air pollutants of Bolpur area

							· · · ·						
	TSPM	PM10	PM2.5	SO2	NO2	03	CO	Mn	Zn	Cd	Pb	Ni	Co
TSPM	1	0.672	0.779	.870*	.872*	.845*	0.729	.819*	0.776	.863*	0.795	0.727	0.523
PM_{10}		1	0.592	0.52	0.267	0.688	0.288	.928**	0.779	0.754	.938**	0.488	0.2
PM _{2.5}			1	.885*	0.717	.950**	0.804	0.773	.949**	.866*	0.804	.958**	0.798
SO_2				1	.838*	.891*	0.653	0.643	0.782	0.708	0.678	0.763	0.717
NO_2					1	0.675	.834*	0.519	0.611	0.7	0.467	0.728	0.529
O ₃						1	0.708	.820*	.906*	.859*	.875*	.866*	.826*
CO							1	0.619	0.746	.844*	0.563	.922**	0.628
Mn								1	.912*	.943**	.983**	0.743	0.39
Zn									1	.936**	.919**	.918**	0.61
Cd										1	.909*	.896*	0.536

The annual average (mean \pm SD) concentration of pollution level in ambient air of Bolpur is shown in table 2. During the study periods, there are significant variations noticed in the mass concentration of TSPM, PM_{10} and $PM_{2.5}$ in various sites which are summarized in Table 2. Average concentration of each site was compared with the national ambient air quality standard (NAAQS, 2009) and found that all particulate matters concentrations were above the National Ambient Air Quality Standard (NAAQS, 2009) levels prescribed by CPCB, India (Table 2).

The proximity sources of PM were roadways near the sampling point, the influence of traffic-induced coarse dust from roadside, vehicular traffic, fossil fuel combustion, which makes difference between these sites. Resuspension is due to the high temperature and wind velocity as a result the ratios of TSPM, $PM_{2.5}$ and PM_{10} shifts to the lower values during the summer months [5]. This is plausible because mechanically produced particles, and in particular resuspension, depend not only on the vehicle frequency but also on the condition of the roadway (e.g., clean/dirty, wet/dry) [6]. Concentrations in winter were consistently higher compared to summer and this can be due to the differences in meteorological parameters as well as variations in source strengths. Significant differences between PM concentrations in summer and winter have been recorded in various pollution studies [7, 8, 9], and higher winter concentrations are attributed to larger combustion source strength in the winter combined with calm weather conditions and a shallow boundary layer which affects pollutant dispersion adversely [10]. Sulphur dioxide (SO₂) emitted during fossil fuel combustion, is a major precursor of new airborne particles. Vehicle pollutants believed to cause adverse health impacts include diesel particulate matter (DPM), which include ultrafine particles; criteria pollutants, such as, NO₂ and carbon monoxide; and organic compounds such as benzene and butadiene [11]. Vehicular emissions, automobile lubricants, wear and tear of tyres and construction activities are the possible sources of Cd in ambient air. Various products like batteries, toys, etc. contains Cd and is more common in household garbage which are other sources of Cd

* Correlation is significant at the 0.05 level (2-tailed). ** Correlation is significant at the 0.01 level (2-tailed).



International Journal for Research in Applied Science & Engineering Technology (IJRASET) ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor:6.887

Volume 5 Issue VIII, July 2017- Available at <u>www.ijraset.com</u>

[12]. The contribution of Zn and Mn in this study may be the earth crust or wind blow soil and vehicular emission as a contributor in PM_{10} [13]. In ambient air Co may originate mainly from coal burning and re-suspension of upper soil layer [14]. Lead is a major constituent of the lead-acid battery used extensively in car batteries [15]. In vehicular emissions CO, oxides of nitrogen (NOx), sulphur dioxide (SO₂), Pb and particulates are the main pollutants find out in the investigators analysis of diurnal and seasonal variations study by Mondal et al [16] and Pandey et al [17].

B. Agglomerative Hierarchical Cluster Analysis

Agglomerative Hierarchical Cluster Analysis was used to find out the similarity among the sampling sites in Bolpur area. The results of agglomerative hierarchical cluster analysis (AHCA) performed on the dataset after standardization is displayed as dendrogram in Figure 6. The clustering pattern in figure shows three statistically significant groups namely, Cluster-I, Cluster-II and Cluster-III from six sampling sites according to spatial resemblance. Cluster-I included sampling sites S_4 , S_2 and S_6 ; Cluster-II includes sampling sites S_3 and S_1 ; and Cluster-III included sampling sites S_4 , S_2 and S_6 are the main market area where population density is very high; sampling sites S_3 and S_1 are well known for high traffic density and vehicular emission throughout the day; and sampling site S_5 is indicating the silent zone where population and traffic density is low.



Figure 6: Dendrogram of multivariate cluster statistical analysis exhibiting three differentiated groups

IV. CONCLUSIONS

The variations of ambient levels of Particulate matter (TSPM, PM_{10} and $PM_{2.5}$), gaseous air pollutants (SO₂, NO₂, CO and O₃) and heavy metals (Mn, Zn Cd, Pb, Ni and Co) were analysed at the semi urban area of Bolpur during April 2014 to February 2015. Emission from anthropogenic activities mainly traffic, automobile vehicles and biomass burning are the major local



International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor:6.887 Volume 5 Issue VIII, July 2017- Available at www.ijraset.com

outdoor sources of pollutants in the absence of industrial activities. These pollutants were found to have variation in accordance to variation of meteorological parameter during this study period. During winter season particulate matter levels were high due to temperature inversion, high humidity, low rainfall and low wind speed which help particles to settle down. For that reason the metal concentration also showed variation in semi urban area. Depending on behavioural characteristic of sites, there are three clusters (I, II and III) which were extracted by statistical analysis which will help to monitor the ambient air quality in Bolpur.

V. ACKNOWLEDGEMENTS

One of the authors (SG) would like to thank University Grants Commission, New Delhi for providing Non-NET fellowship during this study.

REFERENCES

- R. Yadav, L. K. Sahu, G. Beig, N. Tripathi, and S.N.A. Jaaffrey, "Ambient particulate matter and carbon monoxide at an urban site of India: Influence of anthropogenic emissions and dust storms", Environmental Pollution., vol. 225, pp 291-303, 2017.
- [2] K. Karar, and A. K. Gupta, "Seasonal variations and chemical characterization of ambient PM 10 at residential and industrial sites of an urban region of Kolkata (Calcutta), India", Atmospheric research, vol. 81, no. 1, pp. 36-53, 2006.
- [3] M. Obaidullah, I. V. Dyakov, L. Peeters, S. Bram, and J. D. Ruyck," Measurements of particle concentrations and size distributions in three parking garages", Int J Energy Environ., vol. 5, pp. 508-515, 2012.
- [4] T. Foley, E. A. Betterton, and A. M. A. Wolf, "Ambient PM₁₀ and Metal Concentrations Measured in the Sunnyside Unified School District, Tucson, Arizona", Journal of the Arizona-Nevada Academy of Science, vol. 43, no. 2, pp. 67-76, 2012.
- [5] D. K. Deshmukh, M. K. Deb, and S. L. Mkoma, "Size distribution and seasonal variation of size-segregated particulate matter in the ambient air of Raipur city, India", Air Quality, Atmosphere & Health., vol. 6, no. 1, pp 259-276, 2013.
- [6] B. A. Begum, S. K. Biswas, and P. K. Hopke, "Assessment of trends and present ambient concentrations of PM_{2.2} and PM₁₀ in Dhaka, Bangladesh. Air Quality", Atmosphere & Health., vol. 1, no. 3, pp. 125-133, 2008.
- [7] S. K. Guttikunda, and B. R. Gurjar, "Role of meteorology in seasonality of air pollution in megacity Delhi, India", Environmental monitoring and assessment, vol. 184, no. 5, pp. 3199-3211, 2012.
- [8] S. Yadav, A. Tandon, and A. K. Attri, "Characterization of aerosol associated non-polar organic compounds using TD-GC-MS: a four year study from Delhi, India", Journal of hazardous materials, vol. 252, pp. 29-44, 2013.
- [9] S. Tiwari, S. Pervez, P. Cinzia, D.S. Bisht, A. Kumar, and D.M. Chate, "Chemical characterization of atmospheric particulate matter in Delhi, India, Part II: Source apportionment studies using PMF 3.0", Sustainable Environment Research, vol. 23, no. 5, pp. 295-306, 2013.
- [10] P. Pant, A. Shukla, S. D. Kohl, J. C. Chow, J. G. Watson, and R. M. Harrison, "Characterization of ambient PM_{2.5} at a pollution hotspot in New Delhi, India and inference of sources", Atmos. Environ., vol. 109, pp. 178-189, 2015.
- [11] A. C. Lloyd, and T. A. Cackette, "Diesel engines: environmental impact and control", Journal of the Air & Waste Management Association, vol. 51, no. 6,pp. 809-847, 2001.
- [12] T. K. Rout, R. E. Masto, L. C. Ram, J. George, and P.K. Padhy, "Assessment of human health risks from heavy metals in outdoor dust samples in a coal mining area", Environmental geochemistry and health., vol. 35, no.3, pp 347-356, 2015.
- [13] B. Dubey, A. K. Pal, and G. Singh, "Trace metal composition of airborne particulate matter in the coal mining and non-mining areas of Dhanbad Region, Jharkhand, India." Atmospheric Pollution Research, vol. 3, no. 2, pp. 238-246, 2012.
- [14] A. K. Mishra, S. K. Maiti, and A. K. Pal, "Status of PM[^] sub 10[^] bound heavy metals in ambient air in certain parts of Jharia coal field, Jharkhand, India. International Journal of Environmental Sciences", vol. 4, no.2, pp. 141, 2013.
- [15] M. Ceraolo, "New dynamical models of lead-acid batteries", IEEE transactions on Power Systems, vol. 15, no. 4, pp. 1184-1190, 2000.
- [16] R. Mondal, G. K. Sen, M. Chatterjee, B. K. Sen, and S. Sen, "Ground-level concentration of nitrogen oxides (NOx) at some traffic intersection points in Calcutta", Atmospheric Environment, vol. 34, no. 4, pp. 629-633, 2000. J.
- [17] Pandey, M. Agrawal, N. Khanam, D. Narayan, and D. N. Rao, "Air pollutant concentrations in Varanasi, India. Atmospheric Environment. Part B", Urban Atmosphere, vol. 26, no. 1, pp. 91-98, 1992.











45.98



IMPACT FACTOR: 7.129







INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089 🕓 (24*7 Support on Whatsapp)