



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 5 Issue: VIII Month of publication: August 2017

DOI: <http://doi.org/10.22214/ijraset.2017.8053>

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

Coinoculation of Azospirillum with PGPR for Plant Growth Promotion

Amrita Gupta¹, Pramod Kumar Sahu²

¹Department of Biotechnology, M.S. Ramaiah College of Arts, Science and Commerce, Bengaluru, India-560085

²ICAR-National Bureau of Agriculturally Important Microorganisms, Mau, India-275 103

Abstract: *Azospirillum* is nitrogen fixing rhizobacteria found with grasses and cereals. It imparts many beneficial effects on crop plant like improved germination, dinitrogen fixation, phytohormone production, induced systemic resistance, etc. its impact can be even pronounced by co-inoculating it with other plant growth promoting rhizobacteria. Coinoculation or consortium is an inoculant formulation having containing two or more beneficial microorganisms that synergistically enhances plant growth. Countless literature in last few decades has been accumulated on coinoculation (consortium) of agriculturally beneficial microorganisms and certainly has benefits over single inoculation. Here we have reviewed some of the prominent benefits of coinoculation with *Azospirillum*.

Keywords: *Azospirillum*, Coinoculation, PGPR, biofertilizers

I. INTRODUCTION

Microbes are treasure of plant beneficial features and there are wide array of benefits are provided by microorganisms directly or indirectly to the plants (Sahu et al., 2013; Lavanya et al., 2015a, 2015b; Lavanya et al., 2016; Sahu et al., 2016a, 2016b; Renu et al., 2016a, 2016b; Nair et al., 2017; Renu et al., 2017; Meena et al., 2017). All beneficial microorganisms have their own effects on plant growth and there is no single organism which can provide all these benefits together (Brahmaprakash and Sahu, 2012). In the crop production, it is not practically feasible to apply several PGPR in separate application to the crop. As a consequence of this problem, the concept of consortium came so as to derive more benefit from every application, at the same time, coinoculation also gives synergistic effects. Success of inoculation and profit to the farmer are two major benefits which attracts towards consortium. A group of agriculturally beneficial microorganisms containing a potential N₂ fixer (*Azospirillum*), P- solubilizing bacteria, mycorrhiza, phytohormones producer and/ or a biocontrol agent will give a complete package for success in inoculation. Consortium is an inoculant formulation containing two or more beneficial microorganisms for enhancing plant growth (Ocampo et al. 1975; Brahmaprakash and Sahu, 2012; Sahu and Brahmaprakash, 2016; Sahu et al., 2016a). In consortium, its members may provide nutrients to others, removing inhibitory products by utilizing it, co-metabolism, etc. all these adds to enhanced growth of plant. It is very important to gain farmer's confidence on bioinoculants. In situation where one organism failed to produce its effect, the other one takes over and produce at least some beneficial effects on plant.

A. Benefits of coinoculation to farmers

- 1) Direct improvement in yield by plant growth promotion
- 2) Reduction in disease incidence, thus reduce cost of cultivation by cutting down pesticide use
- 3) Replacement of part of nitrogen and phosphorus fertilizers
- 4) Sustaining soil health

B. *Azospirillum* and *Rhizobium* coinoculation

This is one of the very common consortium of *Azospirillum*. Various green house and field trials have shown that coinoculation of *Azospirillum* and *Rhizobium* to the legumes resulted in increased N₂ fixation, number of nodules, nodule function, hormone induced epidermal-cell differentiation in root hairs and increase in yield (Iruthayathas et al. 1983; Sarig et al. 1986; Andreeva et al. 1993). Phytohormones produced by Epidermal-cell differentiation in root hairs increases the number of potential sites for rhizobial infection (Yahalom et al. 1990).

C. *Azospirillum* and AM fungi coinoculation

Arbuscular mycorrhizal (AM) fungal inoculation has got several benefits for the plant including P-mobilization, biocontrol, alleviation of stress, etc. Coinoculation of AM fungi with *Azospirillum* can enhance nitrogen nutrition to the plants and at the same time it enhances mycorrhizal infection to the plants.

There are various success stories of this consortium (Barea et al. 1983; Pacovsky 1988; Subba Rao et al. 1985). Significant amount of N and P can be replaced for sustained plant growth. Application of *Glomus intraradices* with *Azospirillum* enhanced uptake of N, P, Zn, Cu and Fe apart from increasing mycorrhizal infection (Veeraswamy et al. 1992). Spore germination of *G. fasciculatum* was stimulated by cell free extract of *A. brasilense* and *A. lipoferum* (Tilak and Dwivedi 1990).

D. *Azospirillum* and cellulolytic bacteria coinoculation

A different synergism observed in *Azospirillum* inoculation with cellulolytic bacteria where it enhanced straw decomposition. The association of *Azospirillum* with polysaccharide degraders is helpful for both organisms as it consumes the fermentation products and removes the product inhibition for the polysaccharide degraders. This principle is utilized in augmenting straw decomposition by cellulolytic bacteria in a synergistic way. Apart from increasing N₂ fixation by *Azospirillum*, this consortium was found more efficient in decomposing straw than cellulose decomposing bacteria alone (Halsall and Gibson 1985, Halsall et al. 1985; Markus and Kramer 1988). It was reported that straw amended soil has given higher yields by *Azospirillum* (Hegazi 1988). An increase in pectin degradation was reported by *Bacillus* and N₂ fixation by *Azospirillum* when inoculated together (Khammas and Kaiser 1992). An increase of N₂ fixation by 22 folds was recorded coinoculating with chitinolytic fungi (Halsall 1993).

Some of the organisms enhance nitrogen fixation of *Azospirillum* by providing more congenial environment for nitrogen fixation. It was reported that mangrove rhizosphere bacterium *Staphylococcus* sp. increases the N₂-fixation of the *Azospirillum*. Release of aspartic acid from the *Staphylococcus* sp. cells into the medium may be a cause for this (Holguin and Bashan 1996).

E. *Azospirillum* and PSB coinoculation

Alagawadi and Gaur (1992) reported that combined inoculation of sorghum in field trial by *Azospirillum brasilense* and the phosphate-solubilizing bacteria (*Pseudomonas striata* or *Bacillus polymyxa*) achieved significantly higher yields and enhanced N and P uptake as compared with single inoculation.

Consortium of *Azospirillum lipoferum* and the phosphate-solubilizing *Agrobacterium radiobacter* in a pot experiments in barley showed a significant increase in grain yield by inoculation with mixtures of either compared with single inoculations. The N₂ fixation and uptake is significantly higher in the plants was significantly higher when *Azospirillum lipoferum* and *Agrobacterium radiobacter* were combined. The higher positive impact of consortial inoculation on plant growth and development was recorded with the short supply of combined nitrogen in soil. Field experiments with three barley cultivars confirmed the assertion that inoculation with these mixed cultures is superior to single culture inoculations (Belimov et al. 1995).

F. Compatibility check before coinoculation

An organism secretes many metabolites in the vicinity which may or may not be beneficial for other organisms. There are few reports of negative effects of coinoculation. In clover, coinoculation of *Rhizobium* with *Azospirillum* prevented nodulation. Colonization of root hairs by *Azospirillum* blocked infection site for *Rhizobium* to develop nodule (Plazinski and Rolfe 1985). Therefore, compatibility must be studied before constructing the consortium for coinoculation.

II. CONCLUSION

Azospirillum is very prominent bioinoculant for field use. The efficiency of *Azospirillum* can be even higher if the possibilities of consortial application are harnessed. Much intense improvement is required for preparing improved consortial formulation of *Azospirillum* with stable performance at field. Many other coinoculation techniques for wider benefit in each crop can also be tried for better results of applied inoculants.

III. ACKNOWLEDGEMENT

Authors are grateful to their respective institutes for support and encouragement.

REFERENCES

- [1] Alagawadi AR, Gaur AC (1992) Inoculation of *Azospirillum brasilense* and phosphate-solubilizing bacteria on yield of sorghum [*Sorghum bicolor* (L.) Moench] in dry land. *Trop Agric* 69: 347–350
- [2] Andreeva IN, Red'kina TV, Ismailov SF (1993) The involvement of indoleacetic acid in the stimulation of *Rhizobium* legume symbiosis by *Azospirillum brasilense*. *Russ J Plant Physiol* 40: 901–906
- [3] Barea JM, Bonis AF, Olivares A (1983) Interactions between *Azospirillum* and VA mycorrhiza and their effects on growth and nutrition of maize and ryegrass. *Soil Biol Biochem* 15:705–709
- [4] Belimov AA, Kojemiakov AP, Chuvarliyeva CV (1995) Interaction between barley and mixed cultures of nitrogen fixing and phosphate-solubilizing bacteria. *Plant Soil* 173:29–37
- [5] Brahmaprakash, G. P. and Sahu, P. K. 2012, Biofertilizers for Sustainability. *Journal of Indian Institute of Sciences*, 92:1, pp. 37-62 (2017 IF=0.857)

- [6] Halsall DM (1993) Inoculation of wheat straw to enhance lignocelluloses breakdown and associated nitrogenase activity. *Soil Biol Biochem* 25:419–429.
- [7] Halsall DM, Gibson AH (1985) Cellulose decomposition and associated nitrogen fixation by mixed cultures of *Cellulomonas gelida* and *Azospirillum* species of *Bacillus macerans*. *Appl. Environ. Microbiol* 50:1021–1026
- [8] Halsall DM, Turner GL, Gibson AH (1985) Straw and xylan utilization by pure cultures of nitrogen-fixing *Azospirillum* spp. *Appl Environ Microbiol* 49:423–428
- [9] Hegazi NA (1988) Modification of soil environment through straw application versus *Azospirillum* spp. inoculation. In *Azospirillum*. IV. Genetics, physiology, ecology. W Klingmüller (ed). Springer- Verlag, Berlin, Heidelberg. pp. 215–222.
- [10] Holguin G, Bashan Y (1996) Nitrogen-fixation by *Azospirillum brasilense* Cd is promoted when co-cultured with a mangrove rhizosphere bacterium (*Staphylococcus* sp.). *Soil Biol Biochem* 28: 1651–1660
- [11] Iruthayathas EE, Gunasekaran S, Vlassak K (1983) Effect of combined inoculation of *Azospirillum* and *Rhizobium* on nodulation and N₂-fixation of winged bean and soybean. *Sci Hortic* 20:231–240
- [12] Khammas KM, Kaiser P (1992) Pectin decomposition and associated nitrogen fixation by mixed cultures of *Azospirillum* and *Bacillus* species. *Can J Microbiol* 38:794–797
- [13] Lavanya, G., Sahu, P. K., Brahmaaprakash, G. P., 2016, Survival and effectiveness of fluid bed dried formulation of microbial consortium on cowpea (*Vigna unguiculata*). *Environment and Ecology*, 34(4):2440-2444
- [14] Lavanya, G., Sahu, P. K., Manikanta DS, Brahmaaprakash, G. P. (2015b) Effect of fluid bed dried formulation in comparison with lignite formulation of microbial consortium on finger millet (*Eleusine coracana* Gaertn.). *Journal of Pure and Applied Microbiology* 9(2):193-199.
- [15] Lavanya, G., Sahu, P. K., Manikanta DS, Brahmaaprakash, G. P., 2015a, Standardization of methodology for preparation of microbial consortia of agriculturally beneficial microorganisms using fluid bed dryer. *Journal of soil biology and ecology*. 35(1&2): 126-134.
- [16] Markus P, Kramer J (1988) Importance of nonsymbiotic nitrogen-fixing bacteria in organic farming systems. In *Azospirillum*. IV. Genetics, physiology, ecology. Edited by W. Klingmüller. Springer-Verlag, Berlin, Heidelberg. pp. 197 –204
- [17] Meena KK, Sorty AM, Bitla UM, Choudhary K, Gupta P, Pareek A, Singh DP, Ratna Prabha, Sahu PK, Gupta VK, Singh HB, Krishanani KK and Minhas PS, 2017, Abiotic Stress Responses and Microbe-Mediated Mitigation in Plants: The Omics Strategies. *Frontiers in Plant Sciences*, 8(172): 1-25
- [18] Nair SS, Sahu PK and G P Brahmaaprakash, 2017, Microbial Inoculants for Agriculture In Changing Climates. *Mysore Journal of Agricultural Sciences*, 51(1): 27-44
- [19] Ocampo JA, Barea JM, Montoya E (1975) Interactions between *Azotobacter* and phosphobacteria and their establishment in the rhizosphere as affected by soil fertility. *Can J Microbiol* 21:1160–1165
- [20] Pacovsky RS (1988) Influence of inoculation with *Azospirillum brasilense* and *Glomus fasciculatum* on sorghum nutrition. *Plant Soil* 110:283–287
- [21] Plazinski J, Rolfe BG (1985) Analysis of the pectolytic activity of *Rhizobium* and *Azospirillum* strains isolated from *Trifolium repens*. *J Plant Physiol* 120:181–187
- [22] Renu, Manish S Bhoyar, Udai B. Singh, Upasana Sahu, Dipak T Nagrale, Pramod Kumar Sahu, 2017, Characterization of lytic bacteriophage Xcc9SH3 infecting *Xanthomonas campestris* pv. *Campestris*. *Journal of Plant Pathology*, 99(1) 233-238.
- [23] Renu, Pramod K. Sahu, Upasana Sahu, Manish S. Bhoyar, Rajiv K. Singh and Munish Kumar, 2016b, Microbial dynamics in rhizosphere at different growth stages of rice (*Oryza sativa* L.) based on plant growth promoting functions. *Ann. Agric. Res. New Series Vol. 37 (4): 434-444*
- [24] Renu, Pramod Kumar Sahu, Upasana Sahu, Manish S Bhoyar, Munish Kumar, Udai Bhan Singh and R K Pathak, 2016a, Ecological success of compatible microbes in consortia isolated from rice rhizosphere for growth and yield of mung bean (*Vigna radiata* L.). *Journal of Pure and Applied Microbiology*, 10 (4):3231-3239.
- [25] Sahu PK and G.P. Brahmaaprakash (2016). Formulations of biofertilizers- approaches and advances. In: *Microbial inoculants in sustainable agricultural productivity- vol. 2 Functional application* (eds). Singh DP et al., Springer pp. 179-198. (DOI 10.1007/978-81-322-2644-4_12).
- [26] Sahu PK, Lavanya G, Gupta A and Brahmaaprakash GP, 2016a, Fluid bed dried microbial consortium for enhanced plant growth: A step towards next generation bioformulation. *Vegetos*, 29(4): 6-10.
- [27] Sahu PK, Sharma L, Amrita Gupta, Renu (2016) Rhizospheric and Endophytic Beneficial Microorganisms: Treasure For Biological Control of Plant Pathogens. In: *Recent Biotechnological applications in India* (eds. Subhash Santra and A Mallick). ENVIS centre on Environmental Biotechnology, University of Kalyani, WB, pp: 50-63
- [28] Sahu, P. K., Lavanya, G., Brahmaaprakash, G. P., 2013, Fluid bed dried microbial inoculants formulation with improved survival and reduced contamination level. *Journal of soil biology and ecology*. 33(1&2): 81-94
- [29] Sarig S, Kapulnik Y, Okon Y (1986) Effect of *Azospirillum* inoculation on nitrogen fixation and growth of several winter legumes. *Plant Soil* 90: 335–342
- [30] Subba Rao NS, Tilak KVBR, Singh CS (1985) Synergistic effect of vesicular- arbuscular mycorrhizas and *Azospirillum brasilense* on the growth of barley in pots. *Soil Biol Biochem* 17: 119–121
- [31] Tilak KVBR, Dwivedi A (1990) Enhancement of spore germination of *Glomus fasciculatum* by bacterial cell free extracts. *Indian J Exp Biol* 28:373–375
- [32] Veeraswamy J, Padmavathi T, Venkateswarlu K (1992) Interaction effect s of *Glomus intraradices* and *Azospirillum lipoferum* on sorghum. *Indian J Microbiol* 32:305–308
- [33] Yahalom E, Okon Y, Dovrat A (1990) Possible mode of action of *Azospirillum brasilense* strain Cd on the root morphology and nodule formation in burr medic (*Medicago polymorpha*). *Can J Microbiol* 36:10–14



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)