



iJRASET

International Journal For Research in
Applied Science and Engineering Technology



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 5 Issue: VIII Month of publication: August 2017

DOI: <http://doi.org/10.22214/ijraset.2017.8184>

www.ijraset.com

Call: ☎ 08813907089

E-mail ID: ijraset@gmail.com

New functions in BiČech $sg\beta$ -Biclosure spaces

M. Divya Bharathi¹, A. Francina Shalini²

¹M.Phil, Research Scholar, ²Assistant Professor Department of Mathematics

Nirmala College for Women, Coimbatore, India.

Abstract: The aim of this paper is to introduce the concept of BiČech $sg\beta$ -continuous functions, BiČech $sg\beta$ -Irresolute functions, Totally BiČech $sg\beta$ -continuous functions and Slightly BiČech $sg\beta$ -continuous functions in Biclosure spaces and investigate their characterizations.

Keywords: BiČech $sg\beta$ -closed sets, BiČech $sg\beta$ -open sets, BiČech $sg\beta$ -continuous, BiČech $sg\beta$ Irresolute, Totally BiČech $sg\beta$ -continuous functions and Slightly BiČech $sg\beta$ -continuous functions.

I. INTRODUCTION

Čech spaces were introduced by Eduard Čech [3] (i.e., sets endowed with a grounded, Extensive and additive closure operators) and studied by many others [6][12]. BiČech closure spaces were introduced by K.Chandrasekhara Rao, R. Gowri and V. Swaminathan [4]. N. Levine [13] introduced g -closed sets. D. Andrijevic [1] initiated the study of β -open sets and β -closed sets. In this paper, we analyze the concept of BiČech $sg\beta$ -continuous functions and BiČech $sg\beta$ -Irresolute functions, Totally BiČech $sg\beta$ -continuous functions and Slightly BiČech $sg\beta$ -continuous functions in biclosure spaces and discuss some of their basic properties.

II. PRELIMINARIES

Definition 2.1: Two maps k_1 and k_2 from power set X to itself are called BiČech closure operator on X and the pair (X, k_1, k_2) is called a BiČech closure spaces if the following axioms are satisfied

$$k_1(\varphi) = \varphi \text{ \& } k_2(\varphi) = \varphi$$

$$A \subseteq k_1(A) \text{ \& } A \subseteq k_2(A) \text{ for every } A \subseteq X$$

$$k_1(A \cup B) = k_1(A) \cup k_1(B) \text{ and } k_2(A \cup B) = k_2(A) \cup k_2(B) \text{ for all } A, B \subseteq X.$$

Definition 2.2 [5] A subset A in a BiČech closure space (X, k_1, k_2) is said to be

$$k_i\text{-regular open if } A = \text{int}_{k_i}(k_i(A)), i = 1, 2$$

$$k_i\text{-regular closed if } A = k_i(\text{int}_{k_i}(A)), i = 1, 2$$

$$k_i\text{-semi open if } A \subseteq k_i(\text{int}_{k_i}(A)), i = 1, 2$$

$$k_i\text{-semi closed if } \text{int}_{k_i}(k_i(A)) \subseteq A, i = 1, 2$$

$$k_i\text{-pre open if } A \subseteq \text{int}_{k_i}(k_i(A)), i = 1, 2$$

$$k_i\text{-pre closed if } k_i(\text{int}_{k_i}(A)) \subseteq A, i = 1, 2$$

$$k_i\text{-}\alpha\text{ open if } A \subseteq \text{int}_{k_i}(k_i(\text{int}_{k_i}(A))), i = 1, 2$$

$$k_i\text{-}\alpha\text{ closed if } k_i(\text{int}_{k_i}(k_i(A))) \subseteq A, i = 1, 2$$

$$k_i\text{-}\beta\text{ open if } A \subseteq k_i(\text{int}_{k_i}(k_i(A))), i = 1, 2$$

$$k_i\text{-}\beta\text{ closed if } (\text{int}_{k_i}(k_i(\text{int}_{k_i}(A)))) \subseteq A, i = 1, 2.$$

Definition 2.3: A subset A of a BiČech closure space (X, k_1, k_2) is called biclosed if $k_1A = A = k_2A$ and called biopen if its complement is biclosed.

Definition 2.4: A subset A of a BiČech closure space (X, k_1, k_2) is said to be (k_1, k_2) - g biclosed if $k_2(A) \subseteq G$ whenever $A \subseteq G$ and G is k_1 open set in X .

Definition 2.5: A subset A of a BiČech closure space (X, k_1, k_2) is said to be (k_1, k_2) - $\pi g\alpha$ biclosed if $k_{2\alpha}(A) \subseteq G$ whenever $A \subseteq G$ and G is k_1 π -open set in X .

Definition 2.6: Let (X, k_1, k_2) be a BiČech closure space. A subset $A \subseteq X$ is said to be (k_1, k_2) -w -biclosed set if $k_2(A) \subseteq G$ whenever $A \subseteq G$ and G is k_1 semi-open set in X .

Definition 2.7: Let (X, k_1, k_2) be a BiČech closure space. A subset $A \subseteq X$ is said to be (k_1, k_2) -J- Čech-biclosed set if $k_\alpha(A) \subseteq G$ whenever $A \subseteq G$ and G is k_1 semi-open set in X , where $k_\alpha(A)$ is the smallest α -closed set containing A .

Definition 2.8: Let (X, k_1, k_2) be a BiČech closure space. A subset $A \subseteq X$ is called (k_1, k_2) -sg β closed set if $k_{2\beta}(A) \subseteq G$ whenever $A \subseteq G$ and G is semi-open subset of (X, k_1) where $k_{2\beta}(A)$ is the smallest β -closed set containing A .

III. BICECH sg β – CONTINUOUS FUNCTIONS

Definition 3.1: Let (X, k_1, k_2) and (Y, v_1, v_2) be a BiČech biclosure space. A map $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ is called BiČech sg β -continuous if every $f^{-1}(v)$ is BiČech sg β - open set in (X, k_1, k_2) for every biopen set V in (Y, v_1, v_2) .

Definition 3.2: A function $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ is called

- (a) continuous if $f^{-1}(V)$ is biclosed in X for each biclosed set V of Y .
- (b) g-continuous if $f^{-1}(V)$ is g-biclosed in X for each biclosed set V of Y .
- (c) w-continuous if $f^{-1}(V)$ is w-biclosed in X for each biclosed set V of Y .
- (d) J-continuous if $f^{-1}(V)$ is J-biclosed in X for each biclosed set V of Y .
- (e) $\pi g\alpha$ -continuous if $f^{-1}(V)$ is $\pi g\alpha$ -biclosed in X for each biclosed set V of Y .

Proposition 3.3:

- (a) Every BiČech continuous is BiČech sg β – continuous.
- (b) Every BiČech g – continuous is BiČech sg β – continuous.
- (c) Every BiČech w – continuous is BiČech sg β – continuous.
- (d) Every BiČech J-continuous is BiČech sg β – continuous.
- (e) Every BiČech $\pi g\alpha$ -continuous is BiČech sg β – continuous.

Proof: (a) Let f be a BiČech continuous. Let V be a BiČech open set in (Y, v_1, v_2) . Since f is BiČech continuous, $f^{-1}(V)$ is BiČech open set of (X, k_1, k_2) . Every BiČech open set is BiČech sg β – open set.

This implies that $f^{-1}(V)$ is BiČech sg β – open set of (X, k_1, k_2) , for every BiČech open set V in (Y, v_1, v_2) . (i.e.,) f is BiČech sg β – continuous. Therefore every BiČech continuous is BiČech sg β – continuous.

Note: The proof is obvious for others.

Remark 3.4: Converse of the above theorem need not be true which can be seen from the following example.

Example 3.5: (a) Let $X = \{a, b, c\}$, $Y = \{1, 2, 3\}$.

Define a closure operator k_1, k_2 on X by $k_1\{\varnothing\} = \{\varnothing\}$, $k_1\{a\} = k_1\{a, b\} = \{a, b\}$, $k_1\{c\} = k_1\{b, c\} = \{b, c\}$, $k_1\{b\} = \{b\}$, $k_1\{a, c\} = k_1\{X\} = X$, $k_2\{\varnothing\} = \{\varnothing\}$, $k_2\{a\} = k_2\{a, c\} = \{a, c\}$, $k_2\{b\} = k_2\{b, c\} = \{b, c\}$, $k_2\{c\} = \{c\}$, $k_2\{a, b\} = k_2\{X\} = X$. BiČech closed set of $X = \{X, \varnothing, \{b, c\}\}$.

(k_1, k_2) sg β – closed set of $X = \{X, \varnothing, \{a\}, \{b\}, \{c\}, \{a, c\}, \{b, c\}\}$.

Define a closure operator v_1, v_2 on Y by $v_1\{\varnothing\} = \{\varnothing\}$, $v_1\{1\} = \{1\}$, $v_1\{3\} = \{3\}$, $v_1\{1, 3\} = \{1, 3\}$, $v_1\{2\} = v_1\{1, 2\} = v_1\{2, 3\} = v_1\{Y\} = Y$, $v_2\{\varnothing\} = \{\varnothing\}$, $v_2\{3\} = \{3\}$, $v_2\{1, 2\} = \{1, 2\}$, $v_2\{1\} = v_2\{2\} = v_2\{1, 3\} = v_2\{2, 3\} = v_2\{Y\} = Y$.

Biclosed set of $Y = \{Y, \varnothing, \{3\}\}$.

Let $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ be defined by $f(a) = 2$, $f(b) = 3$, $f(c) = 1$. Then f is (k_1, k_2) sg β – continuous but not continuous. Since for the biclosed set $\{3\}$ in Y , the inverse image $f^{-1}\{3\} = \{b\}$ is not BiČech closed set in X .

Example 3.6: (b) Let $X = \{a, b, c\}$, $Y = \{p, q, r\}$. Define a closure operator k_1, k_2 on X by $k_1\{\varnothing\} = \{\varnothing\}$,

$k_1\{a\} = k_1\{a, c\} = \{a, c\}$, $k_1\{b\} = k_1\{b, c\} = \{b, c\}$, $k_1\{c\} = \{c\}$, $k_1\{a, b\} = k_1\{X\} = X$, $k_2\{\varnothing\} = \{\varnothing\}$, $k_2\{b\} = k_2\{c\} = k_2\{b, c\} = \{b, c\}$, $k_2\{a\} = \{a\}$, $k_2\{a, b\} = k_2\{a, c\} = k_2\{X\} = X$.

g- biclosed set of $X = \{X, \varnothing, \{a\}, \{c\}, \{a, c\}, \{b, c\}\}$.

(k_1, k_2) sg β – closed set of $X = \{X, \varnothing, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}\}$.

Define a closure operator v_1, v_2 on Y by $v_1\{\varnothing\} = \{\varnothing\}$, $v_1\{p\} = \{p\}$, $v_1\{q\} = \{q\}$, $v_1\{p, q\} = \{p, q\}$,

$v_1\{r\} = v_1\{p, r\} = v_1\{q, r\} = v_1\{Y\} = Y$, $v_2\{\varnothing\} = \{\varnothing\}$, $v_2\{p\} = v_2\{q\} = v_2\{p, q\} = \{p, q\}$, $v_2\{r\} = v_2\{p, r\} = v_2\{q, r\} = v_2\{Y\} = Y$. Biclosed set of $Y = \{Y, \varnothing, \{p, q\}\}$. Let $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ be defined by $f(a) = q$, $f(b) = p$, $f(c) = r$. Then f is (k_1, k_2) sg β – continuous but not g-continuous. Since for the biclosed set $\{p, q\}$ in Y , the inverse image $f^{-1}\{p, q\} = \{a, b\}$ is not g-biclosed set in X .

Example 3.7: (c) Let $X = \{a, b, c\}$, $Y = \{1, 2, 3\}$. Define a closure operator k_1, k_2 on X by $k_1\{\emptyset\} = \{\emptyset\}$, $k_1\{c\} = k_1\{a, c\} = \{a, c\}$, $k_1\{a\} = \{a\}$, $k_1\{b\} = k_1\{b, c\} = k_1\{a, b\} = k_1\{X\} = X$, $k_2\{\emptyset\} = \{\emptyset\}$, $k_2\{a\} = \{a\}$, $k_2\{b\} = k_2\{a, b\} = \{a, b\}$, $k_2\{c\} = k_2\{a, c\} = \{a, c\}$, $k_2\{b, c\} = k_2\{X\} = X$.

w-biclosed set of $X = \{X, \emptyset, \{a\}, \{a, b\}, \{a, c\}\}$.

(k_1, k_2) sg β – closed set of $X = \{X, \emptyset, \{a\}, \{b\}, \{c\}, \{a, c\}, \{a, b\}\}$.

Define a closure operator v_1, v_2 on Y by $v_1\{\emptyset\} = \{\emptyset\}$, $v_1\{1\} = \{1\}$, $v_1\{2\} = v_1\{1, 2\} = \{1, 2\}$, $v_1\{3\} = v_1\{1, 3\} = \{1, 3\}$,

$v_1\{2, 3\} = v_1\{Y\} = Y$, $v_2\{\emptyset\} = \{\emptyset\}$, $v_2\{1\} = \{1\}$, $v_2\{2\} = v_2\{3\} = v_2\{2, 3\} = \{2, 3\}$,

$v_2\{1, 3\} = v_2\{1, 2\} = v_2\{Y\} = Y$.

Biclosed set of $Y = \{Y, \emptyset, \{1\}\}$. Let $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ be defined by $f(a) = 3$, $f(b) = 1$, $f(c) = 2$. Then f is (k_1, k_2) sg β – continuous but not w-continuous. Since for the biclosed set $\{1\}$ in Y , the inverse image $f^{-1}\{1\} = \{b\}$ is not w-biclosed set in X .

Example 3.8: (d) Let $X = \{1, 2, 3\}$, $Y = \{a, b, c\}$. Define a closure operator k_1, k_2 on X by $k_1\{\emptyset\} = \{\emptyset\}$, $k_1\{2\} = \{2\}$, $k_1\{3\} = k_1\{1, 3\} = \{1, 3\}$, $k_1\{1\} = k_1\{1, 2\} = k_1\{2, 3\} = k_1\{X\} = X$, $k_2\{\emptyset\} = \{\emptyset\}$, $k_2\{2\} = k_2\{2, 3\} = \{2, 3\}$, $k_2\{3\} = \{3\}$, $k_2\{1\} = k_2\{1, 3\} = \{1, 3\}$, $k_2\{1, 2\} = k_2\{X\} = X$.

BiČech J- closed set of $X = \{X, \emptyset, \{1\}, \{1, 2\}, \{1, 3\}, \{2, 3\}\}$.

(k_1, k_2) sg β – closed set of $X = \{X, \emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}\}$.

Define a closure operator v_1, v_2 on Y by $v_1\{\emptyset\} = \{\emptyset\}$, $v_1\{b\} = v_1\{c\} = v_1\{b, c\} = \{b, c\}$,

$v_1\{a\} = \{a\}$, $v_1\{a, b\} = v_1\{a, c\} = v_1\{Y\} = Y$, $v_2\{\emptyset\} = \{\emptyset\}$, $v_2\{a\} = \{a\}$, $v_2\{b\} = v_2\{a, b\} = \{a, b\}$,

$v_2\{c\} = v_2\{a, c\} = \{a, c\}$, $v_2\{b, c\} = v_2\{Y\} = Y$.

Biclosed set of $Y = \{Y, \emptyset, \{a\}\}$. Let $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ be defined by $f(1) = c$, $f(2) = a$, $f(3) = b$. Then f is (k_1, k_2) sg β – continuous but not J-continuous. Since for the biclosed set $\{a\}$ in Y , the inverse image $f^{-1}\{a\} = \{2\}$ is not in J- biclosed set in X .


Example 3.9: (e) Let $X = \{a, b, c\}$, $Y = \{p, q, r\}$. Define a closure operator k_1, k_2 on X by $k_1\{\emptyset\} = \{\emptyset\}$, $k_1\{a\} = \{a\}$, $k_1\{b\} = k_1\{a, b\} = \{a, b\}$, $k_1\{c\} = k_1\{a, c\} = \{a, c\}$, $k_1\{b, c\} = k_1\{X\} = X$, $k_2\{\emptyset\} = \{\emptyset\}$, $k_2\{b\} = k_2\{c\} = k_2\{b, c\} = \{b, c\}$, $k_2\{a\} = \{a\}$, $k_2\{a, c\} = k_2\{a, b\} = k_2\{X\} = X$.

BiČech $\pi\alpha$ – closed set of $X = \{X, \emptyset, \{a\}, \{a, b\}, \{a, c\}, \{b, c\}\}$.

(k_1, k_2) sg β – closed set of $X = \{X, \emptyset, \{a\}, \{b\}, \{c\}, \{a, c\}, \{b, c\}, \{a, b\}\}$.

Define a closure operator v_1, v_2 on Y by $v_1\{\emptyset\} = \{\emptyset\}$, $v_1\{p\} = \{p\}$, $v_1\{q\} = v_1\{r\} = v_1\{q, r\} = \{q, r\}$, $v_1\{p, q\} = v_1\{q, r\} = v_1\{Y\} = Y$, $v_2\{\emptyset\} = \{\emptyset\}$, $v_2\{p\} = \{p\}$, $v_2\{q\} = v_2\{p, q\} = \{p, q\}$, $v_2\{r\} = v_2\{p, r\} = \{p, r\}$, $v_2\{q, r\} = v_2\{Y\} = Y$.

Biclosed set of $Y = \{Y, \emptyset, \{p\}\}$. Let $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ be defined by $f(a) = r$, $f(b) = p$, $f(c) = q$. Then f is (k_1, k_2) sg β – continuous but not $\pi\alpha$ -continuous. Since for the biclosed set $\{p\}$ in Y , the inverse image $f^{-1}\{p\} = \{b\}$ is not in $\pi\alpha$ -biclosed set in X .



Proposition 3.10: Let (X, k_1, k_2) and (Y, v_1, v_2) be biclosure spaces and let $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ be a map. Then f is BiČech $sg\beta$ – continuous if and only if the inverse image of every BiČech closed subset of (Y, v_1, v_2) is BiČech – $sg\beta$ closed in (X, k_1, k_2) .

Proof: Let F be BiČech closed subset in (Y, v_1, v_2) . Then $Y - F$ is BiČech open in (Y, v_1, v_2) . Since f is BiČech $sg\beta$ – continuous, $f^{-1}(Y - F)$ is BiČech $sg\beta$ – open. But $f^{-1}(Y - F) = X - f^{-1}(F)$ thus $f^{-1}(F)$ is BiČech $sg\beta$ – closed in (X, k_1, k_2) . Conversely let G be an BiČech open subset in (Y, v_1, v_2) . Then $Y - G$ is BiČech closed in (Y, v_1, v_2) . Since the inverse image of each BiČech closed subset in (Y, v_1, v_2) is BiČech $sg\beta$ – closed in (X, k_1, k_2) . We have $f^{-1}(Y - G)$ is to be BiČech $sg\beta$ – closed in (X, k_1, k_2) . But $f^{-1}(Y - G) = X - f^{-1}(G)$. Thus $f^{-1}(G)$ is BiČech $sg\beta$ – open. Therefore f is BiČech $sg\beta$ – continuous.

Remark 3.11: The composition of two BiČech $sg\beta$ – continuous need not be BiČech $sg\beta$ – continuous.

Definition 3.12: A Biclosure space (X, k_1, k_2) is said to be a T_d – space if every BiČech $sg\beta$ – open set in (X, k_1, k_2) is BiČech open.

Proposition 3.13: Let (X, k_1, k_2) and (Z, w_1, w_2) be biclosure spaces and (Y, v_1, v_2) be a T_d – space. If $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ and $g: (Y, v_1, v_2) \rightarrow (Z, w_1, w_2)$ are BiČech $sg\beta$ – continuous, then $g \circ f$ is BiČech $sg\beta$ – continuous.

Proof: Let H be BiČech open in (Z, w_1, w_2) . Since g is BiČech $sg\beta$ – continuous, $g^{-1}(H)$ is BiČech $sg\beta$ – open in (Y, v_1, v_2) . But (Y, v_1, v_2) is a T_d – space, hence $g^{-1}(H)$ is BiČech open in (Y, v_1, v_2) . Thus $f^{-1}(g^{-1}(H)) = (g \circ f)^{-1}(H)$ is BiČech $sg\beta$ – open in (X, k_1, k_2) . Therefore, $g \circ f$ is BiČech $sg\beta$ – continuous.

Proposition 3.14: Let (X, k_1, k_2) , (Y, v_1, v_2) and (Z, w_1, w_2) be biclosure spaces. If $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ is BiČech $sg\beta$ – continuous and $g: (Y, v_1, v_2) \rightarrow (Z, w_1, w_2)$ is continuous then $g \circ f$ is BiČech $sg\beta$ – continuous.

Proof: Let H be an BiČech open subset of (Z, w_1, w_2) . Since g is continuous, $g^{-1}(H)$ is BiČech open in (Y, v_1, v_2) . Since f is BiČech $sg\beta$ – continuous, $f^{-1}(g^{-1}(H))$ is BiČech $sg\beta$ – open in (X, k_1, k_2) . But $f^{-1}(g^{-1}(H)) = (g \circ f)^{-1}(H)$. Therefore, $g \circ f$ is BiČech $sg\beta$ – continuous.

IV. BIČECH $sg\beta$ – IRRESOLUTE FUNCTION

Definition 4.1: Let (X, k_1, k_2) and (Y, v_1, v_2) be biclosure space and a map $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ is called BiČech $sg\beta$ – irresolute, if $f^{-1}(G)$ is BiČech $sg\beta$ – open set (closed set) in (X, k_1, k_2) for every BiČech $sg\beta$ – open set (closed set) G in (Y, v_1, v_2) .

Proposition 4.2: Every BiČech $sg\beta$ – irresolute map is BiČech $sg\beta$ – continuous.

Proof: Assume that f is BiČech $sg\beta$ – irresolute. Let V be a BiČech closed set in Y . Every BiČech closed set is BiČech $sg\beta$ – closed. That implies V be a BiČech $sg\beta$ – closed set in Y . Since f is BiČech $sg\beta$ – irresolute, $f^{-1}(V)$ is BiČech $sg\beta$ – closed set in X . Thus $f^{-1}(V)$ is BiČech $sg\beta$ – closed set in X , \forall BiČech closed set V in Y . That implies f is BiČech $sg\beta$ – continuous.

Remark 4.3: The converse is not true as can be seen from the following example:

Example 4.4: Let $X = \{a, b, c\}$ and $Y = \{1, 2, 3\}$. Define a closure operator k_1, k_2 on X by $k_1\{\phi\} = \{\phi\}$, $k_1\{a\} = k_1\{a, c\} = \{a, c\}$, $k_1\{b\} = k_1\{b, c\} = \{b, c\}$, $k_1\{c\} = \{c\}$, $k_1\{a, b\} = k_1\{X\} = X$, $k_2\{\phi\} = \{\phi\}$, $k_2\{b\} = \{b\}$, $k_2\{a\} = k_2\{a, b\} = \{a, b\}$, $k_2\{c\} = k_2\{b, c\} = \{b, c\}$, $k_2\{a, c\} = k_2\{X\} = X$.

Biclosed set of $X = \{X, \phi, \{b, c\}\}$.

BiČech $sg\beta$ – closed set of $X = \{X, \phi, \{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}\}$.

Define a closure operator v_1, v_2 on Y by $v_1\{\phi\} = \{\phi\}$, $v_1\{2\} = v_1\{1, 2\} = \{1, 2\}$, $v_1\{3\} = v_1\{1, 3\} = \{1, 3\}$, $v_1\{1\} = \{1\}$, $v_1\{2, 3\} = v_1\{Y\} = Y$, $v_2\{\phi\} = \{\phi\}$, $v_2\{1\} = \{1\}$, $v_2\{2\} = v_2\{3\} = v_2\{2, 3\} = \{2, 3\}$,

$v_2\{1, 2\} = v_2\{1, 3\} = v_2\{Y\} = Y$.

Biclosed set of $Y = \{Y, \phi, \{1\}\}$.

BiČech $sg\beta$ – closed set of $Y = \{Y, \phi, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}\}$.

Define a function $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ such that $f(a) = 3$, $f(b) = 2$, $f(c) = 1$. Here f is BiČech $sg\beta$ – continuous.

$f^{-1}\{1, 3\} = \{a, c\}$ is not BiČech $sg\beta$ – closed in (X, k_1, k_2) . Therefore f is not BiČech $sg\beta$ – irresolute.

Proposition 4.5: Let (X, k_1, k_2) and (Y, v_1, v_2) be biclosure spaces and $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ be a map. Then f is BiČech $sg\beta$ – irresolute if and only if $f^{-1}(B)$ is BiČech $sg\beta$ – closed in (X, k_1, k_2) whenever B is BiČech $sg\beta$ – closed in (Y, v_1, v_2) .

Proof: Suppose B be a BiČech $sg\beta$ – closed subset of (Y, v_1, v_2) . Then $Y - B$ is BiČech $sg\beta$ – open in (Y, v_1, v_2) .

Since $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ is BiČech $sg\beta$ – irresolute, $f^{-1}(Y - B)$ is BiČech $sg\beta$ – open in (X, k_1, k_2) .

But, $f^{-1}(Y - B) = X - f^{-1}(B)$, so that $f^{-1}(B)$ is BiČech $sg\beta$ - closed in (X, k_1, k_2) . Conversely, Let A be a BiČech $sg\beta$ - open subset in (Y, v_1, v_2) . Then $Y - A$ is BiČech $sg\beta$ - closed in (Y, v_1, v_2) . By the assumption, $f^{-1}(Y - A)$ is BiČech $sg\beta$ - closed in (X, k_1, k_2) . But $f^{-1}(Y - A) = X - f^{-1}(A)$. Thus $f^{-1}(A)$ is BiČech $sg\beta$ - open in (X, k_1, k_2) . Therefore, f is BiČech $sg\beta$ - irresolute.

Proposition 4.6: Let (X, k_1, k_2) , (Y, v_1, v_2) and (Z, w_1, w_2) be biclosure spaces. If $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ is a BiČech $sg\beta$ - irresolute map and $g: (Y, v_1, v_2) \rightarrow (Z, w_1, w_2)$ is a BiČech $sg\beta$ - continuous map, then the composition $g \circ f: (X, k_1, k_2) \rightarrow (Z, w_1, w_2)$ is BiČech $sg\beta$ - continuous.

Proof: Let G be an open subset of (Z, w_1, w_2) . Then $g^{-1}(G)$ is a BiČech $sg\beta$ - open in (Y, v_1, v_2) as g is BiČech $sg\beta$ - continuous. Hence, $f^{-1}(g^{-1}(G))$ is BiČech $sg\beta$ - open in (X, k_1, k_2) because f is BiČech $sg\beta$ - irresolute. Thus $g \circ f$ is BiČech $sg\beta$ - continuous.

Proposition 4.7: Let (X, k_1, k_2) , (Y, v_1, v_2) and (Z, w_1, w_2) be biclosure spaces. If $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ and $g: (Y, v_1, v_2) \rightarrow (Z, w_1, w_2)$ are BiČech $sg\beta$ - irresolute, then $g \circ f: (X, k_1, k_2) \rightarrow (Z, w_1, w_2)$ is BiČech $sg\beta$ - irresolute.

Proof: Let F be BiČech $sg\beta$ - open set in (Z, w_1, w_2) . As g is BiČech $sg\beta$ - irresolute, $g^{-1}(F)$ is BiČech $sg\beta$ - open in (Y, v_1, v_2) . Since, f is BiČech $sg\beta$ - irresolute, $f^{-1}(g^{-1}(F))$ is BiČech $sg\beta$ - open in (X, k_1, k_2) implies $(g \circ f)^{-1}F = (f^{-1}g^{-1}(F))$ is BiČech $sg\beta$ - open in (X, k_1, k_2) . Hence $g \circ f$ is BiČech $sg\beta$ - irresolute.

Proposition 4.8: Let (X, k_1, k_2) and (Z, w_1, w_2) be biclosure spaces and (Y, v_1, v_2) be a T_d - space. If $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ be a BiČech $sg\beta$ - continuous map and $g: (Y, v_1, v_2) \rightarrow (Z, w_1, w_2)$ is a BiČech $sg\beta$ - irresolute, then the composition $g \circ f: (X, k_1, k_2) \rightarrow (Z, w_1, w_2)$ is BiČech $sg\beta$ - irresolute.

Proof: Let V be BiČech $sg\beta$ - open in Z . Since g is BiČech $sg\beta$ - irresolute, $g^{-1}(V)$ is BiČech $sg\beta$ - open in Y . As Y is a T_d - space, $g^{-1}(V)$ is BiČech open in Y . Since f is BiČech $sg\beta$ - continuous, $f^{-1}(g^{-1}(V))$ is BiČech $sg\beta$ - open in X . Thus $(g \circ f)^{-1}(V)$ is BiČech $sg\beta$ - open in X . Hence $g \circ f$ is BiČech $sg\beta$ - irresolute.

V. TOTALLY BIČECH $sg\beta$ - CONTINUOUS FUNCTION

Definition 5.1: Let (X, k_1, k_2) and (Y, v_1, v_2) be a BiČech closure space. A function $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ is called Totally BiČech continuous if the inverse image of every biopen subset of (Y, v_1, v_2) is a BiČech clopen subset in (X, k_1, k_2) .

Definition 5.2: Let (X, k_1, k_2) and (Y, v_1, v_2) be a BiČech closure space. A function $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ is called Totally BiČech $sg\beta$ - continuous if the inverse image of every biopen subset of (Y, v_1, v_2) is a BiČech $sg\beta$ - clopen subset in (X, k_1, k_2) .

Theorem 5.3: A function $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ is totally BiČech $sg\beta$ - continuous if and only if the inverse image of every biclosed subset of (Y, v_1, v_2) is a BiČech $sg\beta$ - clopen subset in (X, k_1, k_2) .

Proof: Assume that f is totally BiČech $sg\beta$ - continuous. Let A be any biclosed subset in Y . Then A^c is a biopen subset in Y . Since f is totally BiČech $sg\beta$ - continuous. Thus $f^{-1}(A^c)$ is BiČech $sg\beta$ - clopen subset in (X, k_1, k_2) . But $f^{-1}(A^c) = X - f^{-1}(A)$ and so $f^{-1}(A)$ is both BiČech $sg\beta$ - closed subset and BiČech $sg\beta$ - open subset in X . Conversely, let G be a biopen subset in Y . Then G^c is biclosed subset in Y . By assumption $f^{-1}(G^c)$ is BiČech $sg\beta$ - clopen subset in (X, k_1, k_2) . But $f^{-1}(G^c) = X - f^{-1}(G)$ and so $f^{-1}(G)$ is both BiČech $sg\beta$ - closed and BiČech $sg\beta$ - open. Hence $f^{-1}(G)$ is BiČech $sg\beta$ - clopen subset in (X, k_1, k_2) . Therefore f is totally BiČech $sg\beta$ - continuous.

Theorem 5.4: Every totally BiČech $sg\beta$ - continuous $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ is BiČech $sg\beta$ - continuous function.

Proof: Let A be any biopen subset in Y . Since f is totally BiČech $sg\beta$ - continuous. Thus $f^{-1}(A)$ is BiČech $sg\beta$ - clopen subset in (X, k_1, k_2) . (i.e) $f^{-1}(A)$ is both BiČech $sg\beta$ - closed subset and BiČech $sg\beta$ - open subset in X . Thus $f^{-1}(A)$ is BiČech $sg\beta$ - open subset in (X, k_1, k_2) . Therefore f is a $sg\beta$ - continuous.

Remark 5.5: The converse is not true as can be seen from the following example:

Example 5.6: Let $X = \{a, b, c\}$, $Y = \{1, 2, 3\}$. Define a closure operator k_1, k_2 on X by $k_1\{\phi\} = \{\phi\}$,

$k_1\{a\} = k_1\{a, c\} = \{a, c\}$, $k_1\{b\} = k_1\{b, c\} = \{b, c\}$, $k_1\{c\} = \{c\}$, $k_1\{a, c\} = k_1\{X\} = X$, $k_2\{\phi\} = \{\phi\}$,

$k_2\{b\} = \{b\}$, $k_2\{c\} = k_2\{b, c\} = \{b, c\}$, $k_2\{a\} = k_2\{a, b\} = \{a, b\}$, $k_2\{a, c\} = k_2\{X\} = X$.

Biclosed set of $X = \{X, \phi, \{b, c\}\}$.

(k_1, k_2) $sg\beta$ - closed set of $X = \{X, \phi, \{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}\}$.

Define a closure operator v_1, v_2 on Y by $v_1\{\phi\} = \{\phi\}$, $v_1\{1\} = \{1\}$, $v_1\{2\} = v_1\{1, 2\} = \{1, 2\}$, $v_1\{3\} = v_1\{1, 3\} = \{1, 3\}$,

$v_1\{2, 3\} = v_1\{Y\} = Y$, $v_2\{\phi\} = \{\phi\}$, $v_2\{1\} = \{1\}$, $v_2\{2\} = v_2\{3\} = v_2\{2, 3\} = \{2, 3\}$,

$v_2\{1, 3\} = v_2\{1, 2\} = v_2\{Y\} = Y$.

Biclosed set of $Y = \{Y, \phi, \{1\}\}$. Biopen set of $Y = \{Y, \phi, \{2,3\}\}$.

Let $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ be defined by $f(a) = 2, f(b) = 1, f(c) = 3$. Then f is (k_1, k_2) $sg\beta$ - continuous but not totally BiCech $sg\beta$ -continuous. Since for the biopen set $\{2,3\}$ in Y , the inverse image $f^{-1}\{2,3\} = \{a,c\}$ is not totally BiCech $sg\beta$ -clopen set in X .

Theorem 5.7: Let $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ and $g: (Y, v_1, v_2) \rightarrow (Z, w_1, w_2)$ be function. Then $g \circ f: (X, k_1, k_2) \rightarrow (Z, w_1, w_2)$

(i) If f is BiCech $sg\beta$ -irresolute and g is totally BiCech $sg\beta$ -continuous then $g \circ f$ is totally BiCech $sg\beta$ -continuous.

(ii) If f is totally BiCech $sg\beta$ -continuous and g is BiCech continuous then $g \circ f$ is totally BiCech $sg\beta$ -continuous.

Proof: (i) Let U be a BiCech open set in Z . Since g is totally BiCech $sg\beta$ -continuous, $g^{-1}(U)$ is BiCech $sg\beta$ -clopen in Y . Since f is BiCech $sg\beta$ -irresolute, $f^{-1}(g^{-1}(U))$ is BiCech $sg\beta$ -open and BiCech $sg\beta$ -closed in X . Since $g \circ f^{-1}(U) = f^{-1}(g^{-1}(U))$, $g \circ f$ is totally BiCech $sg\beta$ -continuous.

(ii) Let U be a BiCech open set in Z . Since g is BiCech continuous, $g^{-1}(U)$ is BiCech open in Y . Also since f is totally BiCech $sg\beta$ -continuous, $f^{-1}(g^{-1}(U))$ is BiCech $sg\beta$ -clopen in X . Hence $g \circ f$ is totally BiCech $sg\beta$ -continuous.

VI. SLIGHTLY BIČECH $sg\beta$ – CONTINUOUS FUNCTION

Definition 6.1: Let (X, k_1, k_2) and (Y, v_1, v_2) be a BiCech closure space. A function $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ is called Slightly BiCech -continuous if the inverse image of every clopen subset of (Y, v_1, v_2) is a BiCech-open subset in (X, k_1, k_2) .

Definition 6.2: Let (X, k_1, k_2) and (Y, v_1, v_2) be a BiCech closure space. A function $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ is called Slightly BiCech $sg\beta$ -continuous if the inverse image of every clopen subset of (Y, v_1, v_2) is a BiCech $sg\beta$ -open subset in (X, k_1, k_2) .

Theorem 6.3: Every BiCech $sg\beta$ -continuous $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ is Slightly BiČech $sg\beta$ – continuous.

Proof: Let $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ be a BiCech $sg\beta$ -continuous function. Let U be a clopen set in Y . Then $f^{-1}(U)$ is BiCech $sg\beta$ -open in X and BiCech $sg\beta$ -closed in X . Hence f is Slightly BiČech $sg\beta$ – continuous.

Theorem 6.4: Every Slightly BiCech continuous is Slightly BiČech $sg\beta$ – continuous.

Proof: Let $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ be Slightly BiCech continuous function. Let U be a clopen set in Y . Then $f^{-1}(U)$ BiCech open in X . Since every open set is $sg\beta$ -open, $f^{-1}(U)$ BiCech $sg\beta$ –open. Hence f is Slightly BiČech $sg\beta$ – continuous.

Remark 6.5: The converse is not true as can be seen from the following example:

Example 6.6: Let $X = \{a, b, c\}$, $Y = \{1, 2, 3\}$. Define a closure operator k_1, k_2 on X by $k_1\{\phi\} = \{\phi\}$,

$k_1\{a\} = k_1\{a, c\} = \{a, c\}$, $k_1\{b\} = k_1\{b, c\} = \{b, c\}$, $k_1\{c\} = \{c\}$, $k_1\{a, c\} = k_1\{X\} = X$, $k_2\{\phi\} = \{\phi\}$,

$k_2\{\phi\} = \{\phi\}$, $k_2\{b\} = \{b\}$, $k_2\{c\} = k_2\{b, c\} = \{b, c\}$, $k_2\{a\} = k_2\{a, b\} = \{a, b\}$, $k_2\{a, c\} = k_2\{X\} = X$.

Biclosed set of $X = \{X, \phi, \{b, c\}\}$. Biopen set of $X = \{X, \phi, \{a\}\}$.

(k_1, k_2) $sg\beta$ – closed set of $X = \{X, \phi, \{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}\}$.

Define a closure operator v_1, v_2 on Y by $v_1\{\phi\} = \{\phi\}$, $v_1\{1\} = v_1\{1, 3\} = \{1, 3\}$, $v_1\{2\} = \{2\}$,

$v_1\{3\} = v_1\{2, 3\} = v_1\{1, 2\} = v_1\{Y\} = Y$, $v_2\{\phi\} = \{\phi\}$, $v_2\{2\} = \{2\}$, $v_2\{3\} = \{3\}$, $v_2\{1\} = v_2\{1, 3\} = \{1, 3\}$,

$v_2\{2, 3\} = \{2, 3\}$, $v_2\{1, 2\} = v_2\{Y\} = Y$.

Biclopen set of $Y = \{Y, \phi, \{2\}, \{1, 3\}\}$.

Let $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ be defined by $f(a) = 3, f(b) = 1, f(c) = 2$. Then f is slightly BiCech $sg\beta$ – continuous but not slightly BiCech continuous. Since for the clopen set $\{1, 3\}$ in Y , the inverse image $f^{-1}\{1, 3\} = \{a, b\}$ is not slightly BiCech open set in X .

Theorem 6.7: Let $f: (X, k_1, k_2) \rightarrow (Y, v_1, v_2)$ and $g: (Y, v_1, v_2) \rightarrow (Z, w_1, w_2)$ be function.

(i) If f is BiCech $sg\beta$ -irresolute and g is slightly BiCech $sg\beta$ -continuous then $g \circ f$ is slightly BiCech $sg\beta$ -continuous.

(ii) If f is BiCech $sg\beta$ -irresolute and g is BiCech $sg\beta$ -continuous then $g \circ f$ is slightly BiCech $sg\beta$ -continuous.

(iii) If f is BiCech $sg\beta$ -continuous and g is slightly BiCech continuous then $g \circ f$ is slightly BiCech $sg\beta$ -continuous.

Proof: (i) Let U be a BiCech clopen set in Z . Since g is slightly BiCech $sg\beta$ -continuous, $g^{-1}(U)$ is BiCech $sg\beta$ -open in Y . Since f is BiCech $sg\beta$ -irresolute, $f^{-1}(g^{-1}(U))$ is BiCech $sg\beta$ -open in X . Since $g \circ f^{-1}(U) = f^{-1}(g^{-1}(U))$, $g \circ f$ is slightly BiCech $sg\beta$ -continuous.

(ii) Let U be a BiCech clopen set in Z . Since g is BiCech $sg\beta$ - continuous, $g^{-1}(U)$ is BiCech $sg\beta$ -open in Y . Also since f is BiCech $sg\beta$ -irresolute, $f^{-1}(g^{-1}(U))$ is BiCech $sg\beta$ -open in X . Hence $g \circ f$ is slightly BiCech $sg\beta$ -continuous.

(iii) Let U be a BiCech clopen set in Z . Since g is BiCech continuous, $g^{-1}(U)$ is BiCech open in Y . Also since f is BiCech $sg\beta$ -continuous, $f^{-1}(g^{-1}(U))$ is BiCech $sg\beta$ -open in X . Hence $g \circ f$ is slightly BiCech $sg\beta$ –continuous.

REFERENCES

- [1] D. Andrijevic, "On β -open sets", Mat vesnik 48(1996), 59-64.
- [2] I. Arockiarani and J. Martina Jency, " On J-čech closed sets in closure spaces", Kerala Mathematical Association.

- [3] E. Čech, "Topological Spaces", Inter science publications John Wiley and son, New York (1996).
- [4] K. Chandrasekhara Rao and R. Gowri, "On biclosure spaces", Bulletin of pure and applied sciences, 25E (2006), 171-175.
- [5] K. Chandrasekhara Rao and R. Gowri, "Regular generalised closed sets in biclosure spaces", Jr. of institute of mathematics and computer science, (Math. Ser.), 19(3) (2006), 283-286.
- [6] K. Chandrasekhara Rao and R. Gowri "On closure Spaces", Varahamihir journal of Mathematical Science 5(2) (2005), 375-378.
- [7] Chawalit Boonpok, "Generalized Biclosed sets in Biclosure Spaces", Int. Journal of Math. Analysis, Vol. 4, 2010, no. 2, 89-97.
- [8] R. Devi and M. Parimala, "On $\alpha\psi$ -closed sets in BiČech Closure Spaces", Int. Journal of Math. Analysis, Vol. 4, 2010, no. 33, 1599-1606.
- [9] Francina Shalini A and Arockiarani SR I, "New functions in Čech $\pi g\beta$ - closure spaces", International Journal of Recent Scientific Research, Vol. 7, Issue 1, pp. 8515-8517.
- [10] Francina Shalini A and M. Divya Bharathi, "sg β -closed sets in Biclosure spaces", International Journal of Applied Research, Vol. 3, Issue 7, 1188-1192.
- [11] Ganes M. Pandya and C. Janaki and I. Arockiarani, " $\pi g\alpha$ -Separation Axioms in BiČech Spaces", Int Mathematical Forum, Vol. 6, 2011, no. 21, 1045-1052.
- [12] C. Janaki and I. Arockiarani "A new class of open functions in Čech spaces", Acta Scientia India, 36M, 4(2010), 649-653.
- [13] N. Levine, "Generalized closed sets in topology", Rend. Circ. Mat. Palermo, 19(1970), 89-96.
- [12] S. Pious Missierl & J. Arul Jesthi, "Slightly S_g^* -continuous functions and totally $-S_g^*$ continuous functions in Topological spaces", IOSR Journal of Mathematics, Volume 11, Issue 2 Ver. I (Mar - Apr. 2015), PP 27-30
- [13] D. Sasikala and I. Arockiarani, "Čech w – closed sets in closure spaces", International journal of Mathematical Archive – 4(10), 013, 30 – 35.
- [14] J. Slapal, "Closure operations for digital topology", Theoret. Comput. Sci., 305(1-3) (2003), 457-471.



10.22214/IJRASET



45.98



IMPACT FACTOR:
7.129



IMPACT FACTOR:
7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call : 08813907089  (24*7 Support on Whatsapp)